

60072

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Only for formulation into a fungicide for control of diseases of wheat.

Fungicide

Storage and Disposal

Pesticide Storage and Disposal

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Thoroughly clean container before reuse unless this is a Ciba-Geigy refillable container. If this is a Ciba-Geigy refillable container, and is to be refilled, do not rinse or introduce any pesticide other than Dividend MG. Reseal and return the container to an authorized Ciba-Geigy refilling facility. If the Ciba-Geigy container is not to be refilled, triple rinse (or equivalent), and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.


Dividend® trademark of Ciba-Geigy Corporation
U.S. Patent No. 5,266,585

© 1995 Ciba-Geigy Corporation

Ciba Crop Protection
Seed Treatment Products
Ciba-Geigy Corporation
Greensboro, North Carolina 27419

CGA 128L14

[GANNONC.LABELD\LBL-D-MS-WORD]DIV-MG - 1/24/95



6

Reg # 478-121

PM-23

193

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

MAR 28 1995

Kathie J. Tryson
Realex
Division of United Industries Corp.
P.O. Box 15842
St. Louis, MO 63114

Dear Ms. Tryson:

Subject: Revised Labeling
Real-Kill Broadleaf Weed Killer
EPA Registration No. 478-121
Your Submission Dated October 17, 1994

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following provision:

Rewrite the last sentence in the Storage and Disposal block to read "Securely wrap in several layers of newspaper and discard container in trash."

A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the referenced label, incorporating the above change, before releasing the product for shipment.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

DK: 305-7546: FHB/PM23		CONCURRENCES					
SYMBOL	7505C						
SURNAME	D. KENNY						
DATE	3/24/95						

193

293

Real-Kill® Broadleaf Weed Killer

EPA Reg. No. 478-121

AC with 4 In EPA MVR Under the Fungicide, as amended, registered 478



Treats Northern and Southern Grasses

Bluegrass, Fescue, Zoysia, Bermuda, St Augustine, Bentgrass, Centipede, Bahia, Rye

Kills the Weeds, not the Lawn • Kills 33+ Weed Types
Roofs and All • Kills 33+ Weeds

Kills Dandelion, Chickweed, Wild Onion, Plantain, Poison Ivy and many others. Weed details on back

ECONOMICAL CONCENTRATE • WATER-BASED
1 quart treats up to 16,000 square feet

Active Ingredients	
1,1-dimethyl-3-(4-chlorophenoxy)acetic acid	7.59%
1,1-dimethyl-3-(2,4-dichlorophenoxy)acetic acid	3.66%
1,1-dimethyl-3-(2,4,6-trichlorophenoxy)acetic acid	0.84%
Inert Ingredients	87.91%
1,1-dimethyl-3-(4-chlorophenoxy)acetic acid equivalent per gallon at 6.30% *** 0.26 fl. oz.	
1,1-dimethyl-3-(2,4-dichlorophenoxy)acetic acid equivalent per gallon at 3.02% *** 0.369 fl. oz.	
1,1-dimethyl-3-(2,4,6-trichlorophenoxy)acetic acid equivalent per gallon at 0.69%	

Keep Out of Reach of Children

CAUTION

See Back for Additional Precautionary Statements

Net Contents: 32 fl. oz. (1 Quart)

(Optional Claims)

Especially effective on all types of hard-to-kill weeds
Dilutable concentrate • For home lawn care • Kills over 33 weed types including dandelions, chickweed, wild onions, plantain and poison ivy • Kills broadleaf weeds in lawns

Real-Kill Broadleaf Weed Killer controls more than 33 troublesome weeds including Dandelions, Clover, Plantain and Wild Onion. It kills even woody weeds like Poison Ivy and Poison Oak in both Northern and Southern lawns.

This concentrated formula covers up to 16,000 sq. ft. per quart. See specific directions for St. Augustine and other Southern grasses.

WEEDS CONTROLLED: Bedstraw, Black Medic, Buckhorn, Burdock, Chickweed, Chicory, Clover, Dandelion, Dock, Ground Ivy, Heal All, Herbit, Knotweed, Lambsquarters, Lespedeza, Mallow, Morning glory, Oxalis, Pepper grass, Pepperweed, Pigweed, Plantain, Porron Ivy, Poison Oak, Purslane, Ragweed, Sheep Sorrel, Shepherd's Purse, Speedwell, Sours (Spotted) Wild Carrot, Wild Garlic, Wild Lettuce, Wild Onion, Yarrow and other problem broad-leaf weeds

STOP. READ ENTIRE LABEL BEFORE USE

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Shake well before using.

MEASUREMENT CALCULATIONS: Calculate square feet by multiplying length by width. 1 fl. oz. = 2 tablespoons (6 teaspoons). 2 fl. oz. = 4 tablespoons. 4 fl. oz. = 1/2 cup. 8 fl. oz. = 1 cup. 16 fl. oz. = 1 pint. 32 fl. oz. = 1 quart.

WHEN TO USE:

- 1) **SPRING:** The best time to treat weeds is when they are small and growing.
- 2) **FALL:** Many problem weeds germinate in the fall and a treatment then will eliminate many of next year's weeds.

HOW MUCH TO USE:

SPOT TREATMENT

To spray small areas or individual weeds, use a pressure sprayer for better control. Do not use a hose end sprayer. Max 1 fl. oz. (2 Tbsp.) in one gallon of water. Shake well before spraying. Be certain to wet each weed.

ENTIRE LAWN

PRESSURE SPRAYERS (Optional Directions)

Measure amount of concentrate from chart below into sprayer tank and dilute with 1 gallon of water. Close tank sprayer and shake well before spraying. Use

A spray pattern 250 sq. ft.

Grass Type
St. Augustine
Bermuda, E.
Bluegrass

HOSE-END CONCENTRATE

- 1) From each 2
- 2) Fall spr not ad
- 3) Set di
- 4) Attach treated treated

EXAMPLE: to the spray received wh of your dal

Grass
St. / ugut Bahia, Cent
Bermuda, Bentgrass
Bluegrass Rye, Zoysia

CONCENTRATE For sprayer application

- 1) From each
- 2) Pour
- 3) Then spray