

STOP (optional). READ ENTIRE LABEL BEFORE USE.

Directions for Use: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Sprayer Instructions:

How to prime and care for your sprayer:

1. After shaking product well, remove cap from container and replace with sprayer hose and cap assembly. Tighten securely.
2. Turn sprayer nozzle one half turn counter-clockwise. Hold sprayer below the level of container and prime by pumping trigger 10-15 times until product is dispensed. Adjust nozzle to fine spray or jet stream as needed.
3. To store, turn nozzle to "OFF" position and store with sprayer above level of liquid in container to prevent leakage.

TO OPERATE: SHAKE BEFORE USING. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct the spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage. This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations. To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest. **DO NOT TRANSPORT OR STORE UNDER 32° F.** If freezing occurs, thaw and agitate vigorously.

Use on the following crops: Asparagus, Beans, Broccoli, Cauliflower, Cabbage, Brussels Sprouts, Celery, Cranberries, Eggplant, Lettuce, Mustard Greens, Kale, Collards, Turnips, Peppers, Potatoes, Radishes, Spinach and Tomatoes to control: asparagus beetles, aphids, leafhoppers, whiteflies, mealybugs, Mexican bean beetles, 12-spotted cucumber beetles, cabbage loopers, cross-striped cabbageworms, diamondback moth larvae, flea beetles, harlequin bugs, imported cabbageworms, stink bugs, green peach aphids, leafhoppers, fireworms, blister beetles, Colorado potato beetles, and webworms.

Use on the following ornamentals: African Violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wanderer, Yew, Aster, Carnations, Chrysanthemums, Geraniums, Gladiolus and Marigolds. To control: aphids, flea beetles, mealybugs, leafhoppers and whiteflies. To kill gypsy moth caterpillars and adults, adult Japanese beetles

Real-Kill®

ROSE & GARDEN INSECT SPRAY II

WATER-BASED

KILLS GYPSY MOTH CATERPILLARS AND JAPANESE BEETLES • FAST ACTING • AN EFFECTIVE INSECTICIDE FOR USE ON VEGETABLE AND ORNAMENTAL PESTS • CAN BE APPLIED ON DAY OF HARVEST • USE ON DAY OF HARVEST • CAN BE USED ON DELICATE ORNAMENTALS • PROTECTS ROSES, VEGETABLES, HOUSEPLANTS, ORNAMENTALS, TREES, SHRUBS, FLOWERS

Active Ingredients:

Pyrethrins	0.02%
*Piperonyl Butoxide, Technical	0.20%
Inert Ingredients	99.78%

*Equivalent to min. 0.16% (butylcarbityl (6-propylpiperonyl) ether and 0.04% related compounds

Keep Out of Reach of Children

CAUTION

See Back Panel for Additional Precautionary Statements

Net Contents 128 fl. oz.

and Eastern and Forest tent caterpillars on contact: on trees and shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On roses and in flower gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Avoid breathing vapor. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquariums before spraying.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly into lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by disposing of equipment washwaters. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

Physical or Chemical Hazards


Do not use or store near heat or open flame.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or comments? Call toll free 800-332-5553 weekdays 9-5 Central Time.

Manufactured in the U.S.A. for Realex, Division of United Industries, P.O. Box 15842, St. Louis, MO 63114-0842.

EPA Reg. No. 478-125 EPA Est. No. 9688-MO-1
Form: REG/A3 © 1994



292

ALTERNATIVE DIRECTIONS

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO OPERATE: SHAKE BEFORE USING. Turn nozzle to "spray" or "stream" position before using. Hold sprayer 18 to 24 inches from foliage or surfaces being treated. Direct the spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage. This insecticide may be used

indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations. To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest. *Do not transport or store under 32° F.* If freezing occurs, thaw and agitate vigorously.

TREATMENT AREAS	PESTS TO CONTROL
Vegetables: Asparagus, Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Collards, Cranberries, Eggplant, Kale, Lettuce, Mustard Greens, Peppers, Potatoes, Radishes, Spinach, Tomatoes, Turnips	Asparagus Beetles, Aphids, Blister Beetles, Cabbage Loopers, Colorado Potato Beetles, Cross-striped Cabbageworms, Diamondback Moth Larvae, Fireworms, Flea Beetles, Green Peach Aphids, Harlequin Bugs, Imported Cabbageworms, Leafhoppers, Mealybugs, Leafhoppers, Mexican bean beetles, 12-Spotted Cucumber Beetles, Stink Bugs, Webworms, Whiteflies
Ornamentals: African Violets, Asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Dogwood, Geraniums, Gladioli, Marigolds, Roses, Wandering Jew	Aphids, Flea Beetles, Leafhoppers, Mealybugs, Whiteflies
Roses and Flower Gardens Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.	
Trees and Shrubs Treat as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.	Eastern and Forest Tent Caterpillars, Gypsy Moth Caterpillars and Adults, Adult Japanese Beetles

ACCEPTED
17 OCT 1994
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered for the purposes indicated under EPA Reg. No. 448-125

