

**Postmaster Methyl Bromide
w/Chloropicrin Warning Agent
(See under label, Copy A)**

FOR GRAIN FUMIGATION in flat or upright Bulk grain storage
Mills, Warehouses, Box cars, Vaults.

Procedure: For fumigation methods, consult the manual "The Use of
Methyl Bromide for Grain Fumigation", furnished by the Manufacturer.
Consult Manufacturer's Manual for specific information.

Caution: wear a full-face gas mask (plus a canister) when releasing
gas in buildings. Thoroughly aerate buildings after fumigation.

Insects Controlled: All stored grain insects including granary
weevil, rice weevil, lesser grain borers, cadelle, saw-tooth grain
beetle, Angoumois grain moth. Kills rodents.

Rate of Application: For space fumigation of grain the usual rate
is 1 to 3 pounds for each 1000 cu. ft. of space for an exposure of
16 to 24 hours.

Usual rates for bulk fumigation with recirculation:

- Shelled corn ----- 1.5 lbs. per 1000 cu. ft.
- Wheat, barley, rye, oats, rice----- 1.5 lbs. per 1000 cu. ft.
- Grain sorghum, milo ----- 1.5 lbs. per 1000 cu. ft.

Use Precautions: Fumigation at grain moisture content above 14%
may injure germination. Do not repeat fumigation where fumigation
likely to result in trace residues in excess of 10 ppm.

Do not fumigate feed stuffs that are intended for use as feed unless
this label unless it is specifically determined that the fumigation
compliance with Federal Food, Drug, and Cosmetic Act and its
regulation.

Consult Manufacturer's Manual for details of fumigation.

**Keep this label away from children. Do not use if label is missing or
illegible. Place away from live stock.**

Seller makes no warranty, expressed or implied, for use of
this product. Buyer assumes all risk of use of this product, whether
or not in accordance with the label.