

40 Lbs. Net Weight

G E N I E

WEED AND FEED PLUS INSECTICIDE

18--4--5

Guaranteed Analysis		Ingredients
Total Nitrogen (N)	18.00%	Superphosphate, Dolomite,
Available Phosphoric Acid (P ₂ O ₅)	4.00%	Sul-Po-Mag, Urea
Soluble Potash (K ₂ O)	5.00%	

DIRECTIONS:

Spread contents of this bag evenly over 5,000 square feet of lawn (50'x100').
(See spreader settings below.)

1. Do not apply to a new lawn, wait until lawn is three months old and has been mowed four times.
2. Do not mow lawn for two days prior to or two days after application as it is desirable to have maximum leaf surface for an effective kill of weeds.
3. Apply to a DRY lawn, but preferably one that has previously been thoroughly moistened to a depth of two or three inches by rain or sprinkler. Weeds that have been hardened from being in dry soil are not easily killed.
4. Take care to apply evenly with an accurate spreader and correct dial setting. Do not turn corners without shutting off spreader. If your spreader is not shown below, proceed with caution using a number smaller than half the largest number on the dial. For many spreaders with diamond, rectangular or oval holes an opening the size of an ordinary wooden pencil will give an approximate setting.
5. Do not water lawn for 24 hours after application. Should a heavy or prolonged rain occur, it may be necessary to re-apply for good weed control. Resume normal watering procedures after 24 hours.
6. Mouse ear chick weed and Japanese clover occasionally require two applications at the standard rate a month apart. Do not make double applications at one time.

Distributed by:

BAZAR

1818 S. E. 2nd Avenue
Portland, Oregon 97214

ACCEPTED

MAY 3 1969

THE OFFICIAL SEAL OF THE
 FEDERAL BUREAU OF INVESTIGATION
 RECEIVED
 483-153
 TO ATTORNEY GENERAL

Pack
(FRONT OF BAG - CON'T)

Herbicide and Insecticide Analysis

Active Ingredients:

2,4-Dichlorophenoxy acetic Acid.....	0.29%
2-(2-Methyl-4-Chlorophenoxy) Propionic Acid.....	0.19%
Dimethylamine salt of dicamba (3,6-dichloro- <u>o</u> -anisic acid)*.....	0.1144%
Dimethylamine salt of related acids.....	0.01846%
Technical Chlordane**.....	1.45%

*Equivalent to 0.0916% Dicamba

**Equivalent to 0.831% Octachloro -4,7 Methanotetrahydroindane and
0.619% related compounds

Inert Ingredients:

Relative to Herbicide and Insecticide content.....	97.95%
TOTAL.....	100.0000%

C A U T I O N

KEEP OUT OF REACH OF CHILDREN

Induce vomiting if swallowed. Avoid contact with skin or eyes. Wash hands after use.

The contained weed killers are injurious to most plants other than grasses, so avoid applying or allowing to drift to flowers, shrubs or valuable plants. Do not store this product near seeds, plant or other fertilizers or chemicals. Avoid breathing dust. Avoid contaminating feed or food stuffs. Do not use on St. Augustine grass, dichondra, carpet grass or clover lawns. Possible injury to bentgrass. Do not apply within the drip line of desirable trees and shrubs. Rinse spreader with water after use.

RECEIVED
 5-8-69
 483-153
 (continued)

Controls dandelion, plantain and sheep sorrel. Also rosette weeds such as lambsquarter, heal-all, prickly lettuce, shepherds purse, yellow rocket, buttercup and others.

Controls clovers, chickweed, ground ivy, knotweed, henbit, purslane, black medic (Japanese clover) and other vining weeds.

Controls lawn moth larvae, cut worms and root weevil larvae.

Controls ants, earwigs, grasshoppers and certain other lawn and turf insects.

5-8-69
483 153
comment

RITE ANE 8-E



in water as an insecticide.

	74.0%
	21.0%
	5.0%
Total	100.0%


,7-methanotetrahydroindane and

ION

in can cause toxic symptoms. Avoid contact with skin, wash with soap and water. Fish may be harmed. Fish may be killed if product.

H OF CHILDREN

ured by
AL CORPORATION
AGERSTOWN, MARYLAND



ONIONS: Thrips — use $\frac{3}{4}$ quart per 100 gallons of water per acre. Do not apply to green or spring onions.

TOMATOES: Flea beetle, Serpentine leaf miner, Tomato hornworm — use $\frac{1}{2}$ to 1 quart to 100 gallons of water per acre. Repeat as necessary. Do not apply after fruit begins to form.

Soil Applications

WHITE GRUBS: Potatoes, strawberries, sugar beets — use $1\frac{1}{2}$ to $2\frac{1}{2}$ quarts per acre in 100 gallons of water to soil surface as a pre-plant application. Disk into the top 3 to 4 inches of soil immediately.

WIREWORMS: Beans, beets, cauliflower, onions, potatoes, sugar beets, sweet potatoes, turnips — use $1\frac{1}{2}$ to $2\frac{1}{2}$ quarts in 100 gallons of water per acre. Apply to soil surface as a pre-plant application. Disk into top 4 to 6 inches of soil immediately.

Shade Trees and Ornamentals

For Bagworms, Leaf beetles, Elm leaf beetles, Leaf miners, Lace bugs, Tree crickets on Arborvitae, Camelia, Elm, Hawthorn, Holly, Junipers, Magnolia, Maple and Oak — use 1 pint in 100 gallons of water. Begin applications in late spring or when insects first appear. Repeat in 2 to 3 weeks when necessary.

DIRECTIONS

Pour recommended amount of this material into partly filled spray tank. Add remainder of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. If concentrate does not mix readily but tends to separate as an oily layer, do not use as injury to plants may result.

RECOMMENDATIONS


Foliage Applications

BEANS: Flea beetle, Green stinkbugs — use 1 quart in 100 gallons of water per acre. Repeat as necessary. Do not apply on green or snap beans after pods begin to form.

COLE CROPS: Broccoli, Brussels Sprouts, Cabbage, Cauliflower: Cabbage worms, Flea beetles, Green stink bugs, Harlequin bugs — use 1 quart in 100 gallons of water per acre. Repeat as necessary. Make no applications after edible portions begin to form.

EGGPLANT: Flea beetle, Serpentine Leaf miner — use 1 quart in 100 gallons of water per acre. Repeat applications as necessary. Make no applications after fruit begins to form.

**WEED and FEED
PLUS**



PACIFIC

20% BONUS

COVERS

3000

SQUARE FEET

Weeds & Feeds
3000 sq. ft. **\$3.95**

**WEED and FEED
PLUS**

PLUS
PLUS
PLUS
PLUS



**WEED and FEED
PLUS**

12 - 4 - 8

CAUTION

NET WT. 22½ LBS.

PACKED BY
PACIFIC SUPPLY COOPERATIVE

**PACIFIC
WEED and FEED PLUS**

DIRECTIONS

SUGGESTED SPREADER SETTINGS

NOTICE

1987