


*Feeds  
Grass*

*Controls  
Lawn  
Moths*

*Kills  
Weeds*



# Weed & Feed 4-in-1

HERBICIDE/INSECTICIDAL CONTENT  
ACTIVE INGREDIENTS  
[Faded text describing ingredients]

### CAUTION

KEEP OUT OF REACH OF CHILDREN  
SEE BACK PANEL FOR ADDITIONAL  
CAUTION STATEMENTS

EPA Reg. No. 483-160

**25 LBS. NET • COVERS 5,000 SQ. FT.**

**Turf King**

**ACCEPTED  
Weed & Feed**



# Weed & Feed 4-in-1

## WILL NOT BURN IF USED ACCORDING TO DIRECTIONS

WEED and FEED with LAWN MOTH CONTROL beautifies the lawn 4 ways:

1. Controls dandelion, plantain, chickweed and clover with 2,4-D and Silvex.
2. Controls lawn moth larvae, wireworm, chiggers, white grubs, ants, earwigs and ticks with Chlordane.
3. Helps produce a thicker, greener turf.
4. Contains Wetting Agent to increase water penetration into soil.

### GENERAL DIRECTIONS

**APPLICATION:** USE AT RATE OF 1 LB TO 200 SQ FT.

For Best Results

1. Mow lawn to normal height of 2 to 3 days before application
  2. Water lawn thoroughly at least one day before application
  3. Apply when weeds are young and actively growing and in early morning when dew is on the grass. If grass is not moist, sprinkle lightly to hold material on grass and help prevent drift onto plants that might be injured by weed killers
  4. Close spreader when stopping, backing up or turning to avoid damage from over application.
- Do not wash material from weed leaves for 1 to 2 days, after which a thorough watering should be done.

**WEED CONTROL:** To avoid damage to new turf apply after grass is at least 4 to 5 months old. Application may be made any time during the growing season when temperatures will not exceed 80°. Do not apply in rain.

**SOIL PEST CONTROL:** Apply as needed over entire lawn. The exact timing of application will vary by insects. Apply at first sign of infestation.

### GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	16.0%
<small>Derived from Ammonium Sulphate Sterilized Organic Matter and Urea</small>	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	2.0%
<small>Derived from Sterilized Organic Matter</small>	
Soluble Potash (K <sub>2</sub> O)	0.0%

#### ALSO CONTAINS

Iron (Fe)	1.0%
<small>Derived from Sterilized Organic Matter</small>	

Potential Acidity: 723 lbs. Calcium Carbonate per ton.

#### RECOMMENDED FERTILIZER SPREADER SETTING\*

Spread one way, being careful not to skip

Make of Spreader	Rate
TURF KING	3½
SCOTT	5

\*As calibration and general condition of spreader can materially affect the rate of flow neither manufacturer nor seller can guarantee exact coverage as indicated by the above recommended dial settings

CAUTION



ed

UTION

REACH OF CHILDREN  
EL FOR ADDITIONAL  
EMENTS

No. 483 160

~~50. FT.~~  
~~1 FT.~~  
Feed

INSECTICIDE  
AC

Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) . . . . . 2.0

Derived from Sterilized Organic Matter

Soluble Potash (K<sub>2</sub>O) . . . . . 0.0

**ALSO CONTAINS**

Iron (Fe) . . . . . 1.0

Derived from Sterilized Organic Matter

Potential Acidity: 723 lbs. Calcium Carbonate per ton.

**RECOMMENDED FERTILIZER SPREADER SETTING\***

Spread one way, being careful not to skip

Make of Spreader	Rate
TURF KING	3 1/2
SCOTT	5

\*As calibration and general condition of spreader can materially affect the rate of flow, neither manufacturer nor seller can guarantee exact coverage as indicated by the above recommended dial settings.

**CAUTION**

Do not handle this material with bare hands. Avoid inhalation of dust or contact with clothing. If material contacts the eyes, get immediate medical attention. If swallowed, keep children and pets off treated areas until this material has been washed away. The grass is thoroughly dry. **KEEP OUT OF REACH OF CHILDREN**

Apply this product only as specified on this label. Excessive application or repeated treatments are unnecessary. Allow one month between treatments on August grass, Carpet grass, Bentgrass, Dichondra or Clover lawns. Do not apply in temperatures expected above 80° F. This material is injurious to many plants and should not be applied when wind is strong or air is quiet to avoid drift onto valuable plants. Vapors from this product may be injurious to susceptible plants growing in the immediate area. Do not store near fertilizers or fungicides.

**ENVIRONMENTAL NOTE.** This product is toxic to fish and wildlife. Birds and animals that frequent treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply near water. Accidents are likely to occur. Do not contaminate water by cleaning of equipment or disposal.

Clean fertilizer spreader several times with chemical cleaners and rinse thoroughly with water. Do not use spreader for other purposes. Do not reuse or burn container. Dispose of in a sanitary landfill area.

**STORE IN A COOL DRY LOCATION**

**Manufactured by Pacific Supply Cooperative**

PORTLAND OREGON 97208

Oct-21-1968

1765-130

# WINRU

## 20% CHLORDANE CONCENTRATE

### ACTIVE INGREDIENTS 100%

*Technical chlordane .....	20%
Petroleum distillate .....	80%
	100%

\*Equivalent to 12.0% Octachloro-4,7-Methanotetrahydroindane and 8.0% related compounds.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.  
HARMFUL IF SWALLOWED.**

DO NOT USE IN EDIBLE PRODUCTS AREAS OF FOOD PROCESSING PLANTS.

Working surfaces must be protected or cleaned before re-use. See right panel for additional cautions.

NET CONTENTS \_\_\_\_\_ GALLONS

Manufactured by  
**WINRU CHEMICAL & SALES COMPANY**  
923 STATE LINE, KANSAS CITY, MISSOURI 64101

TRANSMITTED TO \_\_\_\_\_

### DIRECTIONS (Continued):

This product should not be applied as a fog or spray mist for space treatment for flying insects, however window and door screens and other areas where such flying insects tend to congregate at rest may be sprayed for their control.

**CAUTION:** This solution if brought into contact with the skin can cause toxic symptoms. Avoid inhalation of the vapors of this product. Avoid contamination of feed or foodstuffs. Do not use on dairy animals or in dairy barns or on forage to be fed to animals being finished for slaughter. Do not use on humans or pets. If spilled, wash with soap and plenty of water.

This insecticide should not be stored, sprayed or otherwise exposed toward or near an open flame.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

# WINRU

## 45% EMULSIFIABLE CHLORDANE CONCENTRATE

### DIRECTIONS:

**Subterranean Termites:** Winru 45% Emulsifiable Chlordane may be used as an aid in the control of Subterranean termites.

**Ground Treatments:** Dilute Winru 45% Emulsifiable Chlordane at the rate of 1 part concentrate to 24 parts water and apply in trench at the rate of one gallon of mixed spray per linear foot. Trench should be 30" deep around foundations, but not below footings. Be sure to treat entire area around buildings. Annual inspection should be made for possible reinfestations. Winru 45% Emulsifiable Chlordane when diluted as directed above will not cause phytotoxicity.

**Roaches:** To control exposed stages apply as a 2% emulsion (5 oz. in 1 gallon of water) spraying thoroughly in all cracks and crevices and other hiding places. Repeat treatment as needed.

**Ants:** Apply as a 2% emulsion directly to ant trails and areas around door sills and window frames.

A water dispersible chlordane concentrate containing 45% technical chlordane.

### ACTIVE INGREDIENTS:

\*Technical Chlordane 45%  
Petroleum Distillate 45%

### INERT INGREDIENTS:

10%  
100%

\*Equivalent to 1.4% of Octachloro-4,7-dichlorobiphenyl and 1.0% of related compounds.

### CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid inhalation and skin contact. In case of spillage on skin wash with soap and water. Avoid contamination of feed and foodstuffs.

This solution if brought into repeated or prolonged contact with the skin can cause toxic symptoms.

This insecticide should not be stored, sprayed or otherwise exposed toward or near an open flame.

See right panel for additional cautions.

U.S.D.A. REG. NO. 1190-34

NET CONTENTS

GALLONS

Manufactured by:  
**WINRU CHEMICAL & SALES COMPANY**  
923 STATE LINE  
KANSAS CITY, MISSOURI 64101

**WINRU**  
**INSECT CONTROL**

### DIRECTIONS (Continued):

**Buildings:** Winru 45% Emulsifiable Chlordane may be applied as a residual spray in houses, and similar buildings for the control of crawling ants, cockroaches, etc. Pay particular attention to sleeping quarters of animals (5 oz. in 1 gallon of water) 4 to 6 weeks or oftener, if ex-

Do not use in dairy barns, milking parlors or bulk food storage buildings.

**CAUTION:**  
Use only according to label.

**NOT FOR HOUSEHOLD USE**  
This product is toxic to fish.

TRADE MARK

# WINRU

## 45% EMULSIFIABLE CHLORDANE CONCENTRATE

REGISTERED

13-1967

IONS:

45% Emulsifiable Chlor-  
in the control of Subter-

Winru 45% Emulsifiable  
concentrate to 24 parts  
the rate of one gallon of  
trench should be 30" deep  
below footings. Be sure to  
buildings. Annual inspection  
reinfestations. Winru 45%  
then diluted as directed  
city.

ed stages apply as a 2%  
(water) spraying thorough-  
s and other hiding places.

sion directly to ant trails  
and window frames.

A water dispersible chlordane concentrate containing 45% technical chlordane.

### ACTIVE INGREDIENTS:

\* Technical Chlordane 45%  
Petroleum Distillate 45%

### INERT INGREDIENTS:

10%  
100%

\*Equivalent to 24% of Octachloro-4,7,8-tricycloheptane and 10% of related compounds.

### CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid inhalation and skin contact. In case of spillage on  
skin wash with soap and water. Avoid contamination to  
feed and foodstuffs.

This solution if brought into repeated or prolonged con-  
tact with the skin can cause toxic symptoms.

This insecticide should not be stored, sprayed or other-  
wise exposed toward or near an open flame.

See right panel for additional cautions.

S.D.A. REG. No. 1700-34

NET CONTENTS GALLONS

Manufactured by:  
**WINRU CHEMICAL & SALES COMPANY**  
923 STATE LINE  
KANSAS CITY, MISSOURI 64101

DIRECTIONS (Continued):

1700-34

**Buildings:** Winru 45% Emulsifiable Chlordane may be  
applied as a residual spray in barns, stables, dog  
houses, and similar buildings. It is effective for con-  
trolling ants, cockroaches, brown dog ticks, and spiders.  
Pay particular attention to cracks in stalls and other  
sleeping quarters of animals. Apply as a 2% solution  
(5 oz. in 1 gallon of water) making applications every  
4 to 6 weeks or oftener if exposed to weathering.

Do not use in dairy barns, milk houses, poultry houses,  
grain or bulk food storage bins.

CAUTION:

Use only according to label directions and limitations.

NOT FOR HOUSEHOLD USE:

This product is toxic to fish and wildlife.