

**DIRECTIONS FOR STORAGE APPLICATION**

Large flat storages, such as quonset huts, should be treated by power equipment in as short a time as possible. Nozzles should be removed so that the fumigant is applied in a stream. Fumigant should be applied over the entire surface of the grain. Make application from the outside of the building if at all possible. If it is necessary to enter the building for application, a gas mask suitable for protection of the operator must be used.

**DOSAGE**

When treating small grains in this type storage, a dosage of from 3 to 4 gallons of fumigant per 1000 bushels should be used. When treating corn in this type of storage, a dosage of from 4 to 5 gallons of fumigant per 1000 bushels should be used.

**DIRECTIONS**

**LARGE GRAIN ELEVATORS**


**Dosage:** It is recommended that the fumigant be applied at the rate of 1 1/2 to 2 gallons per 1000 bushels of grain, except for the first and last 500 bushels entering the bin, where the dosage should be 3 gallons.

**Application:** The fumigant should be applied as it is entering the bin. The proper amount of fumigant to be released into the bin between drafts.

**SALE CONTRACT** \_\_\_\_\_



**BIN-FUME**



**GRAIN FUMIGANT**

Bin-Fume Grain Fumigant is designed to effectively control stored grain insects: Rice weevil, granary weevil, cadelle, confused flour beetle, in grain storage facilities.

<b>ACTIVE INGREDIENTS:</b>	100
Carbon Tetrachloride	92.3
Carbon bisulfide	16.3
Pentane	0.4
Sulphur dioxide	1.0

**WARNING: Keep out of reach of children.**

May be fatal if swallowed or inhaled, or if it comes in contact with the eyes. Avoid contact with the skin, eyes, and clothing. If contact is made, normally, in case of contact with the skin, wash the affected area immediately with soap and water.

U.S. PAT. REG. NO. 2,152,175

MADE IN U.S.A.

**DIRECTIONS FARM STORAGE BINS**

**Dosage:** Grain Temperatures should be above 60 degrees Fahrenheit. The following dosages of fumigant, when properly applied, may be expected to penetrate to the bottom of all ordinary farm bins.

**FOR SMALL GRAINS**

Bin Capacity	Fumigant Per 1000 Bushels
500 - 1500	4 1/2 gallons
1500 - 3000	4 gallons
Over 3000	3 gallons

**FOR CORN**

When treating corn, it is necessary to use higher rates of application as follows:

- 5 gallons per 1000 bushels in steel bins
- 6 gallons per 1000 bushels in wooden bins

**Application:** It is very important and essential that the surface of the grain be leveled and also that all crusts or caked layers be broken up to insure even distribution and proper penetration of the fumigant. To avoid breathing the fumigant vapors, it should be applied from the outside of the bin. This may be done by spraying the fumigant evenly over the entire grain surface. A piece of 2 inch iron pipe without a nozzle may be used for this purpose. The bins should be made as tight as possible to prevent loss of the fumigant.

**Note:** Do not reuse empty drums. Return to drum re-conditioner or destroy by perforating or crushing and burning in a safe place.