

**THIS PRODUCT
FOR USE BY TRAINED PERSONNEL ONLY**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (& DOMESTIC ANIMALS)**

CAUTION

Harmful if swallowed. Avoid breathing vapor. Avoid contact with skin, eyes, or clothing. Wear mask or respirator of type recommended by NIOSH (National Institute for Occupational Safety and Health) to give adequate protection against this material. Wear protective clothing. Wash exposed skin with soap and water after handling and before eating. Wash all contaminated clothing with soap and water, separately from other clothing before re-use.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After spraying in plants, bakeries and other food processing plants, wash with potable water all equipment, benches, shelving, etc., where disposed food will be handled. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry, well ventilated place. Keep container closed.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

FOR CONTAINERS ONE GALLON AND UNDER: Do not reuse, empty container. Wrap container and put in trash.

FOR CONTAINERS OVER ONE GALLON:

METAL CONTAINERS: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and Local authorities.

PLASTIC CONTAINERS: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



PYRE-MIST AQUA 1

**A Concentrated PYRENONE™ Micro Emulsion
—Clear—Stable—Leaves No Oily or Objectionable Residues.
Effective, Fast Acting Insecticide**

**For Use in Homes, Institutions, Restaurants
Food Processing Plants, Dairies, Livestock and Outdoors for Mosquito Control
For Control of Fleas and Brown Dog Ticks on Premises And On Pets
Apply with conventional mechanical or compressed air equipment**

ACTIVE INGREDIENTS

Pyrethrins

* Piperonyl Butoxide, Technical

INERT INGREDIENTS

0.1%

1.0%

96.9%

100%

* Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled — Remove victim to fresh air. Apply respiration if indicated.

If on skin — Wash immediately with soap and water.

If in eyes — Flush eyes for at least 15 minutes with water. Call a physician immediately.

See additional precautionary statements.

EPA REG. NO. 485-59

EPA EST. NO. 3624-MO-1

THE INDUSTRIAL FUMIGANT COMPANY

601 East 159th Street, Olathe, Kansas 66061

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned, with potable water, before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control cockroaches, palmetto bugs, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drug-store beetles, meal worms, grain mites and cadelle. Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish spray bookcases. For ants spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 qt. to 750 sq. ft of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized applications to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths. Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON LIVESTOCK: To protect cattle and horses from horn flies, horse flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.
ON PETS: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

OUTDOOR APPLICATION

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest (low marshy places, stagnant pools or ponds and shore lines). Apply at a rate of 2.5 pints per acre (0.1 ounce per 1000 square feet).

Using truck-mounted conventional misting equipment such as Rotomists models 51C, P1 or 100 HT traveling at 5 m p h to treat a 300 foot swath calibrate equipment to deliver 0.9 gallons per minute.

WARRANTY

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use but neither this warranty nor any other WARRANTY of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE express or implied extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

ACCEPTED

OCT 1 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 485-59

NET CONTENTS _____ GALLONS

616

65 5317

PC

