

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid breathing vapor. Avoid contact with skin, eyes, or clothing. Wear mask or respirator of type recommended by NIOSH (National Institute for Occupational Safety and Health) to give adequate protection against this material. Wear protective clothing. Wash exposed skin with soap and water after handling and before eating. Wash all contaminated clothing with soap and water, separately from other clothing before re-use.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After spraying in plants, bakeries and other food processing plants, wash with potable water all equipment, benches, shelving, etc., where exposed food will be handled. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquarium, before spraying.

ENVIRONMENTAL HAZARDS

Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticides, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: For Containers One Gallon and Under: Do not reuse empty container. Wrap container and put in trash. For Containers Over One Gallon:

- Metal Containers Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and Local authorities
- Plastic Containers Triple rinse (or equivalent) Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Close windows and doors and shut off ventilating systems. Use a non-thermal fogger or vaporizer, adjusted to deliver an aerosol spray (two droplets over 50 micron in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of flies, including fruit flies, mosquitoes, silverfish, flying moths, gnats, crickets, cock-



PYRE-MIST Aqua 5

**A CONCENTRATED PYRETHRIN
MICRO EMULSION**

Clear
Stable

Leaves No Oily or Objectionable Residue

Apply with Conventional Mechanical or compressed air equipment designed and adjusted to deliver a true aerosol mist.

ACTIVE INGREDIENTS:

Pyrethrin.....	0.5%
Piperonyl Butoxide, Tech.....	5.0%
INERT INGREDIENTS.....	94.5%
	100.0%

*Equivalent to 4.0% (butylcarbitol) (6-propylpiperonyl) ether and 1.0% related compounds

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician immediately. Gastric lavage is indicated if material was taken internally. **DO NOT INDUCE** vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled — Remove victim to fresh air. Apply respiration if indicated.

If on skin — Wash immediately with soap and water.

If in eyes — Flush eyes for at least 15 minutes with water. Call a physician immediately.

See additional precautionary statements.

EPA REG. NO. 485-61

EPA EST. NO.

THE INDUSTRIAL FUMIGANT COMPANY

601 East 159th Street, Olathe, Kansas 66061

bees, spiders, scorpions, wasps, hornets, clover mites, cheese mites, cheese shippers and these common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, angoumois grain moths, mediterranean flour moths, indian meal moths, tobacco moths, meal moths and grain mites.

MUSHROOM PRODUCTION AND PROCESSING: To control adult MUSHROOM FLEES, fungus gnats, and cecid flies, in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2000 cubic feet of space.

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of house flies, fruit flies, gnats, mosquitos, skipper flies, wasps, hornets, angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths, and exposed or accessible stages of the crawling insects listed under **SURFACE SPRAYING.** Direct the spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space (one gallon to 20,000 to 40,000 cubic feet of space).

SURFACE SPRAYING: To control cockroaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drug-store beetles, meal worms, grain mites and cadellera. Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers, for silverfish spray bookcases. For ants spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gal. to 750 sq. ft of surface.

To control **carpet beetles:** Spray edges of carpeting and under carpeting and rugs. Make localized applications to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control **bedbugs:** Spray mattresses lightly particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control **flea and brown dog ticks:** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control **clothes moths:** Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests and closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

OUTDOOR APPLICATION

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitos may rest, low marshy places, stagnant pools or ponds and shore lines. Apply at a rate of 13 fluid ounces per acre (0.3 fluid ounces per 1000 square feet).

Using truck-mounted conventional misting equipment traveling at 5 mph to treat 300 feet swath calibrate equipment to deliver 30 fluid ounces per minute.

ACCREDITED
DEC 14 1984
EPA REG. NO. 485-61