SELIG'S SELIG'S

REG. U. S. PAT OFFICE

FOR USE IN MILL MACHINERY KILLS MILL INSECTS

100% Active Ingredients
Ethylene Dichloride and Carbon Tetrachloride
Penetrating — Non-Explosive — Non-Inflammable

Approved by Mutual Fire Prevention Bureau

ACCEPTED How 13,1467

FUNGICIDE AND REPENDENTICIDE ACT
FOR ECONOM : POISON REGISTERED UNDER NO TO SECURE

SELCOFUME is to be used as a fumigant in milling machinery to aid in the control of insect infestation in the machine. SELCOFUME should be used in elevators, conveyors, rolls, purifiers, reels, sifters, spouts, elevator heads and boots, and other milling machinery.

Always start at the bottom of the mill when treating machines, and work up because the vapor of SELCOFUME is heavier than air and the gas therefore drifts down to lower levels. Systematic and regular application once weekly are recommended for controlling insect infestation in machines.

Inject SELCOFUME in purifiers and reels using 16 to 20 ounces in each conveyor; in sifters inject 12 to 14 ounces in each section; in flour conveyors inject 2 to 3 ounces per linear foot; in elevator heads inject 12 to 14 ounces in each leg; in elevator boots inject 12 to 14 ounces; in elevators inject 6 to 8 ounces per floor length. Repeat this treatment as often as necessary. Local fumigation or the treatment of individual machines can be conducted on week-ends or any week night after the mill is shut down. Pound elevator legs so that refuse will fall into the boot, and then clean out the boot. Cleanliness throughout the mill will aid in reducing insect infestation.

When properly applied SELCOFUME controls the Granary Weevil, Lesser Grain Borer, Indian Meal Moth, Angoumois Grain Moth, Red Flour Moth, Saw-toothed Grain Beetle, Flat Grain Beetle, Confused Flour Beetle, Hairy Fungus Beetle, Foreign Grain Beetle, Rice Weevil and Cadelle.

SELCOFUME has an anaesthetic action when breathed in concentrated form.

DANGER! HAZARDOUS LIQUID AND VAPOR. MAY BE FATAL IF INHALED OR SWALLOWED. DO NOT TAKE INTERNALLY. DO NOT BREATHE VAPOR. Do not re-enter treated areas without airline respirator or full-face gas mask with canister for protection against organic vapor. Avoid contact with skin, eyes, and clothing.

DANGER! KEEP OUT OF THE REACH OF CHILDREN

LOS ANGELES LOUISVILLE MIAMI NEW ORLEANS



SAN JUAN
DALLAS
HOUSTON
KANSAS CITY

ATLANTA, GEORGIA