

[491-171]

ACCEPTED

SELIG'S *NOV 5, 1970*

WHITE STAR SPRAY

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE, HERBICIDE, AND
FOR ECONOMIC PEST CONTROL
ED USE
TO ATTACHED COMMENTS.

FOR CONTROL OF HORN FLIES, LICE, FLEAS AND MOSQUITOES ON LIVESTOCK

ACTIVE INGREDIENTS — 100% — Petroleum Distillates, N-Octyl Bicycloheptene Dicarboximide, *Technical Piperonyl Butoxide, Pyrethrins.

*Consists of (butylcarbityl) (6-propylpiperonyl) other and related compounds.

DIRECTIONS:

FOR HORSES, COLTS, MULES, CALVES AND CATTLE: Apply as a mist at the rate of two ounces on large matured animals. Use correspondingly smaller quantities on smaller animals. Do not apply directly to the face, or directly on the udders of milking cows, but otherwise apply the mist thoroughly over the entire body. This spray will kill stable flies, horn flies, house flies, mosquitoes, gnats, black flies and sand flies. For best results, animals should be sprayed daily for a week or ten days.

LICE: Spray livestock with two or three ounces, then brush thoroughly into hair to reach lice next to hide, paying attention particularly to areas liable to infestation. This heavy spraying should be repeated when necessary, but not more than once every three weeks.

FLEAS: Treat livestock same as for lice and in addition, thoroughly spray the bedding and sleeping quarters where the infested animals are housed. Repeat applications, if necessary.

CAUTION

DO NOT APPLY THIS SPRAY NEAR AN OPEN FLAME. IN USING CONVENTIONAL HAND SPRAY, HOLD THE GUN IN NEARLY LEVEL POSITION SO AS TO AVOID PROJECTING A STREAM OF LIQUID ON THE ANIMALS. USING THE GUN IN A LEVEL POSITION WILL INSURE A MORE UNIFORM SPRAY AND BETTER COVERAGE.

HARMFUL IF SWALLOWED. DO NOT WET HIDES OF ANIMALS.

KEEP OUT OF REACH OF CHILDREN

LOS ANGELES
LOUISVILLE
MIAMI
NEW ORLEANS

Manufactured by
THE  CO.
ATLANTA

SAN JUAN
DALLAS
HOUSTON
KANSAS CITY

U.S.D.A. REG. NO. 491-171