[491-182]

SELIG'S

VAPORIZER SPRAY X-T

| ACTIVE INGREDIENTS | 200 |
|--------------------------------------|--------------|
| Pyrethrins | ,3U% 400/ |
| *Tashnisal Dinaranul Rutovide | , ww /t |
| N-Octyl Bicycloheptene Dicarboximide | 1.00% |
| M-OctAl RicAciousbieus ricotpoyiungs | 98.10% |
| Petroleum Distillates | |
| | 100.00% |

Equivalent to 0.48% (butylcarbityl)(6-propylpiperonyl) ether and 0.12% related compounds.

Combats flies, mosquitoes, cockroaches. Close all doors and windows in room while using. For flies and mosquitoes use 1/2 ounces per 1,000 cubic feet of closed room space. For cockroaches, use 3 ounces per 1,000 cubic feet of closed room. Keep room closed for 10 minutes following treatment. Do not remain in treated areas. Ventilate the area after treatment is completed. Use in Fogmaster Fog Generator which dispenses approximately 2 ounces in one minute on the low setting. Be sure that a sufficient amount is used to thoroughly fog the entire room so that the vapor actually envelopes the bodies of the insects to be killed. When used on roaches, direct spray into cracks and crevices for control of these pests. After operating Fogmaster for the required length of time, examine room and if live flies, mosquitoes, and cockroaches still appear, repeat procedure. When finished, sweep dead and stunned insects from the floor and dispose of same. Thorough and repeated spraying of infested area is recommended for control of roaches.

EFFECTIVE AGAINST COMMON PESTS FOUND IN GRAIN STORAGE BUILDINGS

When used as a space spray in grain storage buildings, will kill the exposed stages of grain weevils and flying arain moths: (dry fog — one gallon per 50,000 cubic feet of inner space). For best results, application should always be made in the late evening or early morning. Buildings should be closed as tightly as possible.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN HARMFUL IF SWALLOWED

Do not use, pour, spill or store near heat or open flame. Avoid breathing vapor and contact with skin and eyes. Remove pets and bird cages from rooms being sprayed. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. When using the product in these areas apply only when the facility is not in operation.

Atler.to

Callas

Houston

Louicville



INDUSTRIES

New Orleans

Los Angeles

Puerto Rico

Miami

USDA REG NO 491-182

JAN 28,1973

UNDER THE FEDERAL INSECTIONS
FUNDICIDE AND POSSIFICATIONS
FOR ECONOMIC OF SEGURIERS

0472

į