

[491-212]

ACCEPTED
8-18-71
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT
POISON REGULATION
ED UNDER SUBJECT
TO ATTACHED COMMENTS.

SELIG'S DU-KIL SP

A new synthetic pyrethroid for controlling the following flying and crawling insects: Flies, mosquitoes, gnats, wasps, roaches, earwigs, spiders, grain mites, sowbugs and centipedes.

An effective fast acting insecticide formulated especially for use in food processing and bottling plants. Also ideal for insect control in institutions, public buildings, hospitals, hotels, clubs, restaurants and theatres.

ACTIVE INGREDIENTS:

*†(5-Benzyl-3-furyl) methyl 2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate	0.200% w/w
Related Compounds	0.027%
Aromatic petroleum hydrocarbons	0.265%
Petroleum Distillate	99.500%
INERT INGREDIENTS:	0.008%
	<hr/> 100.000%

*Cis/trans isomers ratio; max 30% (±) cis and min 70% (±) trans. †U.S. Patent No. 3,465,007

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

Harmful if swallowed.

Use as a space spray in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation.

Avoid contamination of food and feedstuff. Cover or remove fish bowls. Remove all pets from area to be treated. Do not use, pour, spill or store near heat or open flame.

Do not reuse empty container. Destroy containers by perforating, crushing and burying in a safe place. Drums may be returned to a reconditioner.

DIRECTIONS

For most effective results use in a good sprayer adjusted to deliver a fine mist.

For houseflies, mosquitoes and gnats: Close all doors and windows. Point nozzle upward, direct spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

Wasp: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

For crawling insects: Roaches, earwigs, grain mites, sowbugs, spiders and centipedes: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insect where possible.

Columbus

Dallas

Houston

Los Angeles



**CHEMICAL
INDUSTRIES**
A DIVISION OF NATIONAL SERVICE LTD. INC.

Atlanta
USDA Reg. No. 491-212
0771

Louisville

Miami

New Orleans

Puerto Rico

NET CONTENTS _____ GAL.