

DIRECTIONS FOR USE

GENERAL: Mix 16 fluid ounces (1 pint) of Ambush in one gallon of water. Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive runoff. Apply to all insect breeding sites and harborages. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

Cockroaches: Thoroughly treat the infested area. Spray baseboards, cracks and crevices around window and door frames; the under surfaces of shelves, drawers and work tables; behind and beneath cabinets, refrigerators, sinks and stoves, in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

Ants: Apply to trails and places where ants enter premises such as around door and window frames. Spray hills and runways outdoors.

Silverfish, Firebrats and Paperlice: Apply to localized areas infested with these insects.

Brown Dog Ticks: Spray sleeping quarters of pets around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not use on animals.

Clover Mites, Crickets, Earwigs, Millipedes, Centipedes, Sawbugs, Scorpions and Spiders: Thoroughly spray the interior and exterior perimeter of infested premises. For Clover Mites, give special attention around window frames.

Flies, Flies, Gnats, Mosquitoes, Pankies and Sandflies (outdoors only): Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, door latches, patios and other areas where they form swarms.

For Use in Meat Packing and Food Processing Plants: Use only in the inedible product areas of meat packing plant. Facilities there must be fully sanitized. All exposed food products, equipment, or packaging materials must be removed, covered and thoroughly cleaned off equipment with water before resuming operations. Treat food processing plants early in the season before opening or other processing begins. Repeat applications, if necessary, when plant is not in operation. Spraying unloading racks and outside areas where insects congregate will aid in preventing re-infestation. Good sanitation and proper disposal practices is essential to any effective insect control program.

Insect Control in Research Facilities: The use of AMBUSH in and around research animal quarters for insect control will not produce changes in the normal enzyme activity in these animals.

SELIG'S AMBUSH

**An Emulsifiable Residual Insecticide Concentrate
Contains Baygon¹
To Be Diluted Before Use
For Industrial and Institutional Use Only**

AMBUSH is a highly effective residual insecticide for control of the insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. This product drives insects from their hiding places, gives rapid kill and residual activity.

ACTIVE INGREDIENTS:

2-o-Isopropoxyphenyl methylcarbamate 9.4%
Petroleum Distillate 60.0%

INERT INGREDIENTS 30.6%

¹U. S. Patent No. 3,111,539

²Baygon is a registered trademark of the German trade mark proprietor, L. E. Kusen.

KILLS: Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Flies, Flies, Gnats, Millipedes, Mosquitoes, Paperlice, Pankies, Sandflies, Scorpions, Silverfish, Sawbugs and Spiders.

WARNING:

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

Columbus

Dallas

Houston

Los Angeles



**CHEMICAL
INDUSTRIES**

Atlanta, Georgia 30336

0675

Louisville

Miami

New Orleans

Puerto Rico

RESTRICTIONS

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near food or food products. Avoid contamination of feed, food, food processing equipment, and food handling surfaces. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants. Avoid breathing spray mist.

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Apply product only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of waste.

Do not reuse empty container. Return drums to reconditioner or destroy by perforating and burying in a safe place.


WARNING:

KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Should illness occur, get prompt medical aid. In physician's office, it is anticipated that the use of Ambush will be limited to the control of insects in and around animal quarters. Do not use, pour, spill or store near open flame.

EPA Reg. No. 491-214



491-214