


491-214

PM-12

1/22/90

181



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS
WARNING**

May be fatal if swallowed, inhaled, or absorbed through the skin. Do not breathe vapor. Do not get in eyes, skin, or on clothing. Do not use as a space spray. Provide adequate ventilation of area being treated. Avoid breathing spray mist. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants. Wash thoroughly with soap and water after handling.

Do not store near feed or food products. Avoid contamination of feed, food, food processing equipment and food handling surfaces.

Do not use in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

ENVIRONMENTAL HAZARDS
This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by disposal. Open dumping is prohibited. Do not re-use empty container.

PESTICIDE DISPOSAL:
Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:
Metal Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

3399

01/90

**SELIG'S
AMBUSH**

**An Emulsifiable Residual Insecticide Concentrate
To Be Diluted Before Use**

ACTIVE INGREDIENTS:	
2-(1-Methylethoxy) phenol methylcarbamate	10.0%
*INERT INGREDIENTS	<u>90.0%</u>
	100.0%

*Contains petroleum distillates

**KEEP OUT OF REACH OF CHILDREN
WARNING**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink one or two glasses of water. Do not induce vomiting or give anything by mouth to an unconscious person.

Tell physician - Atropine sulfate is antidote.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists. Should illness occur, get prompt medical aid.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 491-214-AA

EPA Est. No. 491-GA-1

Net Contents:



640 Selig Drive, S.W. * Atlanta, GA 30336 * (404)691-9220

AMBUSH is a highly effective residual insecticide for control of the insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. This product drives insects from their hiding places, gives rapid kill and residual activity.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

KILLS: Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Flies, Fleas, Gnats, Millipedes, Mosquitoes, Paperlice, Punkles, Sandflies, Scorpions, Silverfish, Sowbugs and Spiders.

GENERAL: Mix 1 1/2 fluid ounces (1 pint) of AMBUSH in one gallon of water. Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborage. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

Cockroaches - Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables; behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

Ants - Apply to trails and places where ants enter premises such as around door and window frames. Spray hills and runways outdoors.

Silverfish, Firebrats and Paperlice - Apply to localized areas infested with these insects.

Brown Dog Ticks - Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

Clover, Mites, Crickets, Earwigs, Millipedes, Centipedes, Sowbugs, Scorpions and Spiders - Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites give special attention around window frames.

Fleas, Flies, Gnats, Mosquitoes, Punkles and Sandflies (outdoors only) - Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

Use in Meat Packing and Feed Processing Plants - Use only in the inedible product areas or meat packing plants, including those that are federally inspected. Any exposed food products, equipment, or packaging materials must be removed or covered before spraying. Rinse all equipment with water before resuming operations. Treat food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestation. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

Insect Control in Research Facilities: The use of AMBUSH in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

WARRANTY STATEMENT

This product is designed exclusively for industrial and institutional use by trained professional maintenance personnel. Label directions and precautions must be followed exactly. Selig Chemical Industries will not be responsible for injury, loss, or damage if product is used in any manner not in compliance with label directions or if precautions are not observed.

Not Reviewed. Registrant claims to be in accordance with the registration of Pesticide Product Guidance for Phase 2 In Response, Page 2.12, Active to Inert Change in Status.