

**USE DIRECTIONS FOR SANITIZING**

*Sanitizing of Food Processing Equipment and other Food Surfaces in Food Contact Locations*

For sanitizing food processing equipment, trays, equipment, food utensils, tables, containers, glassware, etc. use 1 oz. per 4 gallons of water. For sanitizing other hard surfaces.

Apply 1 oz. per 4 gallons of water. Apply to 4 gallons of water.

Apply to milk type equipment, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush. To trays, silverware, glasses, cooking utensils and other similar size food processing equipment, use 1 oz. per 4 gallons of water.

Note: Potable water rinse is required.

At 1 oz. per 4 gallons, SELCOL has a hard water tolerance of 100 ppm of hardness, calculated as CaCO<sub>3</sub>, when evaluated by the AOAC (permeal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*).

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. SELCOL in 4 gallons of warm water.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

**DANGER:**

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

**FIRST AID:**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

**SELCOL DISINFECTANT-SANITIZER  
FUNGICIDE-DEODORIZER**

Disinfectant-Sanitizer-Fungicide-  
Deodorizer for Hospital, Institutional,  
Industrial, School, Dairy and  
Farm Use

AOAC Phenol Coefficients

<i>Staphylococcus aureus</i> , ATCC #6538	95.8
<i>Salmonella typhosa</i> , ATCC #6539	71.9

Active Ingredients

Alkyl (C <sub>14</sub> , 60%; C <sub>16</sub> , 30%; C <sub>12</sub> , 5%; C <sub>18</sub> , 5%) Dimethyl Benzyl Ammonium Chloride	5.0%
---	------

Alkyl (C <sub>12</sub> , 68%; C <sub>14</sub> , 32%) Dimethyl Ethylbenzyl Ammonium Chlorides	5.0%
---	------

Inert Ingredients

EPA Reg. No. 491-222 AA	90.0
EPA EST. 491-GA. 1	100.0

Net Contents

Columbus

Dallas

Houston

Los Angeles



**CHEMICAL  
INDUSTRIES**

Atlanta, Georgia 30336  
0376

**DANGER:**

KEEP OUT OF REACH OF CHILDREN. SEE  
LEFT PANEL FOR FIRST AID AND ADDI-  
TIONAL PRECAUTIONARY STATEMENTS.

**FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY**

**USE DIRECTIONS FOR DISINFECTION**

*Disinfection in Hospitals, Nursing Homes, and Other Health Care Institutions*

For disinfecting floors, walls, bedframes, mounted pictures, tables, chairs, garbage cans, latrine fixtures, and other hard surfaces.

Apply 1 oz. SELCOL to 5 gallons of water. Apply to previously treated hard surface with mop or cloth.

At 1 oz. per 5 gallons use level, SELCOL is effective against *Staphylococcus aureus*.

*Disinfection in Institutions, Industry and Farms*

For disinfecting floors, walls, bedframes, mounted pictures, chairs, garbage cans, latrine fixtures, and other hard surfaces.

Apply 1 oz. SELCOL to 5 gallons of water. Apply to previously treated hard surface with mop or cloth.

At 1 oz. per 5 gallon use level, SELCOL is effective against *Staphylococcus aureus* and *Salmonella typhosa*.

*Fungicide Activity*

At 1 oz. per 5 gallons use level, SELCOL is effective against *Trichophyton interdigitale*, *Candida foot fungus*, and mammalian hard surfaces.

*Disinfection of Water*

For hard water, such as water from a well, hard water, or water that has been treated with hard water, SELCOL can be disinfected by a minimum of 1 oz. per 5 gallons of SELCOL.

*Disinfection of Sewage*

For any sewage, such as from a toilet, sink, or other animal quarters, such as stalls and kennels, and all hard surfaces, use 1 oz. SELCOL per 5 gallons of water.

Apply 1 oz. SELCOL to 5 gallons of water. Apply to previously treated hard surface with mop or cloth.

At 1 oz. per 5 gallons use level, SELCOL is effective against *Staphylococcus aureus* and *Salmonella typhosa*.

For disinfecting floors, walls, bedframes, mounted pictures, tables, chairs, garbage cans, latrine fixtures, and other hard surfaces.

Apply 1 oz. SELCOL to 5 gallons of water. Apply to previously treated hard surface with mop or cloth.

At 1 oz. per 5 gallons use level, SELCOL is effective against *Staphylococcus aureus* and *Salmonella typhosa*.

For disinfecting floors, walls, bedframes, mounted pictures, tables, chairs, garbage cans, latrine fixtures, and other hard surfaces.

Apply 1 oz. SELCOL to 5 gallons of water. Apply to previously treated hard surface with mop or cloth.

411-222