

498-195

11/6/2001

1/7

FRONT PANEL

CHAMPION SPRAYON
FIRE ANT KILLER DUST

- For indoor/outdoor residential use only.
- For household use.
- Household ornamental garden insect control.
- Controls fire ants, fleas, ticks, *termites and more.
- Kills insects for up to 8 months when left undisturbed.
- Does not clump or absorb moisture.
- For use on ornamentals.
- Use on roses, flowers and ornamentals.
- Controls many different insect pests.
- To control imported fire ants on lawns and other residential areas.
- Nonstaining and odorless.
- Long-lasting residual action.
- Long-lasting.
- 8 oz. treats 15 fire ant mounds.
- 16 oz. treats 30 fire ant mounds
- 32 oz. treats 60 fire ant mounds
- Fast, effective kill of fire ants.
- Fast, effective kill of stinging ants.
- Easy to use.
- Convenient.
- Kills fire ants on contact.
- Kills termites

ACCEPTED
 NOV 6 2001
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 498-195

ACTIVE INGREDIENT:

*Deltamethrin [(s)-alpha-cyano-3-phenoxybenzyl - (1R,3R)-3- (2-2 dibromovinyl)-2,
 2-dimethyl cyclopropanecarboxylate] 0.05%

OTHER INGREDIENTS: 99.95%

Total 100.00%

NET WT. XX OZ. (XXX g)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See precautionary statements on back panel.

BACK PANEL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

May be harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water. Use with caution when applying to areas adjacent to any body of water. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Store product in original container in a cool, dry, locked place out of the reach of children.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap in several layers of newspaper and discard in trash.

IMPORTANT: Read the entire Directions for Use, Precautions and Disclaimer before using this product. This product is a ready-to-use insecticide dust, which exhibits effective knockdown and long lasting control against the pests listed when used according to label directions. This product can be used for long-lasting pest control in and around homes and their immediate surroundings. Permitted areas of use include but are not limited to:

- Apartments
- Houses
- Household greenhouses
- Animal quarters
- Mobile homes and recreational vehicles
- Ornaments

This product is intended for applications with a hand duster or from the shaker can, to hiding and

runway areas and other places where insects are found.

INDOOR USE

In living areas, make applications in such a manner as to avoid deposits on exposed surfaces or introducing the material into the air.

To apply insecticide directly into cracks and crevices, use a bulbous duster or other suitable equipment. Apply lightly and uniformly to infested area. Pay particular attention to cracks and crevices; service ducts; false floors and ceilings; wall voids; around electrical and telephone fittings and equipment; around water and sewer pipes; under and behind cabinets, refrigerators and sinks; around windows and door frames; along baseboards; in attics and crawl spaces. The amount to be applied will vary with the site but usually be in the range of 2-3 grams of product per square yard (or 0.5 lbs per 1,000 square feet).

ANTS, BOX ELDER BUGS, COCKROACHES (INCLUDING BOTH RESISTANT AND NON-RESISTANT STRAINS), CRICKETS, FLEAS, FIREBRATS AND SILVERFISH: Apply dust thoroughly to all areas where these pests crawl and hide, especially in those areas described above. For ***ANTS***, apply to ant trails, around doors and windows and other places where ants enter premises.

****TERMITES, CARPENTER ANTS AND CARPENTER BEES:*** Apply to voids or channels in damaged wooden members of a structure; to cracks, spaces or bearing joints between wooden members of a structure; and in locations vulnerable to termite attack. Drilling may be necessary prior to applying dust in these locations, but it must be recognized that such drilling and treating of lumber in a particular location may not result in complete coverage of infested and vulnerable voids or galleries. This dust can be applied in attics and crawl spaces, to prevent entry of drywood and Formosan termites, at a rate of 0.5 lbs per 1,000 square feet of floor area. In these treatments, all vents and blowers should be closed. Inspect at 6 months and 1 year to ensure that colony has been killed.

The purpose of such applications of the product is to kill workers or winged forms, which may be present in the treated channels at the time of treatment and to impart a temporary resistance to attack at these places, which may aid in the prevention of immediate reinfestation. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment, but merely a supplement. For active termite infestations, get a professional inspection.

CENTIPEDES, GROUND BEETLES, MILLIPEDES, SCORPIONS, SOWBUGS AND SPIDERS: Apply dust along and behind baseboards, to window and door frames, corners, pipes, storage locations, attics, crawl spaces, and other areas where these pests may enter or crawl.

WASPS AND BEES: It is generally advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Wear protective clothing if deemed necessary to avoid stings. Using hand or powder duster (actisol or similar equipment) or other suitable means, with extension tubes if necessary, thoroughly dust nest, entrance and surrounding areas where insects alight. Nests in wall voids can be located by listening with your ear against the wall. Drill a hole in the area, blow dust in and seal. For ground wasps, dust entrance and surrounding areas. For best results, check nests carefully 1 or 2 days after treatment to ensure complete kill, then remove and destroy nest to prevent emergence of newly hatched insects. If removal is not feasible, retreat nest if necessary. Do not

use this product to treat overhead nests.

WEBBING CLOTHES MOTHS: Use this product as a crack and crevice treatment in closets and other storage areas where these pests are found. This is an adjunct treatment and will not control moth larvae already on the clothes. **DO NOT** apply this product directly to clothing.

CARPET BEETLES (BLACK, FURNITURE, COMMON AND VARIED): Dust where these insects are found, for example: on and under edges of floor coverings, around baseboards, under rugs and furniture, and in closets.

BEDBUGS: Apply dust thoroughly to all areas where these pests are typically found, including folds and tufts of mattresses, coils of springs, cracks and hollow posts of bedsteads, upholstery of chairs and sofas, picture frame moldings. All cracks and crevices in room should be treated. Allow powder to remain in contact with surface for 4-6 hours. Thoroughly clean linens on bed.

STORED PRODUCT PESTS: To aid in the control of exposed stages of:

- | | |
|-------------------------------|----------------------------------|
| CADELLES | MERCHANT GRAIN BEETLES |
| CHEESE MITES | RED FLOUR BEETLES |
| CIGARETTE BEETLES | RICE WEEVILS |
| CONFUSED FLOUR BEETLES | RUSTY GRAIN BEETLES |
| DARK MEAL WORMS | SAW-TOOTHED GRAIN BEETLES |
| DRUGSTORE BEETLES | SPIDER BEETLES |
| GRAIN MITES | TOBACCO MOTHS |
| GRANARY WEEVILS | YELLOW MEAL WORMS |
| LESSER GRAIN BORERS | |

In the home, all food-processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Clean up waste materials, dust, dirt and other debris. Thoroughly treat floors, walls and other surfaces in storage and handling areas. Apply dust in cracks and crevices, behind and under cupboards, and in other difficult-to-reach areas where insects may hide. **DO NOT** apply directly on food, utensils or other areas where food is stored.

ORNAMENTAL PLANT PESTS: For the control of:

- | | |
|----------------------------|--------------------------|
| APHIDS | LEAF ROLLERS |
| BEET ARMYWORMS | ORANGE TORTRIX |
| BLISTER BEETLES | PERIODICAL CICADA |
| BOX ELDER BUGS | PLANT BUGS |
| CERTAIN LEAF MINORS | ROSE CHAFER |
| (BOXWOOD, HOLLY | ROSE MIDGES |
| OAK, SPRUCE NEEDLE) | ROSE SLUGS |
| ELM LEAF BEETLES | TENT CATEPILLARS |
| FLEA BEETLES | THRIPS |

**JAPANESE BEETLES
LEAFHOPPERS**

WEEVILS

SHRUBS AND ORNAMENTALS: Apply when insects or their damage first appear and repeat at weekly intervals or as necessary. Apply 8 ounces per 1,000 square feet, or thoroughly apply to place a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

ROSE, PETUNIAS, CHRYSANTHEMUMS, GERANIUMS, CARNATIONS AND OTHER RELATED FLOWERS: To control many leaf- and flower-chewing insects, apply as soon as the insect appears and repeat at 7-day intervals or as long as insects continue to appear. Apply a thin layer of dust on both upper and lower sides of leaves, stems and flowers.

OUTDOOR USE

PERIMETER TREATMENT: Use this product as a residual treatment for control of **ANTS, BEES, CENTIPEDES, COCKROACHES, CRICKETS, FIREBRATS, FLEAS, GROUND BEETLES, MILLIPEDES, SCORPIONS, SILVERFISH, SLUGS, SOW BUGS, SPIDERS, TICKS AND WASPS.** Make a residual treatment around windows and doors, porches, screens, eaves, patios, garages, under stairways, in crawl spaces where pests are active and may find entrance.

FLEAS AND TICKS IN RODENT BURROWS: To control fleas and ticks in and around rodent burrows around homes and other buildings, apply dust with a shaker can. Thoroughly apply dust to adequately cover mound and around the perimeter (18" from the outer edge). Be sure to treat abandoned rodent burrows which are suspected of containing fleas and ticks. Treat for a minimum of 5 days for light rodent populations, 10 days when populations are dense. Apply as needed for control.

USE DIRECTIONS FOR FIRE ANT TREATMENT

IMPORTED FIRE ANT MOUND TREATMENT: Apply this product with a shaker can. Thoroughly apply dust to adequately cover mound and around the perimeter (18" from the outer edge of the mound). A second treatment may be necessary if complete control is not obtained. **DO NOT DISTURB THE MOUND PRIOR TO APPLICATION.** If possible, treat all colonies in the vicinity. Also, treat colonies which have not yet constructed a mound.

[Alternate directions: Evenly sprinkle one tablespoon (approximately 1/2 oz.) of this product over the top of each mound. Do not water in or disturb the mound. Apply when temperatures are between 65° F and 80° F. Allow 3 to 4 days for maximum control, then repeat as necessary. Food utensils such as a teaspoon or measuring cup should not be used for food after use with pesticides.]

THE USER OF THIS PRODUCT ASSUMES ALL RISKS OF USE, STORAGE AND HANDLING NOT IN ACCORDANCE WITH ITS DIRECTIONS AND CAUTIONS.

Questions about this product? Call (708) 865-1000.
MEDICAL INFORMATION ONLY:
Chem-Tel, Inc. (800) 255-3924

[OPTIONAL RECYCLE SYMBOL]

7/7

[OPTIONAL MADE IN USA GRAPHIC]

[OPTIONAL UPC SYMBOL]

CHASE PRODUCTS CO.
Putting the best at your fingertips.™
P.O. Box 70 • Maywood, IL 60153
www.chaseproducts.com

8-XXXX

BOTTOM OF LABEL
