Eest Available COPY

HIN 7 1976

5-27-76 6-2-76

Whitnire Research Labs, Inc. 3568 Tree Court Ind. Blvd. St. Louis, MO 63122

Gentlement

Subject :

WEITMIRE BARN FOGGER EPA Reg. No. 499-128 Application of May 21, 1976

The labeling referred to shove, submitted in connection with registration under the Pederal Insecticide, Purgicide, and Rodenticide Act, as emended, is acceptable, and a stamped copy is enclosed for your records.

The alternative formulation submitted on the above is acceptable.

The product name change is also acceptable.

Sincerely, the structure states

Charles T. Mitchell maker Mark Product Manager 17 Insecticide Rodenticide Branch Registration Division WH-567

a in things as set a Enclosure : under ; cosarde. Do not

WH-567:CTMitchell:cq:rm 219: WSHB Retyped: CTMitchell: pl: WENE, me 229

ACTIN

. -14 <sup>°</sup> ae de ann Repairs for politika an arguna ant sature lappy

Ferenzia and the matters of during treatment. Adda dig stations statilities s ing transies or the beiner ustral. When us nea tu a

SPECIFICATIONS. Bark Print, Gold, 6752 - ---Bergeles Longe Nora, Child, C. S.

- **N** 

### DIRECTIONS

barn fogge

**BARN USE** — Walk behind animals and direct fog over backs allowing approximately 1 second per animal. For most effective results, apply each morning.

**ANIMAL USE** — From approximately about 2-feet distance, thoroughly fog entire animal as it is being released to pasture. Spot-treat withers, shoulders and back where saliva accumulates from head tossing.

MILKING PARLOR AND MILK ROOM —Close all windows and doors. Fog at the rate of 1 to 2 seconds per 1000 cubic feet. If rooms cannot be closed tight, double the dosage.

WARNING:

Cover milking utensils and milk to prevent contamination from fog and dead or falling insects.

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator. Avoid contact with skin.



Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation. Avoid breathing of vapor. PRODUCT CODE: 05-0250

Stand in middle of room. Direct for

upward and fog in all directions.

Seconds may be counted by saying

one-thousand, two-thousand, etc. Keep room closed for 15 minutes af-

ter treatment. Example: A room 20feet long, 12-feet wide and 8-feet

high requires 2 seconds if closed

**FOR HOME USE** — To kill roaches, spiders, silverfish, wasps, thoroughly fog insects and all possible hiding places such as cracks, baseboards,

sinks, cabinets and shelves. Repeat

Remove pets, birds and cover fish

tight; 4 seconds if not.

treatment when necessary.

aquariums before fogging.

Loui

St.

ries,

Se a

9

White

à

Manufactured

## SPECIFICATIONS

Flash Point: None. CSMA flammability test method. Explosive Point: None. CSMA explosive test method. Dielectric: 30.

# KILLS HOUSE FLIES, HORN FLIES, STABLE FLIES, GNATS AND ROACHES

Whitmire `

# FOR ANIMALS AND MILK HOUSES

#### ----

 ACTIVE
 Pyrethrins I & II
 .500%

 \*Technical Piperonyl Butoxide
 4.000%

 Refined Petroleum Oil
 8.000%

 INERT
 87.500%

INGREDIENTS

\*Equivalent to 3.2% (Butylcarbityl) (6-Propylpiperonyl) Ether and .8% of Related Compounds.

> E.P.A. Reg. No. 499-128AA E.P.A. Est. No. 499-MO-1

NET WEIGHT 2<sup>1</sup>/<sub>2</sub> LBS.

WARNING: Keep out of reach of children, see side panel for additional warnings.