

PM 17 499-145 1/1 1-30-71

Rev. 8/74 Label #18-2590

### SOME WAYS THAT INSECTS ARE ATTRACTED

Flying insects have odor, heat and light sensors on their bodies. These act like radar and guide them to their food, warmth and shelter. Spilled food, garbage cans, rooking give off odors (gas) that insects lock on to and follow a great distance to the source.

### SANITATION

This is the first step and most important of any insect control program. Clean up all spilled foods, garbage, etc. Clean oil surfaces. Scrub garbage cans so they do not give off odors (gas) that attracts insects. Remove pets, birds and cover fish aquariums before spraying.

### DIRECTIONS

**FLIES, MOSQUITOES, GNATS, FLYING ADULT MOTHS:** Obtain cubic feet of space to be treated. Example: 10' high x 10' long and 10' wide = 1000 cubic feet. Close doors and windows. Spray from 1 to 4 seconds per 1000 cubic feet. Disperse in all directions. If possible, have fans operating to speed up circulation of aerosol fog throughout area being treated. Keep spray closed for 15 minutes after treatment.  
**GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, CARPET BEETLES, DRUG STORE BEETLES** in egg, pupa and larval stages: Contact as many insects as possible when treating infested areas such as cracks, crevices, stored products, machinery, and dark secluded areas. For Carpet Beetles treat infested area of carpets, localized areas of floors, baseboards, and into cracks and under carpets where these pests may be found. Spray 10 to 20 seconds per 1000 square feet of floor space. Repeat treatment weekly or as insects appear.

### SPECIFICATIONS

Flash Point: None. CMSA flammability test method.  
Explosive Point: None. CMSA explosive test method.  
Dielectric: 30  
Edible products areas of federally inspected poultry, meat, and egg packing plants must be limited to the space spray treatment only.  
The premises should be cleaned by thorough washing after spraying when the product is applied in federally inspected poultry, meat, and egg packing plants.

### WARNING:

- Contains under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.
- Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment.
- Thoroughly clean before using. When using the product in food areas, apply only when the facility is not in operation.
- Do not remain in treated areas. Ventilate the area after treatment is completed.

**ACCEPTED**  
OCT 18 1974  
UNDER FEDERAL PESTICIDE  
FUNCTIONS AND MONITORING ACT  
FOR ECONOMIC POISON REGISTER  
EPA UNDER NO. 499-145-101 SUBJECT  
TO ATTACHED COMMENTS.

Net Weight 2 1/2 lbs

# ME Insect Fogger

Containing Pyrethrum and Piperonyl Butoxide  
E.P.A. Reg. No. 499-145-AA-16101  
E.P.A. Est. No. 499-MO-1  
ACTIVE: 50%  
Pyrethrum 1 & 11 500%  
Technical Piperonyl Butoxide 4.000%

**INERT** 95.50%  
Equivalent to 3.2% (Butylcarbitol) (Propylparathion) Ether, and .8% of Related Compounds.  
Heavy Duty, High Pressure, Concentrated Insecticide For Use Inside Of Drive-In Food Service Locations, Food Plants, Hospitals, Restaurants, Supermarkets, Motels.

**WARNING:** Keep out of reach of children. See side panel for additional cautions.

## MB CODE 5025

2 1/2 lbs. Net Wt.  
**BEST AVAILABLE COPY**

Manufactured By  
Whitmore Research Laboratories, Inc.  
3568 Tree Court Industrial Blvd.