


[499-182]
[499-182]

Whitmire
FLY FOGGER



FLY FOGGER

When Used According To Directions:
Flash Point: None. CSMA flammability test method.
Explosive Point: None. CSMA explosive test method.
Dielectric: 30.

Dosage required 1 sec. per 1000 cu. ft.
Treatment per unit 1,135,000 cu. ft.
Cost of treatments Approx. 1/2¢ per 1000 cu. ft.
Delivery rate @ 70°F 1 gram per sec.
Cylinder Aluminum
Propellent Gas High pressure

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F. may cause bursting. Never throw container into fire or incinerator.

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation. Avoid inhalation. Avoid skin contact. Do not remain in treated areas. Ventilate the area after treatment is completed.

Remove pets, birds and cover fish aquariums before spraying.

INGREDIENTS

ACTIVE:
Pyrethrins500%
*Technical Piperonyl Butoxide 1.000%
N-octyl bicycloheptene dicarboximide 1.000%
Refined Petroleum Oil 8.000%
INERT: 89.500%
*Equivalent to 0.8% (Butylcarbityl) (6-Propylpiperonyl) Ether and 0.2% of Related Compounds.

A heavy duty, commercial formula insecticide for use in hospitals, restaurants, food plants, supermarkets, motels, homes, hotels, utilities.

E.P.A. REG. NO. 499-182 AA
E.P.A. EST. NO. 439-MO-1

Product Code: 02-0565
Manufactured by
WHITMIRE RESEARCH LABORATORIES, INC.
St. Louis, Missouri 63122

CAUTION: Keep out of reach of children.
See side panel for additional cautions.

This is the first step and most important of any insect control program. Clean up all spilled foods, garbage, etc. Clean all surfaces. Scrub garbage cans so they do not give off odors (gas) that attract insects.

FLYING INSECTS: Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths, Fruit Flies. Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disperse in all locations. Keep area closed for 15 minutes.

MUD DAUBER AND WASP NESTS: Spray mist directly on insects and their nests from approximately 2 foot distance. These bees contacted will fly away from fog.

GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, CARPET BEETLES, DRUG STORE BEETLES in exposed adult and larvae stages: Contact as many insects as possible when treating infested areas such as cracks, crevices, stored products, machinery, and dark secluded areas. For Carpet Beetles, treat infested area of carpets, localized areas of floors, baseboards, and into cracks and under carpets where these pests may be found. Spray 10 to 20 seconds per 1000 square feet of floor space. Repeat treatment weekly or as insects appear.

FLEAS in buildings: Close windows and doors. Hold aerosol 36" above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 square feet, making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatments after 7 days.

ROACHES, SPIDERS, WATERBUGS, ANTS, SILVERFISH, CRICKETS: Hold aerosol close to cracks and mist directly into them, contacting as many insects as possible. Apply freely in all possible hiding places behind baseboards, sinks, cabinets, coffee makers, meter boxes, manholes. Repeat all treatments as necessary.