
PT' 565F Pyrethrum

Insect Fogger With Dual Activators

SANITATION

This is the first step and most important of any insect control program.

Clean up all spilled foods, garbage. Clean all surfaces. Scrub garbage cans so they do not give off odors (gases) that attract insects.



Set is a set of the passence of A. Aug. Wa. 499-200



PT 565F Pyrethrum

DIRECTIONS

The insects: Houseflies. Mosquitoes. Gnats. Wasps. Shiau Flying Worths: Truit fless close at Alphows and doors and direct top upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disease in all locations. Keep area closed for 15 minutes.

And Douber and Wasp Nests: Spray mist directly on insects and their nests from approximately 2 and distance. These bees contacted will fly away from fog.

Figure Stages of Indian Meal Moth, Chocolate Moth, Mediterranean Flour Moth, Angoumois Grain Moth, Close all doors and windows. Onect spray into all parts of room especially around Stored Direction Containers, pallets and darkened areas. Spray 5 to 10 seconds for each 1000 clopic feetor. Spray 5 to 10 seconds for each 1000 clopic feetor. Spray 5 to 10 seconds for each 1000 clopic feetor. Spray 5 to 10 seconds from the second product should be fumigated or treated by other affective methods. To sing an approved product intended for this purpose.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

PRECAUTIONARY STATEMENTS CAUTION Hazards to Humans and Domestic Animals

Avoid inhalation. Avoid skin contact. Do not remain in treated areas Ventilate the area after the remain is completed. Remove pets. birds. and cover fish aquariting before spraying.

Food Processing Areas Of Food Plants: Feeds should be removed or Covered during treatment. All solves processing surfaces should be covered during treatment or thoroughly cleaned before using the product in these areas, apply only when the facility is not in operation.

Physical or Chemical Hazards

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure that in the property of the pr

Whitmire Research Laboratories, Inc., St. Louis, MO 63122

