

Proposed Draft Label

7/18/83

Front Panel:

499-208

Whitmire

# PT<sup>®</sup> 565 Pyrethrum

## Insect Fogger

Manufactured by Whitmire Research Laboratories, Inc., St. Louis, MO 63122  
Only for Sale to, Use, and Storage by Servicepersons.

A heavy duty, commercial formula insecticide for use in hospitals, restaurants, food plants, supermarkets, motels, hotels, utilities, homes, warehouses and other commercial buildings, and agricultural operations such as dairy farms, poultry, swine and beef cattle operations, and horse barns and stables

**ACTIVE INGREDIENTS:**

Pyrethrins	.....	.500%
* Piperonyl Butoxide, Technical	.....	1.000%
N-octyl bicycloheptene dicarboximide	.....	1.000%
Refined Petroleum Oil	.....	8.000%

**INERT INGREDIENTS:** ..... 99.500%

\* Equivalent to 0.8% (Butylcarbyl) & (Propylperonyl) Ether and 0.2% of Related Compounds

E.P.A. REG. NO. 499-182 AA  
E.P.A. EST. NO. 499-MO-1

A heavy duty, commercial formula insecticide for use in hospitals, restaurants, food plants, supermarkets, motels, homes, hotels, utilities

KEEP OUT OF REACH OF CHILDREN.

**CAUTION:**

See side panel for additional  
precautionary statements.

NET WEIGHT: 15 LBS.

400 208

Side Panel:



### PT<sup>®</sup> 565 Pyrethrum Insect Fogger

#### SANITATION

This is the first step and most important of any insect control program. Clean up all spilled foods, garbage. Clean all surfaces. Scrub garbage cans so they do not give off odors (gases) that attract insects.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL — To be 12 pt. type

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in a cool area away from heat or open flame.

Pesticide Disposal: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Container Disposal: Replace cap and discard empty container in trash, sanitary landfill, or by other approved state and local procedures. Do not incinerate or puncture.

**FLYING INSECTS:** Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths, Fruit Flies. Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disperse in all locations. Keep area closed for 15 minutes.

**MUD DAUBER AND BIRDP NESTS:** Spray mist directly on insects and their nests from approximately 2 foot distance. These bees contacted will fly away from fog.

**GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, CARPET BEETLES, DRUG STORE BEETLES** in exposed adult and larval stages. Contact as many insects as possible when treating infested areas such as cracks, crevices, stored products, machinery and dark secluded areas. For Carpet Beetles, treat infested area of carpets, localized areas of floors, baseboards, and into cracks and under carpets where these pests may be found. Spray 10 to 20 seconds per 1000 square feet of floor space. Repeat treatment weekly or as insects appear.

**FLEAS** in buildings. Close windows and doors. Hold aerosol 36" above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 square feet, making sure that all floor, side, and chair surfaces are contacted. Repeat treatments after 7 days.

**ROACHES, SPIDERS, WATERBUGS, ANTS, SILVERFISH, CRICKETS:** Hold aerosol close to cracks and nest crevices to them, contacting as many insects as possible. Apply freely to all possible hiding places behind baseboards, sink cabinets, coffee makers, meter boxes, mailboxes. Repeat all treatments as necessary.

#### DIRECTIONS FOR ADULT FLY CONTROL

##### IN AGRICULTURAL OPERATIONS:

**Dairy Farm Use - MILKING PARLOR AND MILK ROOM:** Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. If rooms cannot be closed tight, double the dosage. Stand in middle of room. Direct fog upward and spray in all directions. Seconds may be counted by saying one thousand, two thousand, etc. Keep room closed for 15 minutes after treatment. Example: A room 20 feet long, 12 feet wide, and 8 feet high requires 2 seconds if closed tight, 4 seconds if not. Do not remain in treated areas. Vaccinate the area after treatment is completed. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

**ANIMAL USE:** From approximately 2 foot distance, thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where saliva accumulates from head tossing.

**STANCHION BARN USE:** Walk behind animals and direct spray over backs allowing approximately 1 second per animal. For most effective results, apply each morning.

**Poultry Operations:** Use only in empty poultry houses (when birds are not present). Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.

**Hog Operations:** In hog houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per hog. Repeat application daily or as necessary.

**Horse Barns and Stables:** In horse and pony barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application, lightly mist over the backs of horses and ponies from a distance of at least 2 feet. Do not spray directly toward horse's face or eyes. Repeat application daily or as necessary.

**Beef Cattle Operations:** In barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal use, thoroughly spray entire animal as it is being released to pasture. Repeat application daily or as necessary.

#### CRACK AND CREVICE APPLICATION

For direct spray into cracks and crevices, use Whitmore Crack and Crevice Equipment.

#### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

BEST DOCUMENT AVAILABLE

499-208

**Side Panel (continued):**

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. When using this product as a space spray in food areas, do not apply while food processing is under way. Except in federally inspected meat and poultry plants, careful crack and crevice application for inspection purposes is permissible at all times. After space spraying in meat packing plants, bakeries, and other food processing plants, wash with potable water all equipment, benches, shelves, etc. where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

In hospitals, remove patients from rooms prior to treatment. Keep rooms vacated for minimum of two hours and ventilate before patients are returned.

**Environmental Hazards**

This product is toxic to fish and birds. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Avoid the product spray as specified on the label.

**Physical or Chemical Hazards**

Contents under pressure. Do not puncture or compress container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers, etc.