

**PRECAUTIONARY STATEMENTS
CAUTION**

**Hazards to Humans and
Domestic Animals**

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. When using this product as a space spray in food areas, do not apply while food processing is under way. Except in federally inspected meat and poultry plants, careful crack and crevice application for inspection purposes is permissible at all times. After space spraying in meat packing plants, bakeries, and other food processing plants, wash with potable water all equipment, benches, shelves, etc. where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

Environmental Hazards

This product is toxic to fish and birds. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Physical or Chemical Hazards

Contains under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 120°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces.

USE DIRECTIONS

Read label of Federal Law to use this product in a manner inconsistent with the label.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal.

Store in original container in a cool area away from heat or open flame. Recycle wherever possible. Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State, and local laws.

CONTAINER: Recycle, reuse, and discard empty container in accordance with applicable Federal, State, and local regulations.

AGRICULTURAL USES: For control of house flies, stable flies, horn flies, mosquitoes, and other biting insects.

GENERAL USES: In barns, milking parlors, and other areas where livestock are housed, spray at the rate of 1 to 2 seconds per 1000 cubic feet. In hog houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. In poultry houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. In horse barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. In feed storage areas, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. In food processing plants, wash with potable water all equipment, benches, shelves, etc. where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

ANIMAL USES: From appropriate facilities, thoroughly spray entire animal as being released to pasture. Do not spray directly toward animal's eyes, about 1 foot behind shoulders, and back where some difficulties from head tossing.

STANCHION BARN USE: Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

(Continued)



**FLYS
-OFF.**

Farm Insect Fogger

WITH DUAL ACTIVATORS

FOR: Dairy Farms (including Milk House, Milking Parlor, Loading Sheds, and Holding Lot), Poultry and Hog Operations, Horse Barns and Stables, Beef Cattle Operations, Kennels, Catteries, Food Storage Areas, and Meat Packing and Food Processing Plants.

KILLS: House Flies, Horn Flies, Stable Flies, Gnats, Mosquitoes, Wasps, Small Flying Moths, Fruit Flies, Adult Fleas, Grain Weevils, Flour Beetles, Chocolate Moths, Carpet Beetles, Drug Store Beetles, Roaches (Waterbugs), Spiders, Ants, Silverfish, Crickets.

ACTIVE INGREDIENTS	
Pyrethrins	0.500%
*Technical Piperonyl Butoxide	1.000%
N-octyl bicycloheptene dicarboximide	1.000%
Refined Petroleum Oil	8.000%
INERT INGREDIENTS	
89.500%	
*Equivalent to 0.8% (Butylarbityl) to Propylpiperonyl Ether and 0.2% of Related Compounds	

EPA Reg. NO. 499-208-AA • EPA Est. No. 499-MO-1



**KEEP OUT OF REACH OF CHILDREN.
CAUTION:**

See side panel for additional precautions.

USE DIRECTIONS (Continued)

Poultry Operations: Use only in empty poultry houses when birds are not present. Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.

Hog Operations: In hog houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

Horse Biting and Biting: In horse and pony barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application, lightly mist over the backs of horses and ponies from a distance of at least 2 feet. Do not spray directly towards horse's face or eyes. Repeat application daily or as necessary.

Beef Cattle Operations: In barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal use, thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

NON-AGRICULTURAL USES

Kennels & Catteries: Remove animals before spraying. Ventilate after treatment is completed.

FLYING INSECTS: Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths, Fruit Flies. Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disperse in all locations. Keep area closed for 15 minutes.

Adult Fleas in Buildings: (Not on Animals) Close windows and doors. Hold aerosol 36" above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 square feet, making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatments after 6 days.

Feed Storage Areas, Meat and Food Processing Plants:

FLYING INSECTS: Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths, Fruit Flies. Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disperse in all locations. Keep area closed for 15 minutes.

GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, CARPET BEETLES, DRUG STORE BEETLES: In exposed adult and larvae stages. Contact as many insects as possible when treating infested areas such as cracks, crevices, stored products, machinery, and dark secluded areas. For Carpet Beetles, treat infested area of carpets, localized areas of floors, baseboards, and into cracks and under carpets where these pests may be found. Spray 10 to 20 seconds per 1000 square feet of floor space. Repeat treatment weekly or as insects appear.

ROACHES, SPIDERS, ANTS, SILVERFISH, CRICKETS: Hold aerosol close to cracks and mist directly into them, contacting as many insects as possible. Apply freely in all possible hiding places behind baseboards, sinks, cabinets, coffee makers, meter boxes, etc. Repeat all treatments as necessary.

**Manufactured By
Whitmire Research Laboratories, Inc.
St. Louis, Missouri 63122**

NET WEIGHT 2 LBS.

**Only for Sale to, Use, and
Storage by Servicepersons.**

© 1989 Whitmire Research Laboratories, Inc. All rights reserved. EPA Reg. No. 499-208-AA. EPA Est. No. 499-MO-1.

300-115384
1767

FEB 8 1984

Mr. M. G. Sarli
Whitacre Research Laboratories Inc.
3568 Tree Court Industrial Boulevard
Saint Louis, MO 63122

Dear Mr. Sarli:

Subject: Whitacre Fly-Off Pans Insect Repellent
EPA Registration No. ~~007480~~
Your Letter Dated December 16, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure
DCR-05649:A. Hayward:RD-20:1/27/84:del.2/20/84:LM

BEST DOCUMENT AVAILABLE

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							

EPA Form 1320-1 (4-81) OFFICIAL FILE COPY