

PRECAUTIONARY STATEMENTS
CAUTION
Hazards to Humans and Domestic Animals

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. When using this product as a space spray in food areas, do not apply while food processing is under way. Except in federally inspected meat and poultry plants, careful crack and crevice application for inspection purposes is permissible at all times. After space spraying in meat packing plants, bakeries, and other food processing plants, wash with potable water all equipment, benches, shelves, etc. where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

Environmental Hazards

This product is toxic to fish and birds. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Physical or Chemical Hazards

Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F. may cause bursting. Do not spray on plastic, painted, or varnished surfaces.

USE DIRECTIONS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in a cool area away from heat or open flame.

Pesticide Disposal: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Container Disposal: Replace cap and discard empty container in trash, sanitary landfill, or by other approved state and local procedures. Do not incinerate or puncture.

AGRICULTURAL USES: (For control of House Flies, Horn Flies, Stable Flies, Gnats, Mosquitoes, Small Flying Moths)

Dairy Farm Use: MILKING PARLOR AND MILK ROOM — Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. If rooms cannot be closed tight, double the dosage. Stand in middle of room. Direct fog upward and spray in all directions. Seconds may be counted by saying one-thousand, two-thousand, etc. Keep room closed for 15 minutes after treatment. Example: A room 20 feet long, 12 feet wide, and 8 feet high requires 2 seconds if closed tight; 4 seconds if not. Do not remain in treated areas. Ventilate the area after treatment is completed. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

ANIMAL USE — From approximately 2-foot distance, thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where saliva accumulates from head tossing.

STANCHION BARN USE — Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

(Continued)



Whitmire

PT 10
DAIRY

AND
Farm Insect Fogger

WITH DUAL ACTIVATORS

FOR: Dairy Farms (including Milk House, Milking Parlor, Loading Sheds, and Holding Lot), Poultry and Hog Operations, Horse Barns and Stables, Beef Cattle Operations, Kennels, Catteries, Food Storage Areas, and Meat Packing and Food Processing Plants.

KILLS: House Flies, Horn Flies, Stable Flies, Gnats, Mosquitoes, Wasps, Small Flying Moths, Fruit Flies, Adult Fleas, Grain Weevils, Flour Beetles, Chocolate Moths, Carpet Beetles, Drug Store Beetles, Roaches (Waterbugs), Spiders, Ants, Silverfish, Crickets.

ACTIVE INGREDIENTS

Pyrethrins	0.500%
*Technical Piperonyl Butoxide	1.000%
N-octyl bicycloheptene dicarboximide	1.000%
Refined Petroleum Oil	8.000%
INERT INGREDIENTS	89.500%

*Equivalent to 0.8% (Butylcarbityl) (6-Propylpiperonyl) Ether and 0.2% of Related Compounds.


EPA Reg. NO. 499-208-AA • EPA Est. No. 499-MO-1



KEEP OUT OF REACH OF CHILDREN.

CAUTION:

See side panel for additional precautions.



499-208 171
PM 17
USE DIRECTIONS Continued

Poultry Operations: Use only in ground poultry houses (when birds are not present). Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.

Hog Operations: In hog houses close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

Horse Barns and Stables: In horse and pony barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application, lightly mist over the backs of horses and ponies from a distance of at least 2 feet. Do not spray directly towards horse's face or eyes. Repeat application daily or as necessary.

Beef Cattle Operations: In barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal use thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

NON-AGRICULTURAL USES

Kennels & Catteries: Remove animals before spraying. Ventilate after treatment is completed.

FLYING INSECTS: Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths, Fruit Flies. Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disperse in all locations. Keep area closed for 15 minutes.

Adult Fleas in Buildings: (Not on Animals) Close windows and doors. Hold aerosol 36" above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 square feet, making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatments after 6 days.

Food Storage Areas, Meat and Food Processing Plants:

FLYING INSECTS: Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths, Fruit Flies. Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disperse in all locations. Keep area closed for 15 minutes.

GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, CARPET BEETLES, DRUG STORE BEETLES in exposed adult and larvae stages: Contact as many insects as possible when treating infested areas such as cracks, crevices, stored products, machinery, and dark secluded areas. For Carpet Beetles, treat infested area of carpets, localized areas of floors, baseboards, and into cracks and under carpets where these pests may be found. Spray 10 to 20 seconds per 1000 square feet of floor space. Repeat treatment weekly or as insects appear.

ROACHES, SPIDERS, ANTS, SILVERFISH, CRICKETS: Hold aerosol close to cracks and mist directly into them, contacting as many insects as possible. Apply freely in all possible hiding places behind baseboards, sinks, cabinets, coffee makers, meter boxes, manholes. Repeat all treatments as necessary.

Manufactured By

Whitmire Research Laboratories, Inc.
St. Louis, Missouri 63122

Copyright 1985

NET WEIGHT 2 LBS.

Authorized by USDA for use in federally inspected meat and poultry plants.

DOT-E 7808