



**PT 3-6-10
Aero-Cide**



Only for sale to, use and storage by servicepersons.



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Do not remain in treated area and re-ventilate the area after treatment is completed. Avoid breathing of vapor. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using and before smoking or eating. Do not consume water, food, or tobacco. Remove all pets and birds. Cover fish tanks or remove before spraying.

NOTICE TO APPLICATOR: During any general fumigation treatment, wear a respirator (respirator model approved by the Mining Enforcement and Safety Administration formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds, and other animals. Do not allow to come in contact with water. Apply this product only as specified on this label.

Food Areas of Food Handling Establishments: Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product as a space spray, apply only when part is not in operation. Except in food areas, inspect for and destroy pests, clean floors and crevices, application for insecticidal purposes is prohibited.

PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not puncture or otherwise damage container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

Do not spray on or into painted or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers, etc.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL: Store in a cool area away from heat or open flame. Do not contaminate water, food, or feed by storage or disposal. Do not reuse empty container. Wrap container and put in trash can.

GENERAL VOLUMETRIC TREATMENT: Flying insects: Flies, Mosquitoes, Gnats, Fruit Flies, Moths, Fruit Flies, Cuckoo Flies, Cattle Flies, Cattle and Horse Flies, and other flies. Spray at a rate of 1 to 3 seconds per 1000 cu. ft. for 15 to 30 minutes. Repeat treatment as necessary.

Moths, Beetles, and Wasps: Spray most directly on insects and their nests from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

Carpet Beetles, Drug Store Beetles, Cigarette Beetles, Confused Flour Beetles, Sawtoothed Grain Weevils, Rice Weevils, Red Flour Beetles, Grain Weevils, and other stored grain and seed insects: Contact as many insects as possible. Spray into cracks and crevices, processing and handling equipment, stored grains, and containers, and other areas where these insects tend to congregate. Spray 20 seconds per 1000 cu. ft. of space for 15 to 30 minutes. Repeat treatment weekly or as insects appear. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Beetles, Sawflies, Spiders, and other insects of basements, attics, and other areas: Spray directly on insects in these locations where possible. Repeat as necessary.

Flies in Buildings: Close windows and doors. Feed aerosol 36" above floor and drift spray from floor and lower walls at a rate of 10 seconds per 1000 cu. ft. making sure that all four walls and other surfaces are contacted. Repeat treatment after 7 days if necessary.

Indian Meal Moth, Chocolate Moth, Warehouse Flour Moth, Angoumois Grain Moth, and other stored grain and seed insects: Contact spray into all parts of room especially around stored product, surfaces, pipes, and fixtures. Spray 10 to 30 seconds per 1000 cu. ft. of space for 15 to 30 minutes. Repeat treatment weekly or as insects appear. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Cowflies: Contact as many insects as possible when flying around a lot of food. In areas where insects are not flying around windows and door frames, spray 10 to 30 seconds per 1000 cu. ft. of space. Repeat as necessary.

Bed Bugs: Spray mattresses, particularly around tufts and seams. Spray bed, sofa, and other furniture. Treat base boards, floor cracks, and crevices, including surfaces and doors. Repeat treatment as necessary.

Roaches, Spiders, House Flies, Ants, Sawflies, Crickets, Centipedes, Sow Bugs, and other household insects: Contact spray directly into their hiding places in many insects as possible. Apply 10 to 30 seconds per 1000 cu. ft. of space. Repeat treatment as necessary.

VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS: This product is for use in inaccessible voids and equipment rooms. For use, application with this product is necessary.

For exposed stages of cockroaches, spiders, white bugs, ants, sawflies, crickets, centipedes, sow bugs, and other household insects: Contact spray into all parts of room especially around stored product, surfaces, pipes, and fixtures. Spray 10 to 30 seconds per 1000 cu. ft. of space for 15 to 30 minutes. Repeat treatment weekly or as insects appear. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

OUTDOOR GROUND APPLICATION: Use this product to control flying insects, including mosquitoes, flies, and other flying insects. Spray 10 to 30 seconds per 1000 cu. ft. of space for 15 to 30 minutes. Repeat treatment as necessary.

Copyright 1983 Whitmire Research Laboratories, Inc., St. Louis, MO

**PT 3-6-10
Aero-Cide**

Insect Fogger

KILLS: Flies, Mosquitoes, Gnats, Small Fly Moths, Fruit Flies, Cluster Flies, Mud Daubers, Wasps, Cockroaches, Waterbugs, Ants, Silverfish Spiders, Crickets, Centipedes, Millipedes, Booklice, Sow Bugs, Fleas, Bed Bugs, Indian Meal Moths, Chocolate Moths, Mediterranean Flour Moths, Angoumois Grain Moths, Clover Mites, Carpet Beetles, Drug Store Beetles, Cigarette Beetles, Confused Flour Beetles, Sawtoothed Grain Beetles, Curculionid Weevils, Rice Weevils, Red Flour Beetles and Grain Mites.

For Use In and Around Meat Packing and Food Processing Plants, Food Storage Areas, Restaurants and other Food Handling Establishments, Hospitals, Supermarkets, Motels, Motel's Resort, Homes, Transportation Equipment (Buses, Bait Shops, Trains, Trucks, Planes), Utilities, Schools, Campgrounds, Warehouses and other Commercial and Industrial Buildings.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide, Technical*	2.00%
n-Octyl Bicycloheptene Dicarboximide	3.33%
Refined Petroleum Oil	8.00%
INERT INGREDIENTS	85.67%

EPA Reg. No. 499-221-AA EPA Est. No. 499-MD-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

For Cockroaches: Contact spray directly on roaches and their nests from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Bed Bugs: Spray mattresses, particularly around tufts and seams. Spray bed, sofa, and other furniture. Treat base boards, floor cracks, and crevices, including surfaces and doors. Repeat treatment as necessary.

For Mosquitoes: Contact spray directly on mosquitoes and their nests from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Flies: Contact spray directly on flies and their nests from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Beetles: Contact spray directly on beetles and their nests from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Moths: Contact spray directly on moths and their nests from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Spiders: Contact spray directly on spiders from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Crickets: Contact spray directly on crickets from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Centipedes: Contact spray directly on centipedes from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Sow Bugs: Contact spray directly on sow bugs from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Fleas: Contact spray directly on fleas from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Fruit Flies: Contact spray directly on fruit flies from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Cluster Flies: Contact spray directly on cluster flies from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Mud Daubers: Contact spray directly on mud daubers from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Wasps: Contact spray directly on wasps from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Ants: Contact spray directly on ants from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Silverfish Spiders: Contact spray directly on silverfish spiders from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Booklice: Contact spray directly on booklice from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Millipedes: Contact spray directly on millipedes from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Centipedes: Contact spray directly on centipedes from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

For Sow Bugs: Contact spray directly on sow bugs from approximately 20 ft. distance. Do not spray directly into food. Application should be made in late evening when insects are most active. Spray into cracks and breeding places, including as many insects as possible.

BES. LABLE



ACCEPTED
with COMMENTS
to EPA Letter #

SEP 15 1983

Under the Fungicide, Insecticide, and Plant
as amended, the pesticide
registered under FPA Reg. No.
494-27-1

300 / 104800
17 / 3

SEP 15 1983

Mr. Richard Vega
Whitire Research Lab., Inc.
3500 Tree Court Industrial Blvd.
St. Louis, MO 63122

Dear Mr. Vega:

Subject: Whitire 15(R) 3-0-10 Zero. Cide
LRA Registration No. 699-221
Your amended Application of August 20, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Also we have no objection to the subject name change.

Sincerely yours,

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (1E-707)

Enclosure

RE:Alleyard HCL-17195:WARG-0321C:AM:Kaven:679-2013:9/13/83

SYMBOL	79-767								
NAME	Alleyard								
DATE	Sept								

EPA 1-117-1-45

OFFICIAL FILE COPY