



**FOR USE WITH SYSTEM III  
OR HAND HELD USE**

# Aero-Cide<sup>®</sup>

PT<sup>®</sup> 3-6-10  
**Pyrethrum**

**PT<sup>®</sup> 3-6-10 Aero-Cide<sup>®</sup>**

## Insect Fogger

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Do not remain in treated area and ventilate the area after treatment is completed. Avoid breathing of vapor. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using and before smoking or eating. Do not consume raw water, food, or feedstuffs. Remove all pots and pans. Cover fish tanks or remove before spraying.

During any general volumetric treatment, wear a approved respirator, only approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute of Occupational Safety & Health under the provisions of 30 CFR Part 11.

**ENVIRONMENTAL HAZARDS**—This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Apply this product only as specified on this label.

**Food Areas of Food Handling Establishments**—Goods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product as a space spray, apply only when pest is not in operation. Escape in Federally inspected meat and poultry plants, carcass racks and device application for inspection purposes is permissible at all times.

**PHYSICAL OR CHEMICAL HAZARDS**—Contains under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces or directly into any electronic equipment such as radios, televisions, etc.

**DIRECTIONS FOR USE**—It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**—Do not contaminate water, food, or feed by storage or disposal. Storage: Store in a cool area away from heat or open flame.

**Pesticide Disposal:** Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

**Container Disposal:** Replace cap and discard empty container in trash, sanitary landfill, or by other approved state and local procedures. Do not incinerate or puncture.

**GENERAL VOLUMETRIC TREATMENT:** Flying insects: Flies, Mosquitoes, Gnats, Small Flying Moths, Fruit Flies, Cluster Flies, Cope of windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cu. ft. for 8 to 15 seconds per 1000 cu. ft. for 8 to 15 ft. size due to delivery rate. Fill room with fog. Disperse in all locations. Keep area closed for 15 min. Suck up and destroy taken insects. Repeat treatment as necessary.

**Head Lice and Head Nits:** Spray mist directly on insects and their eggs from approximately 2 ft. distance. Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Spray into hair and bedding (nits) contacting as many insects as possible.

**Target Beetles:** Drug Store Beetles, Cigarette Beetles, Confused Flour Beetles, Spotted Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites in exposed adult and larval stages. Contact as many insects as possible. Spray into cracks and crevices, processing and handling equipment, around cartons and containers, and other areas where these insects tend to congregate. Spray 20 seconds per 1000 cu. ft. of space. For 8 to 15 ft. size spray 10 seconds per 1000 cu. ft. Repeat treatment weekly or as insects appear. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

**Spiders, Silverfish:** Spray cracks and crevices in basements, door frames, porches, and storage areas. Spray directly on insects in these locations where possible. Repeat as necessary.

**Flies in Buildings:** Close windows and doors. Hold sprayer 30" above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq. ft., making sure that all floor, side, and chair surfaces are contacted. Repeat treatment after 7 days, or as necessary.

**Indian Meal Moth, Chocolate Moth, Mediterranean Flour Moth, Angoumois Grain Moth, Cope of doors and windows:** Direct spray into all parts of room, especially around stored product containers, shelves, and corners. Spray 5 to 10 seconds for each 1000 cu. ft. of space. For 8 to 15 ft. size, spray 5 to 5 seconds per 1000 cu. ft. Repeat area closed for 15 minutes following treatment. Repeat treatment weekly or as necessary. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

**Closet Mites:** Contact as many insects as possible when treating infested areas such as cracks and crevices. It will work around windows and door frames, under siding, and in dark secluded areas. Repeat as necessary.

**Roaches, Spiders, Water Bugs, Ants, Silverfish, Crickets, Centipedes, Sow Bugs, Head Lice:** Spray directly into cracks and most directly into them, contacting as many insects as possible. Apply treat in all possible hiding places behind baseboards, sinks, cabinets, coffee makers, meat boxes. Repeat 2-3 treatments as necessary.

**VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS:** Ants, Fire Ants, etc. Use and equipment voids. For spray application with Whittmore Crack & Crevice Injection equipment.

**For exposed stages of Cockroaches, Spiders, Water Bugs, Ants, Silverfish, Crickets, Centipedes, Sow Bugs and Beetles:** Calculate void volume and treat at a rate of 7 seconds per 1000 cu. ft. for 8 to 15 ft. size, 10 seconds per 1000 cu. ft. Spray by packing material to into cracks or crevices, holes, or other voids. Spray into void space. Repeat treatment as necessary.

**OUTDOOR GROUND APPLICATION:** To kill Flies, Mosquitoes, and Gnats in open areas near buildings and in lawns, etc. Best results are obtained when the temperature is cool (75°F or less) and the wind speed is 1 to 15 MPH or less. Apply at a rate of 15-30 seconds per acre for 8 to 15 ft. size, spray 30-40 seconds per acre. Spray in wide swaths across area to be treated. Allow spray drift to penetrate dense foliage. Repeat treatment as necessary.

© Copyright 1985 Whitmire Research Laboratories, Inc., St. Louis, MO

**KILLS:** Flies, Mosquitoes, Gnats, Small Fly Moths, Fruit Flies, Cluster Flies, Mud Daubers, Wasps, Cockroaches, Waterbugs, Ants, Silverfish, Spiders, Crickets, Centipedes, Milkbeetles, Booklice, Sow Bugs, Fleas, Bed Bugs, Indian Meal Moths, Chocolate Moths, Mediterranean Flour Moths, Angoumois Grain Moths, Clover Mites, Carpet Beetles, Drug Store Beetles, Cigarette Beetles, Confused Flour Beetles, Sawtoothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles and Grain Mites.

For Use in and Around Meat Packing and Food Processing Plants, Food Storage Areas, Restaurants and other Food Handling Establishments, Hospitals, Supermarkets, Motels, Hotels, Resorts, Homes, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes), Utilities, Schools, Campgrounds, Warehouses, and other Commercial and Industrial Buildings.

**ACTIVE INGREDIENTS:**

Pyrethrins	1.00%
Piperonyl Butoxide, Technical	2.00%
n-Octyl Bicycloheptene Dicarboximide	3.33%

**INERT INGREDIENTS** ..... 93.67%


\*Equivalent to 1.6% (Sulfacarbonyl (5-Propoxyphenyl) Ether and 0.4% Residue Compounds)  
EPA Reg. No. 499-221-AA EPA Est. No. 499-MO-1

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

If available: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. Do not induce vomiting. Vomiting may cause absorption. If inhaled: Remove patient to fresh air. Apply artificial respiration if necessary. If on skin: Remove contaminated clothing and wash affected skin areas with soap and water. Get medical attention if irritation persists. If in eyes: Wash eyes with clean water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.



Authorized by USDA for use in Federally inspected meat and poultry plants

18-2362 Rev. Date 2/87

Reviewed, [illegible] [illegible] [illegible]