

R-493-223
 0115
 1/10/1

WhitMire PT 230 Tri-Die Insecticide Spray



PT[®] 230 Tri-Die Insecticide Spray

SHAKE WELL BEFORE USE



PT[®] 230 Tri-Die

Trade Name Accepted

Activated Pyrethrum in Silica Aerogel

CONTACT-REPELLENT-RESIDUAL INSECTICIDE CRACK & CREVICE[®] INJECTION SYSTEM

Kills: Cockroaches, crickets, beetles, spiders, earwigs, sowbugs, pillbugs, millipedes, centipedes, booklice, silverfish, firebrats, ants, cluster flies, ear flies, mealworms, boxelder bugs, fleas, ticks, lice, bedbugs, caterpillars, cheese mites, grain mites, cigarette beetles, drugstore beetles, confused flour beetles, red flour beetles, dark mealworms, yellow mealworms, sawtoothed grain beetles, flat grain beetles, granary weevils, rice weevils, drywood termites, powder post beetles, ground beetles, and scorpions.

For use in hospitals, schools, supermarkets, motels, hotels, homes, apartments, transportation equipment (buses, boats, ships, trucks, trailers, trains, planes), utilities, meat packing and food processing plants, food storage areas, restaurants and other food handling establishments, warehouses, and other commercial buildings.

Authorized by USDA for use in federally inspected meat and poultry plants.

ACTIVE INGREDIENTS	
Pyrethrins	0.3%
*Piperonyl Butoxide Technical	3.0%
Silica Gel	4.0%
Petroleum Distillate	5.7%
INERT INGREDIENTS	87.0%

*Equivalent to 2.4% (butyl arbutyl) 6-propylperonyl ether and 0.6% related compounds.

EPA Reg. No. 499-223-AA EPA Est. No. 499-MO-1

KEEP OUT OF REACH OF CHILDREN
 WARNING:

See side panel for additional cautions.

NET WT. 1 LB. 2 OZS.

© Copyright 1983 WhitMire Research Laboratories, Inc.

ACCEPTED
 MAY 22 1985
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the purposes registered under EPA Reg. No. 499-223.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Avoid eye irritation. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before smoking or eating. Avoid contamination of food and foodstuffs. Do not use in serving areas while food is exposed.

Notice to Operator: In order to avoid any increase in the risk of eye and skin irritation when dispensing PT 230 Tri-Die, confined areas or when prolonged exposure to spray mist occurs, wear a pesticide dust respirator fully approved by the Mining Enforcement and Safety Administration, formerly the U.S. Bureau of Mines, and by the National Institute of Occupational Safety and Health under the provisions of 30 CFR Part 117.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES - Flush eyes with plenty of water for several minutes.

IF SWALLOWED - Call a physician or Poison Control Center immediately. Gastric lavage may be indicated. Medical aid given internally. DO NOT INDUCE VOMITING. Do not use emetics. Cause aspiration pneumonia.

IF INHALED - Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN - Remove contaminated clothing and wash affected areas with soap and water.

PHYSICAL OR CHEMICAL HAZARDS

Flammable: Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 170°F may cause bursting.

Do not spray on paints, PT 230 Tri-Die is flammable. Do not spray on plastic, painted or varnished surfaces. Do not spray directly into any electronic equipment such as radios, televisions, computers, motors, or into outlets and switches or any other location where the powdery residue may cause electrical contacts and circuits.

DIRECTIONS FOR USE
 This is a variation of Federal Law to use this product in a manner consistent with this labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry place away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Vessel containing form of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Recycle, cap and discard container in trash, incinerate (landfill) or by other approved state and local procedure. Do not incinerate or puncture.

SHAKE WELL BEFORE USE: Before shaking after use, invert container and spray until air comes out in clear, uniform spray mist.

How to Use WhitMire Crack and Crevice Injection Equipment

Do not use as a space spray. Except when used as a contact spray for flea control, WhitMire's PT 230 Tri-Die is for use only with injection tube for injecting into cracks and crevices where insects may be hiding, feeding, and breeding. Do not inject into cracks, crevices, hoses, and other small openings where insects may be a problem. Release applicator in second spray period in serious infestations. 10 ticks per square foot 12" apart. In light infestations or for preventive treatment, place spots approximately 3 ft apart in areas with 2 cracks. Move the injector tip along the crack while spraying. Move at the rate of one linear foot per second in heavy infestations and three feet per second in light infestations. See additional directions below for effective control of specific insects.

INDOOR PESTS -

Crawling Pests: Cockroaches, crickets, beetles, spiders, earwigs, sowbugs, millipedes, centipedes. Spray into hiding places such as cracks and crevices, moist areas, openings around pipes and sinks, under refrigerators, behind baseboards, coffee makers, in dark storage areas, and in other areas where these pests are found. Spray directly on insects in these locations where possible. Repeat as necessary.

Flies, Silverfish, Firebrats: Spray cracks and crevices in flycatcher, door frame, bookcase, and storage area. Spray directly on insects in these locations where possible. Repeat as necessary.

WHITMIRE RESEARCH LABS, INC.
 St. Louis, MO 63122

Ants: Pour ant colonies and spray 5-10 seconds directly into nest. Spray into holes and points of entry around doors, windows, structural cracks and crevices. Spray directly on insects in these locations where possible. Repeat as necessary.

Microzing stages of: Fleas, ticks, lice, ear lice, beetles, and boxelder bugs. Contact as many insects as possible when treating infested areas such as cracks and crevices, wall voids, around windows and door frames, under siding in machinery and in other infested areas. Repeat as necessary.

Flea, ticks, and lice in animal quarters: Treat bedding, floor and floor coverings and all areas where pests may hide at the rate of one second spray per square foot. Treat cracks and crevices, behind moldings, shelving, baseboards, pipe openings, and all other cracks and crevices at the rate of one second spray per square foot. Repeat as necessary. A deposit of PT 230 Tri-Die will remain in cracks and crevices to prevent re-infestation. Repeat application when other insects dominate.

Bedbugs: Take bed apart and treat cracks, joints, and crevices in mattress. Mattresses and box springs should be treated especially at tufts, folds, and edges. Bedside frames, headings and all cracks and crevices in the room should also be treated. Do not use in hospital patient quarters.

Stored product pests: Cuckoo, cheese mites, grain mites, cigarette beetles, drugstore beetles, confused flour beetles, red flour beetles, dark mealworms, yellow mealworms, sawtoothed grain beetles, flat grain beetles, granary weevils, and rice weevils, sawtoothed grain beetles, granary weevils, and other pests. Apply in cracks and crevices and other hiding places. Do not use in hospital patient quarters.

Drywood Termites: Spot treatment for local infestations. Probe to locate galleries, holes and nest. 5-10 seconds spray per hole into the active galleries.

Powder Post Beetles: Treat infested wood using injection tube. Apply small amounts directly into and around entrance holes. Systemic treatment of wood by boron or other preservative. Use wood preservative in areas, cracks, spaces, and junctures where human contact with treated surfaces will be minimal. For spray application with WhitMire Crack & Crevice Injection Equipment: For exposed surfaces of construction, earwigs, spiders, sowbugs, millipedes, centipedes, booklice, silverfish, firebrats, ants, and beetles. Calculate total horizontal surface area and treat at the rate of one second per square foot. Spray by inserting injection tube into cracks or crevices. Move injector tip along the crack while spraying. Repeat as necessary.

APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS - Places other than private residences in which food is held, processed, prepared or served, including when family is not in operation.

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY: Includes areas for receiving, storage, packing, cleaning, bottling, wrapping, boxing, preparing, and the warehousing and enclosed processing systems (meat, dairy, vegetable, poultry, serving areas) when food is exposed to the public in operation. Do not use in residential food areas.

Apply in small amounts directly into cracks and crevices using injection tube in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces, in wall, equipment legs and bases where insects hide. Care should be taken to avoid depositing the contact onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A CRACK AND CREVICE TREATMENT ARE NOT PERMITTED

For application ideas for both food and non-food areas, refer to the WhitMire Branch Location Chart and Pest Management Manual.

OUTDOOR PESTS

Cockroaches, sowbugs, pillbugs, centipedes, earwigs, ground beetles, earwigs, crane flies, scorpions, and ants. Clean up debris around premise. Spray in cracks around doors, windows, under siding at foundation level, beneath guards on loading docks, into cracks in foundations and terraces and in other places where these pests may enter building. Repeat application as necessary.