

499-289

04/04/2000

1/1

waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Sealing areas (when food is exposed and facility is in operation) is also considered a food area.

Apply prescribed amounts of PT* 259 as a Crack and Crevice* application using the supplied injection tube or other Whitmire Micro-Gen equipment in points between different elements of construction, between equipment and floor, openings leading to voids and hollow spaces in walls, equipment legs and bases where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food, food utensils or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD-HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE* TREATMENT ARE NOT PERMITTED.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Waste resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: This container may be recycled in aerosol recycling centers. At present, there are only a few such centers in the country. Before offering for recycling, empty the can by using the product according to the label (DO NOT PUNCTURE!). If the recycling is not available, wrap the container and discard in the trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Do not get in eyes, on skin, or on clothing. Harmful if swallowed or absorbed through the skin. Avoid breathing of spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid contamination of food, utensils, and food preparation areas. Should illness occur, get prompt medical aid.

To Physician: Atropine sulfate is antidotal.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Apply this product only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS

Extremely Flammable. Contents under pressure. Keep away from fire, sparks, and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

Do not use this product in or on electrical equipment due to the possibility of shock hazard. DO NOT SPRAY ON PLASTIC, PAINTED, OR VARNISHED SURFACES or directly into any electronic equipment such as radios, televisions, computers, etc. Aerosols work best when can temperature is above 70° F. If can temperature is below 60° F, store at room temperature until a temperature above 60° F is reached.


Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



A Prescription Treatment* brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
www.wmmg.com

© 1999 Whitmire Micro-Gen Research Laboratories, Inc.



KILLS:

Ants, Booklice, Boxelder Bugs, Carpenter Ants, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs, Elmleaf Beetles, Flour Beetles, Grain Weevils, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders, Ticks, and Trogoderma, (Cabinet, Khapra, and Warehouse Beetles).

FOR USE IN AND AROUND:

Apartments, Homes, Hospitals, Hotels, Food Storage Areas, Inedible Product Areas of Meat Packing Plants, Food Processing Plants, Motels, Nursing Homes, Restaurants and Other Food Handling Establishments, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains and Planes), Warehouses and Other Commercial and Industrial Buildings.

* Do not use in aircraft cabins.

ACTIVE INGREDIENT:

Propoxur 1.0%

OTHER INGREDIENTS: 99.0%

KEEP OUT OF REACH OF CHILDREN CAUTION FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

See side panel for additional precautionary statements


EPA Reg. No. 499-289 • EPA Est. No. 499-MO-1

NET WEIGHT:

800-777-8570



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES INC.



DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

APPLICATION RESTRICTIONS

Do not apply this product in a way that will contact any person or pet, either directly or indirectly. Keep people and pets out of the area during application. Do not apply in aircraft cabins.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

APPLICATOR MUST WEAR: long sleeved shirt and long pants, chemical-resistant gloves, and shoes plus socks.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove protective clothing and equipment immediately after handling this product. Wash the outside of gloves before removing. Keep and wash protective clothing and equipment separately from other laundry.

HOW TO USE WHITMIRE MICRO-GEN CRACK AND CREVICE* INJECTION SYSTEM

CRACK AND CREVICE* APPLICATIONS: Use Whitmire PT* 259 with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment. Inject PT* 259 into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. Calculate the void's cubic area and treat at the rate of 5 to 10 seconds per 3 cubic feet. Several holes may be required in long-running voids. Repeat as necessary.

INDOOR TREATMENTS

CRAWLING INSECTS - Cockroaches, Dermestids, Scorpions, Spiders, Earwigs, Crickets, Sowbugs, Millipedes, Centipedes: Apply as a Crack and Crevice* application into cracks, crevices and void spaces where insects may be harboring. Inject into openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewer, floor drains and around and in meter boxes. Repeat as necessary.

TICKS: Apply as a Crack and Crevice* application into infested and egg laying sites near pet resting areas such as cracks and crevices, behind baseboards, door and window frames.

BOOKLICE & SILVERFISH: Apply as a Crack and Crevice* application into cracks and crevices associated with baseboards, door frames, bookcases and other harborage sites.

ANTS: Apply as a Crack and Crevice* application into cracks, crevices and voids where ants may be traveling and/or harboring, around doors, window frames, wall voids and other structural cracks and crevices.

CARPENTER ANTS: Locate source of infestation. Inject 5-10 seconds of product into insect tunnels and cavities. Treat trails, points of entry or exits from voids such as around doors and windows.

FOR EXPOSED ADULT AND LARVAL STAGES OF GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, DRUGSTORE BEETLES, TROGODERMA: Apply as a Crack and Crevice* application into cracks, crevices and voids where these insects may be harboring, cracks and crevices in cabinets, pantries, wall voids and in machinery.

FOR ACTIVE AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Apply as a Crack and Crevice* application into cracks, crevices and voids where these insects may be harboring or hibernating. Treat cracks and crevices, wall voids, around window and door frames.

APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS

Food Handling Establishments are places other than private residences in which food is held, processed, prepared or served.

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE* TREATMENT ONLY: Includes areas for receiving, storage, age, packing (canning, bottling, wrapping, boxing), preparing, edible