


FRONT PANEL:

4/24/92



For Use With System III[®]
 Whitmire
 Engage™
 Residual Injection System

KILLS: Ants, Bees, Booklice, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs, Elmleaf Beetles, Fire Ants, Flour Beetles, Grain Weevils, Hornets, Millipedes, Scorpions, Silverfish, Sawbugs, Spiders, Springtails, Termites, Ticks, Trogodermas, Yellow Jackets, Wasps, Wood Infesting Borers and Beetles, and Wood Wasps.

For use in and around apartments, homes, hospitals, hotels, food storage areas, inedible product areas of meat packing plants, food processing plants, motels, nursing homes, restaurants and other food handling establishments, supermarkets, transportation equipment (buses, boats, ships, trains, planes) and warehouses and other commercial and industrial buildings.

Authorized by USDA for use in inedible areas of Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Chlorpyrifos (0,0-Diethyl 0-(3,5,6-Trichloro-2-Pyridyl) Phosphorothioate) 0.5%

INERT INGREDIENTS: 99.5%

**KEEP OUT OF REACH OF CHILDREN
 CAUTION
 STATEMENT OF PRACTICAL TREATMENT**

IF SWALLOWED: Call a physician or poison control center immediately. Gastric lavage is indicated. Do not induce vomiting. Vomiting may cause aspiration pneumonia.

IF INHALED: Remove patient to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: If symptoms of cholinesterase inhibition are present, atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

See side panel for additional precautionary statements

EPA Reg. No. 499-292 EPA Est. No. 499-MO-1
 NET WEIGHT:



272

))

Whitmire PT[®] 279 ^{Enzyme™}
Residual Injection System

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO USE WHITMIRE CRACK AND CREVICE[®] INJECTION SYSTEM: Use Whitmire PT[®] 279 only with the supplied actuator and injection tubes or other Whitmire equipment. Inject PT[®] 279 into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For closed voids calculate the void's cubic area and treat at the rate of five to ten seconds per three cubic feet. Several holes may be required in long-running voids. Repeat as necessary.

INDOOR TREATMENTS

Crawling Insects: Cockroaches, Dermestids, Scorpions, Spiders, Earwigs, Crickets, Sowbugs, Millipedes, Centipedes: Inject with the prescribed dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewer, floor drains and meter boxes. Repeat as necessary.

Ticks: Thoroughly treat infested and egg laying sites near pet resting areas such as cracks and crevices, behind baseboards, door and window frames.

Booklice & Silverfish: Inject into cracks and crevices associated with baseboards, door frames, bookcases and other harborage sites. Repeat as necessary.

Ants: Inject cracks, crevices and voids where ants may be traveling and/or harboring, including, but not limited to, around doors, window frames, wall voids and other structural cracks and crevices.

For exposed adult and larval stages of Grain Weevils, Flour Beetles, Chocolate Moths, Drugstore Beetles, Trogodermas: Inject into cracks and crevices and/or voids where these insects may be harboring. Infested areas include, but are not limited to, cracks and crevices in cabinets, pantries, wall voids and in machinery. Repeat treatment as necessary.

APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS: Places other than private residences in which food is held, processed, prepared or served.

BEST AVAILABLE COPY

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE[®] TREATMENT ONLY:

Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) is also considered a food area.

Apply prescribed amounts of PT[®] 279 directly into cracks and crevices or voids using injection tube or other Whitmire equipment in points between different elements of construction, between equipment and floor, openings leading to voids and hollow spaces in walls, equipment legs and bases where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food, food utensils or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD-HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE[®] TREATMENT ARE NOT PERMITTED.

INDOOR AND OUTDOOR TREATMENTS

Carpenter Ants: Locate source of infestation. Inject 5-10 seconds of product into insect tunnels and cavities. Treat trails, points of entry or exits from voids such as around doors and windows. Treatment of trails on tree trunks or fences is permitted.

For active and hibernating stages of Boxelder Bugs, Elmleaf Beetles, Cluster Flies and Clover Mites: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames. Repeat as necessary.
Side Panel (cont.)

OUTDOOR TREATMENTS

Subterranean ants such as Fire Ants, Argentine Ants and Harvester Ants, etc.: To treat nests, use the Whitmire soil injector. To maximize insecticide distribution, insert injector into the nest full length if possible. Inject the nest at 3-6 sites at a rate of 5-10 seconds per site. For very large nests (over 36" diameter), increase the number of injection sites. Space injection sites in a circular pattern on the nest surface with one site in the center. To quiet active ants on nest surface, sweep surface with spray from a distance of 12 inches. For best results, apply when temperature is 65-85°F, or in early morning or late evening hours. Treat new nests as they appear.

CARPENTER BEES: Locate entrance holes. Inject 5 - 10 seconds of product into entrance holes. Seal bee entrances after application.

PERIMETER TREATMENT: For crawling insects, apply PT[®] 279 to cracks and crevices where insects may enter premises. Areas to be treated include, but are not limited to, wasp holes, under siding, wall voids, soffits, around attic vents, cracks and crevices openings around windows, doors and pipes. Treatment of infested fences and tree holes is permitted.

BEST AVAILABLE COPY

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Replace cap and discard container in trash, sanitary landfill or by other approved state and local procedures. Do not incinerate or puncture.

PRECAUTIONARY STATEMENT HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Do not get in eyes, on skin, or on clothing. Harmful if swallowed or absorbed through the skin. Avoid breathing of spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Avoid contamination of food, utensils, and food preparation areas. Should illness occur, get prompt medical aid. To Physician-Atropine sulfate is antidotal.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and birds. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard. Do not spray on plastic, painted or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers, telephones, etc.

BEST AVAILABLE COPY

LIMITATION OF LIABILITY

Manufacturer warrants that this product conforms to the chemical description on the label. Buyer assumes all risks of use in handling which are at variance in any way with the directions on the label. **MANUFACTURER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. DAMAGES CAUSED BY THIS PRODUCT ARE LIMITED TO REPLACEMENT OF THE PRODUCT OR RETURN OF THE PURCHASE PRICE.**

A Prescription Treatment[®] brand insecticide from
Whitmire Research Laboratories, Inc.
· 3568 Tree Court Ind. Blvd.
St. Louis, Missouri 63122

Copyright, 1992

