

PM04 499-292

7/8/99

1 of 1

INDOOR AND OUTDOOR TREATMENTS

CARPENTER ANTS: Locate source of infestation. Inject 5-10 seconds of product into insect tunnels and cavities. Treat trails, points of entry or exits from voids such as around doors and windows. Treatment of trails on tree trunks or fences is permitted.

FOR ACTIVE AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames. Repeat as necessary.

OUTDOOR TREATMENTS

CARPENTER BEES: Locate entrance holes. Inject 5 - 10 seconds of product into entrance holes. Seal bee entrances after application.

PERIMETER TREATMENT: For crawling insects, apply PT\* 279 to cracks and crevices where insects may enter premises. Areas to be treated include, but are not limited to, weep holes, under siding, wall voids, soffits, around attic vents, cracks and crevices, openings around windows, doors and pipes. Treatment of infested fences and tree holes is permitted.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: This container may be recycled in the few but growing number of communities where steel aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the label. (DO NOT PUNCTURE!). If recycling is not available, wrap the container and discard in the trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Do not get in eyes, on skin, or on clothing. Harmful if swallowed or absorbed through the skin. Avoid breathing of spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before re-use. Avoid contamination of food, utensils, and food preparation areas. Should illness occur, get prompt medical aid. To Physician: Atropine sulfate is antidotal.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Apply this product as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard. Do not spray on plastic, painted or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers, telephones, etc.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



A Prescription Treatment\* brand insecticide from: Whitmire Micro-Gen Research Laboratories, Inc. 3568 Tree Court Industrial Blvd. St. Louis MO 63122-6682 www.wmmg.com

© 1999 Whitmire Micro-Gen Research Laboratories, Inc.

Whitmire PT\* 279

ENGAGE

Residual Injection System


For Commercial Use Only.

KILLS: Ants, Booklice, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs, Elmleaf Beetles, Grain Weevils, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders, Springtails, Ticks and Troglodermas.

FOR USE IN AND AROUND: Apartments, Homes, Hospitals, Hotels, Food storage areas, Inedible product areas of meat packing plants, Food processing plants, Motels, Nursing homes, Restaurants and other food handling establishments, Supermarkets, Transportation equipment (buses, boats, ships, trains, planes), Warehouses and other commercial and industrial buildings.

\* Do not use in aircraft cabins.

Authorized by USDA for use in feedlot areas of dead-end inspected meat and poultry plants.



ACTIVE INGREDIENTS:

Chlorpyrifos 0.5% OTHER 99.5%

EPA Reg. No. 499-292 EPA Est. No. 499-MO-1 KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED: Call a physician or poison control center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If a person is unconscious, do not give anything by mouth and do not induce vomiting.

IF INHALED: Remove patient to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: If symptoms of cholinesterase inhibition are present, atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

See side panel for additional precautionary statements

NET WEIGHT: 1 lb. 4 oz. (20 oz.)



WHITMIRE MICRO-GEN RESEARCH LABORATORIES INC. 800-777-8570

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

HOW TO USE WHITMIRE MICRO-GEN CRACK AND CREVICE\* INJECTION SYSTEM: Use Whitmire PT\* 279 only with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment. Inject PT\* 279 into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one second of product. For light infestations, move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations, move injector tip along at one linear foot per second. For dust-like voids, calculate the voids cubic area and treat at the rate of 3.0 to ten seconds per three cubic feet. Several holes may be required in long-running voids. Repeat as necessary.

APPLICATION OF PT\* 279 ENGAGE\* IN INDOOR RESIDENTIAL SETTINGS OTHER THAN AS A CRACK AND CREVICE\* TREATMENT IS NOT PERMITTED.

INDOOR TREATMENTS

CRAWLING INSECTS - Cockroaches, Dermestids, Scorpions, Spiders, Earwigs, Crickets, Sowbugs, Millipedes, Centipedes: Inject with the prescribed dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewers, floor drains and meter boxes. Repeat as necessary.

TICKS: Thoroughly treat infested and egg-laying sites near pet resting areas such as cracks and crevices, behind baseboards, door and window frames.

BOOKLICE & SILVERFISH: Inject into cracks and crevices associated with baseboards, door frames, bookcases and other harborage sites. Repeat as necessary.

ANTS: Inject cracks, crevices and voids where ants may be traveling and/or harboring, including, but not limited to, around doors, window frames, wall voids and other structural cracks and crevices.

FOR EXPOSED ADULT AND LARVAL STAGES OF GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, DRUGSTORE BEETLES, TROGODERMAS: Inject into cracks and crevices and/or voids where these insects may be harboring. Infested areas may include, but are not limited to, cracks and crevices in cabinets, pantries, wall voids and in machinery. Repeat treatment as necessary.

APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS

Food handling establishments include places other than private residences in which food is held, processed, prepared or served.

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE\* TREATMENT ONLY: Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) are also considered food areas.

Apply prescribed amounts of PT\* 279 directly into cracks and crevices or voids using injection tube or other Whitmire Micro-Gen equipment in points between different elements of construction, between equipment and floor, openings leading to voids and hollow spaces in walls, equipment legs and bases where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food, food utensils or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A CRACK AND CREVICE\* TREATMENT ARE NOT PERMITTED.