

PM 12

499-317

1 of 12

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 499-317	DATE OF ISSUANCE February 15, 1991
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	TERM OF ISSUANCE <i>Until Reregistration</i>	
NAME OF PESTICIDE PRODUCT Whitman 1-B Insecticide		
NAME AND ADDRESS OF REGISTRANT (Include ZIP code) <div style="text-align: center;"> ┌ ┐ </div> Whitman Research Laboratories, Inc. 3500 Tree Court St. Louis, MO 63122-6629 <div style="text-align: center;"> └ ┘ </div>		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>THIS PRODUCT IS CONDITIONALLY REGISTERED IN ACCORDANCE WITH FIFRA SECTION 3(c)(7)(F), PROVIDED THAT YOU:</p> <ol style="list-style-type: none"> 1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) or 4(a) when the Agency requires all registrants of similar products to submit such data. 2. Make the labeling changes listed below before you release the product for shipment: <ol style="list-style-type: none"> a. Change the label "EPA Registration No. 499-317." b. Check the directions for use for the label, delete the label directions for use for the product. 3. Submit all data required for registration/reregistration of your product when you release the product for shipment. Refer to EPA R-77-00109 for a further description of the data requirements. 		
<input checked="" type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL <i>Dennis H. Edwards</i>	DATE 2/15/91	

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Dennis H. Edwards, Jr.
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosures

3/27/90

WHITMIRE 1-6 INSECTICIDE

Controls numerous pests.

To be applied only by or under the supervision of Pest Control Operators or other trained personnel responsible for insect control programs.

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 6.42

INERT INGREDIENTS:

. 93.62

Contains 0.5 pound of chlorpyrifos per gallon.
Contains aromatic petroleum solvent.

KEEP OUT OF REACH OF CHILDREN

ACCEPTED
with **COMMENTS**
in EPA Letter Dated

WARNING

FEB 15 1991

PRECAUTIONARY STATEMENTS

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 497-307

Hazards to Humans and Domestic Animals

MAY CAUSE IRREVERSIBLE EYE DAMAGE -- HARMFUL IF SWALLOWED -- HARMFUL IF ABSORBED THROUGH SKIN -- MAY CAUSE SKIN IRRITATION

Do not get in Eyes, On Skin Or Clothing -- Wear Eye Protection -- Avoid Breathing Vapors And Spray Mist -- handle Concentrate in A Ventilated Area -- Wash Thoroughly With Soap And Water After Handling -- Remove Contaminated Clothing and Wash Before Reuse -- Keep Away From Food, Feedstuffs and Water Supplies

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Do not induce vomiting. Contains an aromatic petroleum solvent. Do not give anything by mouth to unconscious person.

If On Skin: Immediately wash with plenty of soap and water. Get medical attention.

If In Eyes: Flush eyes with plenty of water. Get medical attention immediately.

If Inhaled: Remove to fresh air. Apply artificial respiration if indicated.

Note to Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE

**Do Not Use or Store Near Heat or Open Flame
Do Not Cut or Weld Container**

Environmental Hazards

Whitacre 1-6 is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment wash waters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

473.2 ml/16 fl. oz.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Whitmire 1-6 insecticide is an emulsifiable concentrate designed for use as a spray to control various pests injurious to trees, turf and ornamental plants. The pests controlled are listed in the accompanying tables. Whitmire 1-6 is compatible with insecticides, miticides, and fungicides commonly recommended except for alkaline materials such as Bordeaux mixture and lime. A small amount of spray mixture should be prepared to check for compatibility before a large volume of spray is mixed.

Attention: Do not allow adults, children or pets on treated areas until spray has dried. Keep out of fishpools and other bodies of water. Do not treat vegetable gardens. Do not allow livestock to graze in treated areas. Do not feed treated grass cuttings (hay) or seed screenings to livestock nor use hay for livestock bedding. Do not use in poultry houses. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

PEST CONTROL INDOORS

Use Whitmire 1-6 insecticide to control pests listed in the following table by application at the recommended dosages. Use a 0.25% spray to control light infestations and a 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray, dilute Whitmire 1-6 insecticide with water.

Applications may be made within residential building, including homes and apartment buildings. Applications may also be made within nonfood areas of industrial, institutional, and commercial buildings, including hospitals, stores, manufacturing plants, and warehouses. Applications may be made within food handling establishments. This includes, but is not limited to, restaurants, grocery stores, bakeries, bottling plants, canneries, and grain mills.

Apply as a coarse, low pressure spray (20 psi or less) or with a paint brush to localized areas. Treat where insects are found or normally occur, including dark corners of rooms and closets; floor drains; cracks and crevices in walls; along and behind baseboards; beneath and behind sinks, stoves, refrigerators, refrigerator units, and cabinets; and around plumbing and other utility installations.

In food-handling establishments use only as a spot treatment. Spot treatment may encompass crack and crevice treatments by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs and bases, or which occur at points between different elements of construction, or between equipment and floors. Equipment capable of delivering a pinstream of spray should be used. Avoid applying spray to the point of runoff. Repeat treatment as needed, but not more often than once every 7 days in restaurants and similar food service establishments, or more often than once every 14 days in other types of food-handling establishments. In case of emergency (call Jack), Whitmire 1-6 insecticide may be applied after 2 days from last treatment, but emergency use should be limited to once per month. Applications of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

NOTE: Where quick "knockdown" and/or "flushing action" is desired, pyrethrins may be added to the spray mixture containing Whitmire 1-6 insecticide. For best results, use pyrethrins at 0.25% in water-based sprays. Consult labeling of pyrethrin products used to determine the amount needed to give the desired spray concentrations. Also read and observe all use and handling precautions for these products.

Do not apply water-based sprays of this product in conduits, motor housing, junction and switch boxes, or other electrical equipment because of possible shock hazard.

Do not allow spray to contact food, feedstuffs, or water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying. Do not introduce the spray into the air or allow the spray to contact food or food-contacting surfaces.

Use Whitmire 1-6 insecticide to control the pests listed below by application as a 0.25% or 0.5% spray.

Pest*	Amount of Whitmire 1-6 to Water					Specific Directions
	: Gallon of Water-based Spray					
	0.25% Spray		0.5% Spray			
	oz.	ml	oz.	ml		
Ants(1) Cockroaches (Such as: American, brown-banded, German, Oriental, Smokey brown)(2) Confused flour beetles Crickets Earwigs Firebrats Indian meal moths Mediterranean flour moths Millipedes Red flour beetles Rice weevils Saw-toothed grain beetles Silverfish Spiders	5 1/3	160	10 2/3	320		Use the low rate (0.25%) to control light infestations and the high rate (0.5%) to quickly reduce heavy infestations or for extended residual control. 1. For ants apply to <u>ant trails</u> and <u>around doors</u> and windows and wherever else the pest may find entrance. 2. A period of 4 to 7 days is normally required for maximum effect on cockroaches.
Brown dog ticks(1) Carpet beetles(2) Clover mites	5 1/3	160	10 2/3	320		1. For the control of brown dog ticks, thoroughly apply the spray as a spot treatment to infested areas, such as <u>pet beds</u> and <u>resting quarters</u> ; nearby cracks and crevices; along <u>baseboards</u> , windows and door frames, and localized areas of floor and floor coverings where these pests may be present. Old bedding should be removed and replaced with clean, fresh bedding after treatment. Do not treat pets with this product. 2. For the control of carpet beetles, thoroughly apply the spray as a spot treatment along <u>baseboards</u> and edges of carpeting, under carpeting, rugs and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

*Numbers in parentheses refer to Specific Directions.

TURF AND OTHER OUTDOOR USES

Use Whitmire 1-6 to control the pests listed in the following table by application at the recommended dosages and in accordance with the directions given below. Dilute Whitmire 1-6 in water and apply as a coarse, low-pressure spray using suitable application equipment. Use one gallon of diluted spray per 1,000 sq. ft. of area. Thoroughly water immediately after treatment to wash the insecticide into the turf, except as noted. For best results, the lawn should be moist at time of treatment. Spray when pest first appear; retreat when needed.

Pest*	Amount of Whitmire 1-6 in water to make 1 gal.	Specific Directions
Ticks (Such as American dog, cattle fever, Gulf coast, lone star)(1) brown)(2)	2 fl. oz.	1. For control of ticks, infesting corrals, stockyard holding pens and other outdoor livestock areas, wet in corrals and pens, especially in corners, feeders and shelters and other areas likely to serve as harborage sites for ticks that have removed themselves from their host. Spray surfaces to be treated, but do not create excessive runoff. NOTE: This application is intended as a preventive only. Do not use as a direct spray on livestock care to avoid creating mist or drift which may contact with feed, water or livestock. Livestock not be allowed in treated areas until spray has dried.
Ants (including foraging fire ants) Armyworm (Such as: Beech, Fall, Yellow-striped) Centipedes Chiggers(1)X Chinch bugs Clover mites Cutworms Crickets Earwigs Flory skipper Fire ants (mounds)(2)X Fleas Gnats Grasshoppers Greenbugs June beetle Leafhoppers Lucerne moth Millipedes Mites (Such as: Clover, Perennial grass, Winter grain) Mosquitoes(3)X Sod webworm(4)(10-15 mites)X Sawflies Springtails Ticks(1)	4 fl. oz. 1999 sq. ft.	1. Use Whitmire 1-6 insecticide for area control of and Chiggers infesting non-cropland areas such as sides, footpaths, and trails, picnic and camping parks and other recreational areas where these are present and create a nuisance or a possible health problem. Do not allow use of treated area during application or until spray has dried. Apply Whitmire 1-6 in water at the rate of 1 pint per acre (equivalent to 4 fl. oz. per 1,700 sq. ft.) using a hydraulic sprayer, mist applicator, insect sprayer, or other suitable hand or power-operated equipment. Treat low underbrush, grassy areas, and ground surface and debris using enough spray to obtain thorough coverage, usually 40 to 100 gal per acre. 2. For individual fire ant mounds, apply Whitmire 1-6 in a drench, diluted at the rate of 2 fl. oz. (60% 1 gallon of water. Gently sprinkle a total of 1 gallon of the diluted emulsion over the mound each mound using a gentle spray (such as a spritz can). Thoroughly wet mound and surrounding area a 4 ft. diameter (12 sq. ft.). For best results apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants cause migration, reducing product effectiveness. 3. Mosquitoes coming to rest on areas treated for control of turf pests will be killed for varying periods time after treatment, depending on exposure of areas to weathering conditions. 4. For sod webworms delay watering or mowing the treated area for 12 to 14 hours after treatment.
European crane fly Turfgrass weevil (<i>Hypodamia</i>)(1)	2 fl. oz. 12 fl. oz.	1. Spray suspected problem areas in mid-April and a mid-May or as recommended by your local Agricultural Extension Service Specialist.
White grubs (Such as: Black turfgrass stem-borer, European chafer, Southern masked chafer, and Japanese beetle larvae)(1)	12 to 16 fl. oz.	1. For white grubs, spray when grubs are young and feeding near the soil surface, usually during June and August or as recommended by your local Agricultural Extension Service Specialist. Immediately after spraying, irrigate the treated area with 1/2 to 1 of water to wash the insecticide into the soil.
Bluegrass billbugs(1)	8 to 16 fl. oz.	1. For bluegrass billbugs spray early in the season when they first appear. Retreat as needed.

*Numbers in parentheses refer to Specific Directions.

ORNAMENTALS (OUTDOOR AND NURSERY)

Use Whitmire 1-6 insecticide to treat flowers, shrubs, evergreens, vines, shade and flowering trees found to be infested with pests listed in the following table. Dilute Whitmire 1-6 with water according to directions given in the table and apply using suitable hand or power-operated spray equipment in a manner to provide complete and uniform coverage. For best results apply a coarse spray to thoroughly wet both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 20 day intervals, if needed. Consult your State Agricultural Experiment Station or Extension Service Specialist for application timing and other specific use information.

Note: Ornamentals: Environmental factors have significant effects on phytotoxic expression. Whitmire 1-6 has been tested on numerous ornamental plants without causing serious phytotoxicity. However, do not use on azaleas, camellias, poinsettias, rose bushes, or variegated ivy because of possible injury to these plants.

Pest*	Amount of Whitmire 1-6 in water to make 1 gal.	Specific Directions
<p>Adelgids (Such as: Cooley and Eastern spruce galls, Pine bark)</p> <p>Ants (including <u>foraging fire ants</u>)</p> <p>Aphids (Such as: Apple, Chrysanthemum, Cottonwood, Elm leaf, Peach, Rose, Spirea, Woolly)</p> <p>Armyworms (Such as: Fall, Yellowstriped)</p> <p>Bagworms(1)</p> <p>Barkbeetle bugs</p> <p>Cankerworms</p> <p>Catalpa sphinx</p> <p>Chiggers</p> <p>Citrus mealybugs</p> <p>Cockroaches (Such as: American, Brown-headed, German, Oriental, Smoky brown)</p> <p>Elm sawfly</p> <p>Fall webworm</p> <p>Grasshoppers</p> <p>Green fruitworm</p> <p>Herbivores</p> <p>Jackpine budworm</p> <p>Juniper webworm</p> <p>Katydid</p> <p>Leaf bugs</p> <p>Leafhoppers</p> <p>Leafrollers(2)</p> <p>Maple leafcutter(3)</p> <p>Mites (Such as: Clover, Red spider, Southern red)(4)</p> <p>Slender caterpillar</p> <p>Orange tortrix</p> <p>Periodical cicada</p> <p>Plant bugs</p> <p>Poplar tentmaker</p> <p>Psyllids</p> <p>Pine caterpillar</p> <p>Pine chafer</p> <p>Scavflies, exposed (Such as: Pine, Pin oak, Redheaded)</p> <p>Sawflies</p> <p>Spittlebugs</p> <p>Spruce budworm</p> <p>Tent caterpillars (Such as: Eastern, Forest, Western)</p> <p>Thornbug</p> <p>Walrus caterpillar</p> <p>Western spruce budworm</p> <p>Whiteflies</p> <p><u>Yellow-necked caterpillar</u></p>	<p>2/3 fl. oz.</p>	<p>1. For bagworms treat when insects are in the crawler stage.</p> <p>2. For effective control of leafrollers spray should be applied before leaves are tightly rolled.</p> <p>3. For effective control of maple leafcutter on maple trees apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production.</p> <p>4. For effective control of spider mites when large numbers of eggs are present, apply a second spray 1 to 3 days in the South or 7 to 10 days in the North after initial treatment to control newly-hatched nymphs.</p>

*Number in parentheses refer to Specific Directions.

Pest*	Amount of Whitacre 1-6 in water - 8 oz. 1 gal.	Specific Directions
<p>Aphids (Beet)</p> <p>Beetles (Paler rose, Native elm bark)(1)</p> <p>Cecidians</p> <p>Leafhoppers ✓</p> <p>Manzanita weevils</p> <p>Mealybugs ✓</p> <p>Miner weevils</p> <p>Moths (Such as: Brownail, Cypress tip, Douglas fir tussock, European pine shoot, Gypsy (2), Holly bud, Nantucket pine tip, Podara, Pitch pine tip, Subtropical pine tip, Tussock)</p> <p>Curculionids (Such as: California, Orange-striped, Redhumped)</p> <p>Redhumped caterpillar</p> <p>Thrips (exposed)</p> <p>Weevils (Such as: Blackvine (3), Yellow Pine, Pine reproduction)</p> <p>Beetles (Such as: Cottonwood leaf (1), Elm leaf, Pine, Willow leaf)</p>	<p>1 1/3 fl. oz.</p>	<ol style="list-style-type: none"> 1. To reduce twig and branch feeding by bark beetles applications should be made in the spring or early summer. 2. To kill migrating and invading gypsy moth larvae treat trunks and foliage. 3. Blackvine weevils are night feeders. Late afternoon spraying will give control in some areas.
<p>Beetles(1): Clearwing moths (Such as: Ash, Dogwood, Lesser peachtree, Lilac, Oak, Rhododendron), Metallic wood (Such as: Spruce Birch, Flatheaded appletree, Twined chestnut), Longhorned beetles (Such as: Locust, Red oak)</p> <p>Leafminers</p> <p>Needleminers (Such as: Jeffrey pine, Lodgepole pine, Spruce)</p> <p>Scale crawlers (Such as: Cottonyushion, Cottonyaspic, Eucynema, Fletcher, Florida wax, Golden oak, Semicircular, Louisiana, Magnolia, Oak horned, Oystershell, Pine needle, San Jose, Tea, White Birch)</p>	<p>2 2/3 fl. oz.</p>	<ol style="list-style-type: none"> 1. For borers, apply Whitacre 1-6 to the trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service Specialist for proper time to treat. Apply uniformly as a coarse low-pressure spray.
<p>Borer (Such as: Cottonwood, Peachtree)(1)</p>	<p>8 fl. oz.</p>	<ol style="list-style-type: none"> 1. For psachrae borers apply Whitacre 1-6 in water to flowering trees and shrubs of the genus Prunus as a trunk spray before newly-hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs.
<p>Beetles (Such as: Ambrosia, Anobidae, Black turpentine, Cottonwood leaf, Elm leaf, European elm bark, Flea, Fuller, Vase, Japanese, June, Native elm bark, Pine, Willow leaf)(1)</p>	<p>11 1/2 fl. oz.</p>	<ol style="list-style-type: none"> 1. For preventive treatment apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infected trees. <p>For remedial treatment apply the spray to the main trunk of infected trees when damage occurs but before adult beetles begin to emerge.</p>

* Number in parentheses refer to specific directions.

11 of 12

DORMANT SPRAY (TREE AND FOREST PESTS)

Use Whitmire 1-6 insecticide as a dormant or delayed dormant spray at the rates indicated to control the listed insects. Spray the entire tree by application to runoff using suitable ground spray equipment. Use the higher dosage of Whitmire 1-6 insecticide for severe infestations. Use oil as recommended by your State Agricultural Experiment Station or Extension Service Specialist.

Restrictions: Make only one application during the dormant season. Do not allow meat or dairy animals to graze in treated areas.

Pests	Amount of Whitmire 1-6 in water to make 1 gal.	Specific Directions
Aphids (Such as: Honey plum, Eony apple, Veinly apple) Borer (Such as: Peach twig) Cutworms (Such as: Climbing) Leafrollers (Such as: Pandemic) Pear psylla adults Plant bugs Scale (Such as: San Jose)	2/3 to 1 1/3 fl. oz.	Mix with 1 to 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water.

FRUIT TREES AND NUTS

Use Whitmire 1-6 insecticide at the dosages indicated by application as a foliar spray to control pests listed in the following table. Mix the required dosage of Whitmire 1-6 in water. Spray for thorough and complete coverage of the foliage, using suitable hand- or conventional power-operated spray equipment. Treat when pests first appear or in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service specialist for specific use information. Do not allow livestock to graze in treated areas. Attention: In California do not work with or contact trees within two days after treatment.

Pests	Amount of Whitmire 1-6 in water to make 1 gal.	Specific Directions
Apples: Apple aphid Apple maggot Codling moth European apple sawfly Eyespotted bud moth Fruitree leafroller Green fruitworm Lesser appleworm Obliquebanded leafroller Orange tortrix Oriental fruit moth Pandemic leafroller Plum curculio Redbanded leafroller Rusy apple aphid San Jose scale Tarnished plant bug Tufted apple budmoth Variegated leafroller Veinly apple aphid	1 fl. oz.	Make no more than 3 applications per season. Do not apply last treatment within 15 days before harvest or apply last two treatments closer than 21 days apart.
Almonds: Navel orangeworm Peach twig borer San Jose scale	3 1/3 fl. oz.	Make no more than 3 applications per season to almonds and filberts and no more than 2 applications per season to walnuts. Do not apply within 15 days before harvest.
Walnuts: Codling moth Walnut scale	3 1/3 fl. oz.	
Pecans: Black pecan aphid Hickory shootworms Pecan leaf scorch mite Pecan nut casebearer Pecan weevil Phylloxera spp.	2 2/3 fl. oz.	Make no more than 3 application per season. Do not apply within 15 days of harvest. Do not allow livestock to graze in treated orchards.

13 of 12

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 100°F for extended periods of time. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: METAL DRUM Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Container Disposal: PLASTIC Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.