

PM03

499-323

7/9/88

1/1

PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. Do not induce vomiting. Vomiting may cause aspiration pneumonia.
IF INHALED: Remove patient to fresh air. Apply artificial respiration if indicated.
IF ON SKIN: Remove contaminated clothing and wash affected skin areas with soap and water. Get medical attention if irritation persists. If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Do not remain in treated area and ventilate the area after treatment is completed. Avoid breathing of vapor. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using and before smoking or eating. Do not contaminate water, food, or foodstuffs. Remove all pets and birds. Cover fish tanks or remove before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Apply this product only as specified on this label.

FOOD/FEED AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS

Before using this pesticide compound, exposed food products and packaging materials must be removed from the room or protected. After using this compound, equipment and utensils must be washed with an effective cleaning compound and thoroughly rinsed with potable water before operations are resumed. Before food product may be placed in the area where the material is being used, the area should be sufficiently free of odor to prevent product contamination.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers, etc.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in a cool area away from heat or open flame.
PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal state and local procedures.
CONTAINER DISPOSAL: This container may be recycled in the few but growing number of communities where steel aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the label (DO NOT PUNCTURE!). If recycling is not available, wrap the container and discard in the trash. For best results product should be used when can temperature is above 60°F. If temperature is below 60°F, store at room temperature until a temperature of above 60°F is reached.

A Prescription Treatment® brand Insecticide from:
Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682

© 1997 Whitmire Micro-Gen Research Laboratories, Inc.

Contains no CFC's or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



CLEAR ZONE

FARM FLY SPRAY



KILLS AND REPELS Flies • Gnats • Mosquitoes

Fleas, Ants, Cluster Flies, Cockroaches, Crickets, Deer Flies, Face Flies, Gnats, Hornets, Horn Flies, Horse Flies, House Flies, Lice, Mosquitoes, Mud Daubers, Small Flying Moths, Spiders, Stable Flies, and Wasps.

FOR USE IN AND AROUND, BUT NOT LIMITED TO:
Dairy Farms (including Milk House, Milk Parlor, Loading Sheds, and Holding Lot), Hog Operations, Horse Stables, Poultry Operations, Beef Cattle Operations, Kennel Farms, Restaurants and other Food/Feed Handling Establishments.
Authorized for use by USDA in federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:	
Pyrethrins	0.5%
Piperonyl Butoxide, Technical	4.0%
INERT INGREDIENTS:	95.5%
*Equivalent to 3.2% (Efficacy) (1,1-Dichloro-2,2-Diethyl Ether and 0.3% Related Compounds)	

EPA Reg. No. 499-323 EPA Est. No. 499-MO-1
KEEP OUT OF REACH OF CHILDREN
CAUTION
See side panel for additional precautionary statements.

NET WEIGHT: 20 oz.



Whitmire Micro-Gen
Research Laboratories, Inc.
800-777-5570

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

DAIRY FARM USE: Milking Parlor and Milk Room - Close all windows and doors. Direct fog upward and in all directions at a rate of 1 to 3 seconds per 1000 cu. ft. Keep room closed for 15 minutes after treatment. Open and ventilate before reoccupying. Cover milking utensils and milk to prevent contamination from spray and dead falling insects.

ANIMAL USE (Do not spray sick animals): From approximately 2 foot distance, thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animals' eyes. Spot treat withers, shoulders and back where saliva accumulated from head tossing.

STANCHION BARN USE: Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double dosage. Repeat application daily or as necessary.

HORSE BARN AND STABLES: Close all windows and doors. Spray at a rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double the dosage. For animal application, lightly mist over the backs of horses and ponies from a distance of at least two feet. Repeat application daily or as necessary.

POULTRY OPERATIONS: Use only in empty poultry houses (when birds are not present). Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double the dosage. Treat daily or as necessary.

HOG OPERATIONS: In hog houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes after application. Ventilate the area after treatment is completed. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly toward animals' face or eyes. Repeat application daily or as necessary.

MUD DAUBER AND WASP NESTS: Spray mist directly on insects and their nests from approximately 2 ft. distance. Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Spray into hiding and breeding places containing as many insects as possible.

FLEAS: General Volumetric Treatment. Hold aerosol 36" above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq Ft. making sure that all floor areas are contacted. Change litter before application. Keep area closed for 15 minutes. Remove animals before spraying. Open and ventilate before reoccupying. Repeat treatment after 7 days or as necessary.

KENNELS AND CATTERIES: Remove animals before spraying. Ventilate after treatment is completed.

FLYING INSECTS (Cluster Fly, Flies, Fruit Flies, Gnats, Mosquitoes, Small Flying Moths): GENERAL VOLUMETRIC TREATMENT: Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cu. ft. Disperse in all locations. Keep area closed for 15 min. Open and ventilate before reoccupying. Repeat treatment as necessary.

FOR EXPOSED ADULT AND LARVAL STAGES OF - Carpet Beetles, Cigarette Beetles, Confused Flour Beetles, Drug Store Beetles, Grain Mites, Granary Weevils, Red Flour Beetles, Rice Weevils, and Sawtoothed Grain Beetles: Contact as many insects as possible. Inject into cracks and crevices of pantries, cabinets, food processing and handling equipment and other places where insects harbor. Apply around cartons, and containers, and other areas where these insects tend to congregate.

GENERAL VOLUMETRIC TREATMENT: Apply 20 seconds per 1000 cu. ft. of space. Repeat treatment weekly or as insects appear. Infested stored products should be fumigated or treated by other effective methods using an approved product labeled for this purpose.

OUTDOOR GENERAL APPLICATION: Flies, Gnats and Mosquitoes in open areas near building and in campgrounds. Best results are obtained when wind speed is five (5) MPH or less. Apply at a rate of 60-80 seconds per acre. Spray in wide swath across area to be treated. Allow spray drift to penetrate dense foliage. Repeat treatment as necessary.

ACCEPTED

JUL 09 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 499-323