	499-323	•	08/ 1999	1/2	
CEPA Environmental Protection Ager Washington, DC 20460		Form Appro	Registration Amendment Other	OPP Identifier Number	
Application for Pesticide - Section I					
1. Company/Product Number		2. EPA Product Manager		3. Proposed Classification	
4. Company/Product (Name)		Masson Johnson		None Restricted	
5. Name and Address of Applicant (Include ZIP Code) Whitmire Mimo-Gen Research laboratories, Inc 3568 Tree Court Industrial Blod. 84. Longe, Mo 63120-6682 Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name			
Section - II					
Amendment - Explain below. Resubmission in response to Agency letter Notification - Explain below.	dated	Final printed labels in response to Agency letter dated "Me Too" Application. Other - Explain below.			
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Deletion of Use site: "Restaurants/Food Handling Establishments. Inadvertently submitted as notification, with other minor revisions, on 7/22/99. Resubmitting as an amendment to be compliant with PRNs 98-10 & 91-1.					
Section - III					
1. Material This Product Will Be Packaged In:					
Child-Resistant Packaging Yes* No * Certification must be submitted Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per II "Ye	r Soluble Packaging Yes No Is" No. per age wgt containe	P P	ntainer Metal Mestic Glass Paper Other (Specify)	
3. Location of Net Contents Information 4. Size(s) Retail Container 5. Location of Label Directions On Label On Label On Labeling accompanying product					
6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled Other					
Section - IV					
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Steven Weiss		Title Telephone No. (Include Area Code) Recistation Sorcialist (636) 861-4292			
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment o. /Stamped)					
2. Signature		3. Title Registration Specialist 5. Date 9/21/99			
4. Typod Name Steven We135		9/21/90	<u> </u>		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool area away from heat or open flame

PESTICIDE DISPOSAL: Pesticide that cannot be used according to tabet instructions must be disposed of according to applicable Federal, state and local procedures.

CONTAINER DISPOSAL: This container may be recycled in the few but growing number of communities where steel aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the labet (DO NOT PUNCTURE!). If recycling is not available, wrap the container and discard in the trash.

PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately, Gastric lavage is indicated if material was taken internally. Do not induce vomiting. Vomitling may cause aspiration pneumonia.

IF INHALED: Remove patient to fresh air. Apply artificial respira-

IF ON SKIN: Remove contaminated clothing and wash affected skin areas with soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water. Get medical attention it irritation persists.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmlut if swallowed or absorbed through skin. Do not remain in treated area and ventitate the area after treatment is completed. Avoid breathing of vapor. Avoid contact with skin, eyes, or cothing. Wash thoroughly after using and before smoking or eating. Do not contaminate water, food, or foodstuffs. Remove all pets and birds. Cover tish tanks or remove before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to tish, birds, and other wildlife. Oo not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Apply this product only as specified on this tabel.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces or direction any electronic equipment such as radios, televisions, computers, etc.

For best results product should be used when can lemperature is above 60°F. It temperature is below 60°F, store at room temperature until a temperature of above 60°F is reached.

Contains no CFC's or other ozone depleting substances. Federal regulations prohibit CFC propelants in aerosols.



A Prescription Treatment® brand Insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc. 3568 Tree Court Industrial Blvd. St. Louis MO 63122-6682

www.wmmg.com

§ 1999 Whilmire Micro-Gen Research Laboratories, Inc.



Whitmire PT 20

Dairy and Farm Insect Fogger and Repellant



KILLS AND REPELS Flies • Gnats • Mosquitoes

Fleas, Arts, Cluster Flies, Cockroaches, Crickets, Deer Flies, Face Flies, Gnats, Hornets, Horn Flies, Horse Flies, House Flies, Lice, Mosquitoes, Mud Daubers, Small Flying Moths, Spiders, Stable Flies, Wasps, Carpet Beetles, Centipodes, Cigarette Beetles, Clover Mites, Confused Flour Beetles, Drug Store Beetles, Fruit Flies, Grain Mites, Granary Weevils, Mitigaters, Mud Daubers, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles and Sow Bugs.

FOR USE IN AND AROUND, BUT NOT LIMITED TO: Dairy Farms (including Milk House, Milk Parlor Leaf ed Sheds, and Holding Lot), Hog Operations, Horse Statzles, Poultry Operations, Beef Cattle Operations, Kennels, Farms, Campgrounds, Cattlenes and Horse Barns.

Authorized for use by USDA in federally inspected meat and poultry plants

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
	95.5%
*Enumeration States (factorises to profession	

EPA Reg. No. 499-323 EPA Est. No. 499-MO-1 KEEP OUT OF REACH OF CHILDREN

CAUTIONSee side panel for additional precautionary statements.

NET WEIGHT: 20 oz.



TIFICATION

OCT 8 1999

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

DAIRY FARM USE: Milking Parlor and Milk Room - Close all windows and doors. Direct tog upward and in all directions at a rate of 1 to 3 seconds per 1000 cu. ft. Keep room closed for 15 minutes after treatment. Open and ventilate before reoccupying. Cover milking utensils and milk to prevent contamination from spray and dead falling insects.

ANIMAL USE (Do not spray sick animals): From approximately 2 foot distance, thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animals's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

STANCHION BARN USE: Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes following application. Ventifate area after treatment is compilee. If area cannot be closed, double the dosage. Repeal application daily or as necessary.

HORSE BARNS AND STABLES: Close all windows and doors. Spray at a rate of 1 to 2 seconds per 1600 cu. ft. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double the dosage. For animal application, lightly mist over the backs of horses and ponies from a distance of at least two feet. Repeat application daily or as necessary.

POULTRY OPERATIONS: Use only in emply poultry houses (when birds are not present). Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. It. Keep area closed for 15 minutes following application. Ventilate area after treatment is complete. If area cannot be closed, double the dosage. Treat daily or as necessary.

HOG OPERATIONS: In hog houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cu. ft. Keep area closed for 15 minutes after application. Ventilate the area after treatment is completed. If area cannot be closed, double the dosage. For animal application, direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly toward animal's face or eyes. Pepeal application daily or as necessary.

MUD DAUBER AND WASP NESTS: Spray mist directly on insects and their nests from approximately 2 ft. distance. Contacted insects will fly away from top. Applications should be made in late evening when insects are all rest. Spray into hiding and breeding places contacting as many insects as possible.

FLEAS: General Volumetric Treatment: Hold aerosol 36" above floor and direct spray loward floor and lower walls at a rate of 10 seconds per 100 sq. leet making sure that all floor areas are contacted. Change litter before application. Keep area closed for 15 minutes. Remove animals before spraying. Open and ventitate before reoccupying. Repeat treatment after 7 days or as necessary.

KENNELS AND CATTERIES: Remove animals before spraying. Ventilate after treatment is completed.

FLYING INSECTS (Cluster Fly, Flies, Fruit Flies, Gnats, Mosquiloes, Small Flying Moths): GENERAL VOLUMETRIC TREATMENT: Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cu. fl. Disperse in all locations. Keep area closed for 15 min. Open and ventilate before reoccupying. Repeat treatment as necessary.

FOR EXPOSED ADULT AND LARVAL STAGES OF - Carpet Beelles, Cigarette Beelles, Confused Flour Beelles, Drug Store Beetles, Grain Miles, Granary Weevils, Red Flour Beetles, Rice Weevils, and Sawtoothed Grain Beelles; Contact as many Insects_as possible. Inject into cracks and crevices of partires, cabinets and other places where insects harbor. Apply around cartons, and conticess_and other areas where these insects tend to congregate. GMERAL Vol. UMETRIC TREATMENT: Apply 20 seconds per 1000 but it. of space. Repeat treatment weekly or as insects apper. Helstel Jored products should be fumigated or treated by other effective* methods using an approved product intended for this purpose.

OUTDOOR GROUND APPLICATION: Flies, Gnats and Mosquiloes in open areas near building and in campgrounds. Best results are obtained when wind speed is fire (5) MPH or less, Apply at a rate of 60-86 seconds per acre. Spray in wide swaths across area to be treated. "Allow spray driff to pendrate dense foliage. Repeal treatment as accessary.