

499-413

2-14-2001

OUTDOOR BUILDING TREATMENTS

FOR USE ON: Outside surfaces of Manufacturing, Industrial and Food Processing Plants, Warehouses... SUBTERRANEAN ANTS SUCH AS ARGENTINE ANTS and HARVESTER ANTS: Inject the nest at 3-6 sites at a rate of 5-10 seconds per site...

RESTRICTED USE PESTICIDE For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

WHITMIRE TC 100 INTERN®

Residual Crack & Crevice® and Wood Injection system

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

HOW TO USE WHITMIRE MICRO-GEN CRACK AND CREVICE® INJECTION SYSTEM: Use Whitmire TC 100 INTERN® with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment...

INDOOR TREATMENTS

FOR USE IN: Food and Non-Food Areas of Manufacturing, Industrial and Food Processing Plants, Warehouses, Ship Holds and Railroad Boxcars... CRAWLING INSECTS - Cockroaches, Dermestids, Scorpions, Spiders, Earwigs, Crickets, Sowbugs, Millipedes, Centipedes...

OTHER OUTDOOR TREATMENTS

TERMITE SPECIFIC INFORMATION/INSTRUCTIONS: This treatment is intended for localized termite infestations only... FIRE ANTS: Inject the nest at 3-6 sites at a rate of 5-10 seconds per site...

KILLS: Ants, Booklice, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs...

FOR USE IN AND AROUND: Apartments, Homes, Hospitals, Hotels, Food Storage Areas, Food Processing Plants, Motels, Nurseries, Homes, Restaurants and Other Food Handling Establishments, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Planes) and Warehouses...

ACTIVE INGREDIENTS: Chlorpyrifos 0.5% OTHER INGREDIENTS: 99.5% Contains petroleum distillates.

EPA Reg. No. 499-413 EPA Est. No. 499-MO-1 KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person. IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

STORAGE & DISPOSAL

Store in a cool dry area away from heat or open flame and inaccessible to children. PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. ENVIRONMENTAL HAZARDS This product is toxic to fish, birds, and other wildlife.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.

A PRESCRIPTION TREATMENT® brand insecticide from: Whitmire Micro-Gen Research Laboratories, Inc. 3568 Tree Court, Industrial Blvd. St. Louis, MO 63122-6662



NET WEIGHT: 1 lb. 1.5 oz. (17.5 oz.)

WHITMIRE MICRO-GEN RESEARCH LABORATORIES INC. 800-777-8570

ACCEPTED FEB 14 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 499-413

APPLICATIONS IN FOOD HANDLING PROCESSING ESTABLISHMENTS

FOR USE IN: Food and Non-Food Areas of Food Processing Plants, FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE® TREATMENT ONLY: Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), processing - edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups).

INDOOR AND OUTDOOR TREATMENTS

WOOD DESTROYING INSECTS FOR USE ON: Outdoor Fence Posts and Utility Poles in Non-Residential Settings as well as Chromatals in Industrial Plant Sites only. LOCALIZED TREATMENT OF WOOD-INFESTING INSECTS: Beetles and Spores (such as the following families: Anobiidae, Eosinichidae, Euprosopidae, Cerambycidae, Lyctidae, Oedemeridae, Sesiidae), Carpenter Ants, Carpenter Bees, Termites and Wood Wasps.

TERMITES AND CARPENTER ANTS LIVING INSIDE TREES: Determine the highest and lowest elevation of the colony. Drill holes into colony at highest elevation and inject 5 to 10 seconds. Do the same for lowest elevation. AERIAL TERMITE CARTON IN TREES: Drill or punch hole into center of carton. Inject for 5 to 10 seconds. If carton is greater than 12 inches across, use another injection site.

