

499-419

1/11/2001

1/3

RESTRICTED USE PESTICIDE
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

DURATION

PT 275 MC Microencapsulated Dursban

Liquid Concentrate

ACTIVE INGREDIENT:
Chlorpyrifos 20.0%
OTHER INGREDIENTS: 80.0%

EPA Reg. No. 499-419 EPA Est. No. 499-MO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See supplemental label for additional Precautionary Statements and Directions for Use.

NET CONTENTS: 1 Pint



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.
800-777-8570

PT 275 MC Container Label v3.2.4 (Rev. 1/01)

ACCEPTED
JAN 11 2001
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
Registered under
EPA Reg. No. 499-419

ground size, avoiding unnecessary runoff. Allow 7 to 10 days to achieve maximum effectiveness. This product has shown residual control of fire ants in excess of 30 days. If retreatment is necessary, limit applications to once per every 14 days.

HORTICULTURAL USE

Commercial Greenhouses, Interiorscapes and Landscapes

CONTROLS: Ants, Aphids, Beetles, Centipedes, Cockroaches, Crickets, Flea Beetles, Flies, Gnats, Leafhoppers, Leaf Miners, Mealybugs, Scale, Scorpions, Spring Bugs and Thrips.

MIXING AND DILUTION INSTRUCTIONS:

Heavy Infestation: 0.5 fl. oz. DURATION™ PT® 275 MC per gallon of water.

Preventative and Maintenance: 0.25 fl. oz. DURATION™ PT® 275 MC per gallon of water.

Spray upper and lower foliage surfaces of plants uniformly.

SPECIAL NOTICE: This product has demonstrated excellent plant safety; however, all varieties of the plants listed have not been tested. When treating large numbers of plants of a single variety, spray a few plants and observe for phytotoxicity prior to full scale application.

FOR USE INDOORS AND OUTDOORS ON:

Bedding plants (including): impatiens, marigolds, petunias, geraniums, garden mums, verbena, New Guinea Impatiens, and zinnia.

Cut Flowers (including): roses, chrysanthemums, carnations, snapdragons, and orchids.

Flowering hanging baskets (including): Lobelia and Lantana.

Foliage (including): Boston fern, dracaena, ficus, schefflera, and philodendron.

Non-flowering plants (including): poinsettia, chrysanthemum, forest coleus, lilies, African violet, geranium, hibiscus, kalanchoe, begonia, gloxinia, cyclamen, cineraria, coleocalyx, and croton.

Trees & shrubs (including): azalea, barberry, cotoneaster, euonymus, holly, hyl, juniper, oak, pine, madrodam, rose, spirea, spruce, viburnum, and yew.

USE IN AND AROUND LIVESTOCK HOUSING

MAY BE USED IN AND ON OUTSIDE SURFACES OF OTHER BUILDINGS

FOR USE IN AND AROUND: Livestock Housing and Housing Areas (such as Hog Barns, Horse Barns, Empty Chicken Houses, Rabbit Hutches, Dairy Areas, Milkrooms, Calf Hutches, Calving Pens and Parlors).

Do not apply to lactating or food animals. Do not contaminate milk, food or drinking water. Remove all animals prior to making application to housing. Do not apply or allow insecticide to drift onto animals. Do not allow animals to enter treated areas until surfaces are dry.

For residual control of house flies and stable flies on outdoor surfaces and in animal areas such as dog kennels, hog barns, horse barns, empty chicken houses, rabbit hutches, dairy areas, milkrooms, calf hutches, calving pens and parlors. Treat normal fly resting areas such as ceilings, screens, upper corners of the area, around windows and doors, light fixtures and other surfaces known to attract flies. Apply as a coarse wet spray. Using a mixture of 2.0 to 5 fl. oz. of DURATION™ PT® 275 MC per gallon of water, apply at a rate of 1 gallon diluted product per 1,000 sq. ft. of surface. For best results initiate treatments at the beginning of the fly season and repeat during periods of heavy fly activity.

For control of other pests and to help prevent infestations of buildings, treat a band of soil 6 to 10 feet wide around and adjacent to the building. Also treat the building foundation and sides where pests are active and may find entrance. Treat other insect entry points such as doors, windows, under siding and eaves of structure. DURATION™ PT® 275 MC may be applied in compressed air, backpack or power spray equipment. Depending on type of equipment used to apply product and landscaping of perimeter, use 4.2 to 4.175 ounces per 1,000 sq. ft. Alternate directions are to use 4.5 to 26.875 fl. oz. of product per 50 gallons of water applied at 10 gallons of dilution per 1000 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container only, preferably in a locked storage area. Do not store in a manner where cross contamination with other pesticides, fertilizer, food or feed could occur. If spilled during storage or handling, soak up spillage with absorbent material and dispose of in accordance with Pesticide Disposal instructions listed below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse container, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Do not apply to humans, their clothing or bedding. Do not allow children or pets to contact treated surfaces until spray has dried. Do not contaminate food or use on household pets. Cover fish tanks prior to application.

Avoid contamination of food, feedstuffs, or water supply. Do not contaminate food preparation surfaces, kitchen utensils, dishes, or feed storage containers. All food contact surfaces and cooking utensils should be covered during treatment or thoroughly washed with an effective cleaning compound followed by a potable water rinse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, fish and aquatic invertebrates. Birds feeding in treated areas may be killed. Clean up spilled product to reduce exposure to wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not spray where electrical short circuits might result, such as wall outlets, conduits, motors, switches, etc. Do not spray directly into any electronic equipment, such as radio, televisions, computers, telephones, etc.

RESTRICTED USE PESTICIDE

To be sold for retail sale to and applied use only by commercial Certified Applicators responsible for insect control programs or persons under the direct supervision of such commercial applicators and only for those uses covered by the Certified Applicator's certification.

DURATION®

PT® 275 MC Microencapsulated Dursban

Liquid Concentrate

Contains Dursban® Insecticide

FOR USE IN AND AROUND: Apartments, Campgrounds, Commercial Buildings, Food and Non-Food Areas of Manufacturing, Industrial Plants and Food Processing Plants, Food Storage Areas, Hobby Commercial Greenhouses, Homes, Hospitals, Motels, Food Service Establishments, Interiorscapes, Kennels, Landscapes, Motels, Nursing Homes, Office Buildings, Resorts, Restaurants, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Cargo Holds of Ships, Railroad Boxcars, Commercial Trucks, Trains, Trucks, Planes), Utilities, Warehouses, Commercial and Industrial Buildings, Dog Kennels, Hog barns, Horse barns, Empty chicken houses, Rabbit hutches, Dairy areas, Milkrooms, Calf hutches and Calving pens and parlors.

CONTROLS: Angoumois Grain Moths, Aphids, Ants, Bees, Beetles, Boxelder Bugs, Caddisflies, Carpet Beetles, Carpenter Ants, Carpenter Bees, Centipedes, Cigarette Beetles, Clover Mites, Cluster Flies, Cockroaches, Confused Flour Beetle, Crickets, Darkling Beetles, Domestic Beetles, Dried Fruit Beetles, Drugstore Beetles, Earwigs, Elmleaf Beetles, Fire Ants, Firebrats, Flea Beetles, Fleas, Flies, Gnats, Grain Mites, Indian Meal Moths, Leafhoppers, Leaf Miner, Mealworms, Mediterranean Flour Moths, Millipedes, Moths, Northern Fowl Mites, Psocids, Red Flour Beetles, Saw-Toothed Grain Beetles, Scale, Scorpions, Silverfish, Sowbugs, Soiders, Spring Bugs, Termites, Trogloderma, Wasps, Weevils (Granary Weevils and Rice Weevils) and Yellowjackets.

*Dursban is a registered trademark of Dow/Banco.

ACTIVE INGREDIENTS:

Chlorpyrifos 20.0%

OTHER INGREDIENTS:

..... 80.0%

EPA REG. NO. 499-419 EPA EST. NO. 499-MQ-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to do so. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

See page 4 for additional Precautionary Statements.

A Prescription Treatment® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.

3568 Tree Court Industrial Blvd.

St. Louis, MO 63122-6682

www.wmrg.com

© 2000 Whitmire Micro-Gen Research Laboratories, Inc.



WHITMIRE MICRO-GEN

RESEARCH LABORATORIES INC.

800-777-8570



DURATION™ PT® 275 MC

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

GENERAL INFORMATION: DURATION™ PT® 275 MC is a flowable microencapsulated concentrate containing chlorpyrifos. DURATION™ PT® 275 MC may be sprayed on any surface which will not be damaged or stained by water. Heavy applications may leave a visible deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

MIXING INSTRUCTIONS: Thoroughly clean spray equipment before using this product. Shake concentrate well before diluting. When diluting, first add approximately one-half of the water to the spray tank and then add the proper amount of this product. Add the rest of the water and agitate the sprayer thoroughly. Agitate sprayer occasionally during use to ensure even coverage. Shake or re-agitate sprayer if spraying is interrupted or if dilution is left in sprayer overnight. When emptying the bottle, triple rinse with water and add the rinse to the spray tank. If spray screens are used, they should be 50 mesh or larger.

In situations where conventional application methods have not or are not likely to provide adequate coverage, foam generating equipment or similar machines can be used to provide a continuous coverage. Use dry foam inside wall voids to avoid any water damage or electrical short circuits.

PEST CONTROL-INDOORS INDOOR TREATMENTS

FOR USE IN: Food and Non-Food Areas of Manufacturing, Industrial, and Food Processing Plants; Warehouses; Ship Holds; Railroad Boxcars; Livestock Housing and Holding Areas (such as Hog Barns, Horse Barns, Empty Chicken Houses, Rabbit Hutches, Dairy Areas, Milkrooms, Calf Hutches, Calving Pens and Partens); Dilute DURATION™ PT® 275 MC with water using the spray dilution chart.

SPRAY DILUTION CHART: Dilute DURATION™ PT® 275 MC with water as specified below.

Amount of Finished Spray	0.25% Spray	0.5% Spray
1 gallon	1.5 fl. oz.	3 fl. oz.
3 gallons	4.5 fl. oz.	9 fl. oz.
5 gallons	7.5 fl. oz.	15 fl. oz.
10 gallons	15 fl. oz.	30 fl. oz.

The higher rate should be used for clean-outs, for longer residual control or for high insect infestations.

DURATION™ PT® 275 MC should be used as a crack and crevice residual product where insects may be hiding, living and breeding and/or for selected spot treatment.

CENTIPEDES, COCKROACHES, CRICKETS, DERMESTITID BEETLES, EARWIGS, FIREBRATS, MILLIPEDES, PSOCIDS, SILVERFISH, SOWBUGS, SPIDERS: Make spot treatment and/or crack and crevice injections into areas where insects are normally found for example cracks and crevices, void areas, moist areas, openings around pipes, sinks and under refrigerators. Repeat as necessary.

FLIES: Make spot applications of this product on window frames, screens, trash cans, ceilings and upper wall surfaces, and other surfaces where flies rest in the dark at night or frequent during the day.

CARPET BEETLES (DERMESTITIDS): Thoroughly apply the spray as a spot treatment along edges of carpet, under carpet edge (if necessary), under furniture, base and throw rugs. Use as a crack and crevice application in closets and other shelving where infested items are stored. Do not treat clothing with this product. Repeat application as necessary.

TICKS AND FLEAS: Thoroughly apply as a spot treatment in infested and egg laying areas in pet beds, pet resting quarters, nearby cracks and crevices and behind baseboards. For fleas, make sure to treat window frames. Replace old pet bedding with clean fresh bedding before treatment. Do not treat pets with this product. Repeat application as necessary.

FOR EXPOSED ADULT AND LARVAL STAGES OF GRAIN MITES, STORED PRODUCT BEETLES, GRAIN WEEVILS, AND STORED PRODUCT MOTHS: Inject into cracks and crevices and/or voids where these insects may be harboring. Infested areas include, but are not limited to, cracks and crevices in cabinets, pantries, wall voids and in machinery. Repeat treatment as necessary.

ANTS: Inject cracks, crevices and voids where ants may be traveling and/or harboring, around doors, window frames, and other structural cracks, crevices and voids. Spot treatment of trails and feeding areas may also be made.

HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks, crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices in weep holes, wall voids, around window and door frames, and under siding. Repeat application as necessary. Treat wall area around entry points on exterior where insects congregate.

CARPENTER ANTS: Make spot and/or crack and crevice applications to any trails, inside or outside around doors and windows, and other places where ants enter premises. Areas which may be treated include but are not limited to attics, crawl spaces, ceilings, garages, soffits, wall voids, weep holes, and other void areas. Where possible, treat ant nest or infested wood directly. Repeat application as necessary.

WOOD DESTROYING INSECTS - Localized treatment of wood-infesting insects: Beetles and Borers (such as the following families: Anobiidae, Bostrichidae, Buprestidae, Cerambycidae, Lyidae, Odontomidae, Sesiidae), Termites and Wood Wasps. Locate suspected infestation and/or insect tunnels and cavities. Drill pilot holes into insect tunnels, gallery and inaccessible closed voids. Drill holes 8 to 10 inches apart to insure thorough coverage. Inject 5 to 10 seconds per hole. Do not contact electrical wiring, plumbing, etc. when drilling. Use the 0.5% or 1.0% rate. (For termite use, rate limited to 0.5%.)

APPLICATIONS IN FOOD PROCESSING ESTABLISHMENTS

This product may be applied within residential and non-residential buildings or structures and within industrial, institutional and commercial buildings. Applications can be made in food service establishments such as restaurants, cafeterias, taverns, hotels, motels, mobile homes, school and institutional dining areas, dormitories, mobile canteen, vending machines, groceries and markets.

FOR USE IN: Food and Non-Food Areas of Manufacturing, Industrial and Food Processing Plants. Applications can also be made in Non-food areas of food manufacturing and food processing establishments including garbage rooms, lavatories, drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling). Application in food areas of food service establishments in processing plants is limited to spot and/or crack and crevice treatments.

Treat where insects are found or naturally occur. Application of a 0.25% solution is suitable for control of light pest infestations (i.e. maintenance applications). Application of a 0.5% solution provides residual control of heavy infestations (i.e. clean-out applications) and where extended residual control is desired. A period of 4 to 7 days is normally required to observe maximum effectiveness on insect populations; therefore, limit retreatments to no more often than once per every 7 days (14 days in food areas of food service establishments). Apply spray solution in such a

manner as to minimize spray drift to non-target surfaces. For spot treatments, an individual spot should not exceed 2 sq. ft.

PEST CONTROL-INDOORS AND OUTDOORS

PEST CONTROL-OUTDOORS

OUTDOOR BUILDING TREATMENTS

FOR USE ON OUTSIDE SURFACES OF: Manufacturing, Industrial, and Food Processing Plants; Warehouses.

PERIMETER SPRAY: To help prevent infestations of boulongs, treat a band of soil 6 to 10 feet wide around and adjacent to the building. Also treat the building foundation to a height of 2 to 3 feet and where pests are active and may find entrance. Treat other insect entry points around doors, windows, under siding and eaves of structure, under decks and on fences. This product may be applied in compressed air equipment or power spray equipment. Depending on type of equipment used to apply product and landscaping of perimeter, use 0.5 to 1.25 fl. oz. per 1,000 sq. ft. Alternate directions are to use 5 to 8.75 fl. oz. of product per 50 gallons of water applied at 10 gallons of dilution per 1,000 sq. ft.

ANTS, CRICKETS, COCKROACHES AND OTHER CRAWLING INSECTS: For treatment of localized infestations of these insects in outdoor areas which are landscaped, planted, mulched or unmaintained, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat applications as infestation warrants and as reinfestations occur.

FLIES AND MOTHS: Spray outside surfaces of screens, doors and window frames or wherever these insects enter. Also treat surfaces around light fixtures on porches, in garages, and other places where insects light or congregate. Treat normal fly resting areas on ceilings, upper corners of the area. Repeat as necessary. Spray inside and outside surfaces of dumpsters and other trash holding containers. Spray trash corrals and other trash storage areas. Spray so as to lightly moisten the target surface. For best results, begin treatments at the start of the fly season and repeat weekly during periods of heavy fly activity.

HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks, crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices in weep holes, wall voids, around window and door frames, and under siding. Repeat application as necessary. Treat wall area around entry points on exterior where insects congregate.

CARPENTER ANTS: Make spot and/or crack and crevice applications to any trails, inside or outside around doors and windows, and other places where ants enter premises. Areas which may be treated include but are not limited to attics, crawl spaces, ceilings, garages, soffits, wall voids, weep holes, and other void areas. Where possible, treat ant nest or infested wood directly. Repeat application as necessary.

FLEAS: Use 1 to 2 1/2 fl. oz. of this product to treat 1,000 sq. ft. Treat animal runs, kennel, pet resting areas, shaded areas and other areas where fleas are likely to be found. This product may be tank mixed with approved insect growth regulators.

BEES, WASPS AND YELLOW JACKETS: As a preventative or remedial treatment to help prevent these insects from building nests or workers from congregating on structures follow perimeter treatment directions under outdoor section of label. Pay particular attention to soffit and eave areas.

FOR TREATMENT OF YELLOWJACKET OR WASP NESTS: Liberally treat the wasp nest. Treat inside the nest for yellow jackets and the surrounding area. For treatment of nests inside wall voids inject into the nest sufficient quantity to treat the complete nest. More than one point of application may be required. Most applications should require no more than half a gallon of diluted material. Pay particular attention to what the water may contact such as electrical wiring. Do not use where there is a shock hazard.

THIS PRODUCT DOES NOT PROVIDE INSTANT KNOCKDOWN. THE BEST TIME TO TREAT IS LATE IN THE DAY. COLONY SHOULD BE DEAD IN 2-3 DAYS. DEPENDING ON THE SITUATION PROTECTIVE EQUIPMENT FOR THE APPLICATOR MAY BE REQUIRED.

CARPENTER BEES: Locate entrance holes. Treat galleries through entrance holes. Seal bee entrances after application.

WOOD DESTROYING INSECTS - Localized treatment of wood-infesting insects: Beetles and Borers (such as the following families: Anobiidae, Bostrichidae, Buprestidae, Cerambycidae, Lyidae, Odontomidae, Sesiidae), Termites and Wood Wasps. Locate suspected infestation and/or insect tunnels and cavities. Drill pilot holes into insect tunnels, gallery and inaccessible closed voids. Drill holes 8 to 10 inches apart to insure thorough coverage. Inject 5 to 10 seconds per hole. Do not contact electrical wiring, plumbing, etc. when drilling. Use the 0.5% or 1.0% rate. (For termite use, rate limited to 0.5%.)

SCORPIONS: Remove accumulations of lumber, framework and other materials which serve as harborage sites. Before applying firewood or lumber apply this product as a residual spray to surfaces immediately below such materials. Band treatments may also be helpful in reducing pests immigrating from surrounding areas.

SEWER MANHOLE SHAFTS: To control mosquitoes in sewers, add 1/2 fluid ounce of product per each gallon of water and apply approximately 0.5 gallons of spray solution per each 8 vertical feet of manhole shaft. Avoid application directly to drainage water. Most efficient coverage is achieved, but not limited to, by spraying the vertical walls of the sewer manhole shaft using a power sprayer fitted with a long extended wand tip with a 90 degree spray nozzle.

NOTE: Do not discharge spray or unneeded material directly into the sewer system. Do not apply product within 36 hours of predicted heavy rainfall.

NOTE: Persons applying this product in the manhole must wear chemical resistant gloves, a respirator approved by MICH for organic chemical vapors, goggles and long sleeved clothing.

DURATION™ PT® 275 MC can be used as a residual spray on outdoor surfaces and/or as a crack, crevice and void treatment of buildings, porches, window frames, eaves, patios, garages, refuse dumps and other areas where pests congregate or enter premises. Repeat treatment as necessary to maintain effectiveness. Use the 0.25% or 0.5% rate.

DURATION™ PT® 275 MC can be used as a residual spray to control pests in outdoor void areas such as meterboxes and manholes. Repeat treatment as necessary.

OTHER OUTDOOR TREATMENTS

TERMITES: This treatment is intended for localized termite infestations only. For use outside structures only. This product cannot be used for spot and local treatments to existing structures after 12-31-02. Spot treatments are allowed only where there is clear evidence that reinfestation or barrier disruption has occurred. Spot treatments not to exceed 25% of the amount required to treat the entire structure at the label rate. Areas treated with 0.5% or less must be inspected annually for signs of reinfestation. To inhibit termite infestations treat exposed tubing, cracks or cold joints in concrete or other structural interstices. Treatment of exposed surfaces should be done to cover thoroughly but excess run-off should be avoided. This type of application is intended to be a supplemental treatment for control of drywood and above ground subterranean termites and other wood-infesting insects. It is not intended to be a substitute for soil treatment or for extensive infestation of drywood or other wood-infesting insects. Use a 0.5% rate, 3 fl. oz. of concentrate per gallon of water. To treat for severe infestations or for treatments under slabs in basements and buildings with insufficient crawlspace, use a registered termiticide.

In situations where conventional application methods have not or are not likely to provide adequate coverage, foam generating equipment or similar machines can be used to provide a continuous barrier treatment of filled porches, chimney bases, soil under slabs and treatment of wall voids are examples where foam applicators may be useful.

FIRE ANT MOUNDS: For individual mounds apply this product as a mound drench at a rate of 1.0 fluid ounce per each 2 gallons of water. Satisfactory results can be expected by applying 2 gallons of dilution per mound. Adjust volume according to

3/3