	03,
Please read instructions on reverse before completing form	n. - /

03/10/2005

\$epa	Environmenta Washi	United States I Protection Age Ington, DC 20460	-	Registrati Amendme X Other	
	······································	Application for	· · · · · · · · · · · · · · · · · · ·	······································	· · · · · · · · · · · · · · · · · · ·
1. Company/Product Numbe	499-427		2. EPA Product Man Richard Gebken	agor	3. Proposed Classification
4. Company/Product (Name Whitmire PT 3-6-10C Ac)	onc	PM#	10	None Restricted
5. Name and Address of Ap Whitmire Micro-Gen Re 3568 Tree Court Indus St. Louis, MO 63122-6	pplicant <i>(Include ZIP Co</i> esearch Labs, Inc trial Blvd		(b)(i), my product i to:		e with FIFRA Section 3(c)(3) I in composition and labeling
Check if this	s is a new address		Product Name		
		Sec	ction - II		
Amendment - Explain Resubmission in resp X Notification - Explain	oonse to Agency letter	datad	Agency lett	opplication.	MAR 1 0 2005
"This notification is consistent with th ormula of this product. I understand PR Notice 98-10 and 40 CFR 152.46	that it is a violation of 18 U.S.	C. Sec. 1001 to willfully make a ion of FIFRA and I may be sub	any faise statement to EPA, if	urther understand that if this	labeling or the confidential statement of notification is not consistent with the lerms and 14 of FIFRA "
. Material This Product Wil	l Be Packaged In:				
hild-Resistant Packeging Yes No Certification must e submitted	Unit Packaging Ves No If "Yes" Unit Packaging wgt.	No. per If "Yes	Soluble Packaging Yes No s" No. per ge wgt container	P G P	ntainer Aetal Iastic ilass aper Ither (Specify)
o soonnittoo	· · · · · · · · · · · · · · · · · · ·				
1 1		4. Size(s) Retail Contai	iner	5. Location of Label [Directions
Labei C	Container	4. Size(s) Retail Contai	iner	[]	Directions
Labei C	Container	Lithograph Papar glued Stenciled		[]	Directions
Label C Manner in Which Label is	Container Affixed to Product	Lithograph Paper glued Stencied ~ Sect	(Other 10n - IV	·	
Label C Manner in Which Label is Contact Point <i>(Complete</i> ame	Container Affixed to Product items directly below fo	Lithograph Peper glued Stencifed ~ Sect or identification of indivi- Title	(Other 10n - IV	f necessary, to proce	
Label C Manner in Which Label is Contact Point <i>(Camplete</i> ame H. Wayne Moran	antainer Affixed to Product <i>items directly below fo</i> ments I have made on y knowlinglly false or r	Lithograph Peper glued Stencifed ~ Sect or identification of indivi- Title	Lion - IV idual to be contacted, i Reg. and R&D	f necessery, to proce Tel 	ss this application.) ephone Nr. (Include Area Code) CC-861-4218 6. Dece Application Received • . (Stamped)
Label C Manner in Which Label is Contact Point <i>(Complete</i> ame H. Wayne Moran I certify that the stater I acknowledge that an both under applicable Signature	antainer Affixed to Product <i>items directly below fo</i> ments I have made on y knowlinglly false or r	Lithograph Peper glued Stenciled ~ Sect or identification of indiv. Title Dir., R Certification this form and ell attach misleading statement m 3. Title	Lion - IV idual to be contacted, i Reg. and R&D	f necessary, to proce. Tel 	ss this application.) ephone Nr. (Include Area Code) CC-861-4218 6. Dece Application Received • . (Stamped)
. Manner in Which Label is . Contact Point <i>(Complete</i> ame H. Wayne Moran I certify that the stater I acknowledge that an both under applicable Signature	Container Affixed to Product <i>items directly below fo</i> ments I have made on y knowlinglly faise or r law.	Lithograph Peper glued Stenciled ~ Sect or identification of indiv. Title Dir., R Certification this form and ell attach misleading statement m 3. Title	Lion - IV idual to be contacted, i Reg. and R&D ments therato are true may be punishable by fir	f necessary, to proce. Tel 	ss this application.) ephone Nr. (Include Area Code) CC-861-4218 6. Dece Application Received • . (Stamped)

WHITMIRE PT[®] 3-6-10C AERO-CIDE[®]

[FRONT PANEL]

WHITMIRE PT 3-6-10C AEROCIDE[®]

[ALTERNATE BRAND NAME #1: "Prescription Treatment[®] brand ULD[®] HydroPy-300[®]" DESCRIPTOR: "Pyrethrin Concentrate"] [ALTERNATE BRAND NAME #2: "Prescription Treatment[®] brand ULD[®] HydroPy-300[®] CF" DESCRIPTOR: "Water-Based Contact Insecticide"] NOTIFICATION

MAR 1 0 2005

PYRETHRIN CONCENTRATE

KILLS:

Angoumois Grain Moths, Ants, Boxelder Bugs, Cadelles, Cheese Mites, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Earwigs, Fruit Flies, Gnats, Granary Weevils, House Flies, Indian Meal Moths, Mosquitoes, Rice Weevils, Spiders, Tobacco Moths and Wasps.

FOR USE IN AND AROUND:

Apartments, Cafeterias, Dormatories, Food Service and Preparation Areas, Grocery and Convenience Stores, Hotels, Health Care Facilities, Motels, Office Buildings, Restaurants, Schools and Universities. See page 4 for additional insects and use sites.

ACTIVE INGREDIENTS:

Pyrethrins, <u>a botanical insecticide</u>	3.0%
Piperonyl butoxide, technical*	
n-Octyl bicycloheptene dicarboximide**	
OTHER INGREDIENTS: * Equivalent to 4.8% (butylcarbityl)(6-propylpiperonyl) ether and 1.2% related compounds. ** MGK-264 Insecticide Synergist	<u>81.0%</u> TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

See complete label for additional Precautionary Statements.

EPA Reg. No. 499-427 • EPA Est. No. 499-MO-1

ULTRA LOW DOSAGE THROUGH MICRON GENERATION

NET CONTENTS:



WHITMIRE MICRO-GEN RESEARCH LABORATORIES, INC. 800-777-8570

[ADDITIONAL LABELING]

3/10

PRECAUTIONARY STATEMENTS

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 1-800-225-3320 for emergency medical treatment information.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

GENERAL RESTRICTIONS

Do not apply directly to food. Do not apply as a space or ULV spray while food processing is underway. Cover or remove any food in the treatment area before application. Cover all food processing surfaces and equipment or wash with an effective cleaning compound followed by a potable water rinse.

Food processing operations do not have to be stopped while applying as a wet spray with care and in accordance with directions and cautions above in establishments which do NOT operate under Federal meat, poultry, shell egg grading and egg products inspection programs.

In the home, cover all food processing surfaces, dishes and utensils in the treatment area or thoroughly wash before use.


Remove pets, birds, and cover fish aquariums and turn off air pump motor before spraying. In health care facilities or schools, remove patients or students prior to treatment. Thoroughly ventilate treated area for two hours before reoccupying treated areas.

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

Page 2 of 8 V4.0.7

WHITMIRE PT[®] 3-6-10C AERO-CIDE[®]

DILUTION RATES



Use undiluted in mechanical sprayers which produce particles of aerosol size (80% of droplets less than 30 microns), or diluted in pump sprayers or automated misting systems. Use the 0.125% rate for flying insects only. Use the 0.25% rate for maintenance. Use the 0.5% rate for clean outs or high insect infestations. Use the 0.05% rate only for automated misting or fogging applications outdoors.

Amount of Concentrate per Gallon of Water*
2 fl. oz.
5 1/3 fl. oz.
10 2/3 fl. oz.
21 1/3 fl. oz.
42 2/3 fl. oz.

* Do not dilute with spray oil.

NOTE: 1/3 fl. oz. = 2 tsp and 1/2 fl. oz. = 1 tbsp.

APPLICATION INSTRUCTIONS

SPACE SPRAY (INDOORS): Apply using a mechanical aerosol generator which creates particles of aerosol size (80% of droplets less than 30 microns). Prior to application, close all doors and windows, shut off all ventilating equipment and extinguish all flames in the area to be treated. Do not remain in treated areas after application. Leave room(s) closed for at least one (1) hour. Ventilate thoroughly before re-entry.

CRACK AND CREVICE® TREATMENT: Use diluted or undiluted. Inject PT 3-6-10C into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. Apply in such a manner to limit dripping and run-off on structural surfaces.

VOID INJECTION: Use diluted or undiluted. Use to flush and kill insects harboring in closed or inaccessible voids. Calculate the cubic area of the void and treat at the rate of five to ten seconds per three cubic feet. Several holes may be required in long-running voids. Apply in such a manner to limit dripping and run-off on structural surfaces.

SPOT TREATMENT AND GENERAL SURFACE SPRAY: Use diluted or undiluted. Hold spray nozzle 18 to 24 inches from areas to be treated. Treat surfaces until moist, but in such a manner to limit dripping and run-off.

INDOOR TREATMENTS

FLYING INSECTS (such as Bees, Cluster Flies, Flies, Gnats, Hornets, Mosquitoes, Moths, Small Flying Moths, Wasps, Webbing Clothes Moths and Yellow Jackets): Apply as a space spray or general surface spray. *SPACE SPRAY:* apply undiluted at the rate of 0.5 fl. oz. of concentrate per 1,000 cu. ft. of space. Point nozzle upward and direct spray mist to all parts of the room. *GENERAL SURFACE SPRAY:* use the 0.125% or 0.25% rate and direct spray toward upper corners of room especially around windows and other light sources which attract these insects. Do not remain in treated area(s) after application. Ventilate thoroughly before re-entry.

CRAWLING INSECTS (such as Ants, Booklice, Carpet Beetles, Centipedes, Clothes Moths, Clover Mites, Cockroaches, Crickets, Dermestid Beetles, Earwigs, Firebrats, Fleas, Millipedes, Scorpions, Silverfish, Sowbugs and Spiders): Apply as general surface spray, Crack and Crevice treatment or void injection. Use the 0.25% or 0.50% rate. *GENERAL SURFACE SPRAY*: Apply as a coarse, wet spray on walls, floors and other surfaces. Treat food storage areas, handling areas and unloading and processing areas. *CRACK AND CREVICE TREATMENT AND VOID INJECTION*: Apply to cracks and crevices or structural voids in these places as well as beneath sinks and inside and around the base of food conveying, processing and handling equipment.

STORED PRODUCT PESTS (such as Angoumois Grain Moths, Caddies, Carpet Beetles, Chocolate Moths, Cigarette Beetles, Dermestid Beetles, Dried Fruit Beetles, Drug Store Beetles, Flour Beetles, Grain Mites, Granary Weevils, Indian Meal Moths, Meal Worms, Mediterranean Flour Moths, Psocids, Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Trogaderma and Weevils): Apply as a space spray or a general surface spray. Clean all food preparation areas thoroughly after treatment and before food is exposed. Use only in areas where food is not exposed. *SPACE SPRAY*: apply undiluted at the rate of 0.5 fl. oz. of concentrate per 1,000 cu. ft. of space. Direct treatment towared upper corners of room and ceiling. *GENERAL SURFACE SPRAY*: use the 0.25% or 0.50% rate. Direct output into corners and under pallets, hitting as many insects as possible.

BED BUGS, LICE AND TICKS: Apply as a space spray, spot treatment or Crack and Crevice treatment. First, thoroughly clean air matresses and springs. *SPACE SPRAY:* apply undiluted at the rate of 0.5 fl. oz. of concentrate per 1,000 cu. ft. of space. *CRACK AND CREVICE TREATMENT OR VOID INJECTION:* use the 0.25% or 0.50% rate and apply to cracks and crevices or structural voids around baseboards, floorboards, bedboards and walls. *SPOT TREATMENT:* Take bed apart. Treat joints and channels if hollow, such as square or round tubing. See that the interior framework is treated. Treat mattresses, especially tufts, folds and edges. Also treat other areas where bed bugs may be hiding.

FLEAS: Apply as a *GENERAL SURFACE SPRAY* undiluted or using the 0.25% or 0.50% rates. Direct treatment onto pet resting areas and floor around these areas. Treat pet bedding the same way. Also treat entire inside surface of dog houses. Do not allow pets to contact treated areas until spray has dried. Remove or cover animal feed before spraying. To control the source of flea infestations, pets inhabiting the treated premises should be treated with a product registered for application to pets.

DARKLING BEETLES: After emptying poultry houses, apply as a space spray or general surface spray. Treat litter before cleaning. Keep closed for one hour, remove litter and retreat all surfaces. Ventilate for 24 hours before replacing poultry. *SPACE SPRAY*: apply undiluted at the rate of 0.25 fl. oz. of concentrate per 1,000 cu. ft. of space. *GENERAL SURFACE SPRAY*: use the 0.25% or 0.50% rate.

INDOOR AND OUTDOOR TREATMENTS

BOXELDER BUGS, CARPENTER ANTS, CLUSTER FLIES AND ELMLEAF BEETLES: Apply as a *CRACK AND CREVICE TREATMENT* or as a void injection to voids and other areas where these insects are found. Use undiluted or at the 0.25% or 0.50% rates.

FLIES, GNATS, HORN FLIES, HOUSE FLIES, STABLE FLIES (AROUND LIVESTOCK): For fast knockdown of flying insects, apply as a *SPACE SPRAY* undiluted or using the 0.25% or 0.50% rates. Direct fog upward in all directions. Ventilate the area after treatment is completed. Remove or cover animal feed before spraying. Do not apply to lactating or food animals. Do not contaminate milk, food or drinking water.

OUTDOOR TREATMENTS

MOSQUITOES, FLIES AND GNATS: Dilute to 0.050% - 0.125% when applying with mechanical, electrical, fogging or automated misting equipment, or when applying as a *GENERAL SURFACE SPRAY*. To control these pests in open areas near homes, buildings and in campgrounds, treat shrubbery and vegetation and other areas where these pests may rest. Structural surfaces, shrubbery and vegetation around stagnant pools, marshy areas, ponds and shorelines may be treated. Treatment of vegetation in water from land is also permitted by allowing wind to move product into vegetation, but application of this product directly to any body of water is prohibited.

AUTOMATED MISTING SYSTEMS: Automatic spraying systems must not be programmed to release pesticide where people, pets, food or feed are directly exposed. A legible copy of the label must be securely attached to the equipment pesticide holding container or on the outside of the equipment timer.

TRUCK MOUNTED EQUIPMENT: When using truck mounted equipment, spray at a dosage rate of fifteen (15) oz. per minute from a location up-wind of the areas where insects are to be controlled. Keep speed of vehicle to approximately five (5) MPH. Figure application travel path for a convergance of a three hundred (300) foot swath. Repeat application as required for control.

HORNETS, WASPS AND YELLOW JACKETS: Apply as a *GENERAL SURFACE SPRAY* using the 0.25% or 0.50% rate. Application should be made in late evening when insects are at rest. Spray liberally into hiding and nesting places, contacting as many insects as possible.

6/10

FOR USE IN:

Hobby Greenhouses, Interiorscapes and Landscapes.

KILLS:

Aphids, Armyworms, Bees, Beetles, Bugs (Boxelder, Lace and Plant), Caterpillars (Cabbage Worms, Rose Slugs, Hornworms), Centipedes, Clover Mites, Crickets, Earwigs, Firebrats, Flies, Fruitworm, Fungus Gnats, Fungus Gnat Larvae, Grasshoppers, Leafhoppers, Leafminers, Leaf Rollers, Loopers, Mealybugs, Millipedes, Mosquitoes, Psylla, Rose Aphids, Scale Insects, Scorpions, Sowbugs, Spiders, Spider Mites, Thrips, Wasps, Whiteflies and Yellow Jackets.

APPLICATION INSTRUCTIONS

This product is designed for diluted use (0.05% - 0.25%) in sprayers. DO NOT DILUTE PRODUCT WITH SPRAY OIL. Treat upper and lower foliage surfaces of plants uniformly.

SPRAY DILUTION CHART

Pyrethrin % Desired	Amount of Concentrate per Gallon of Water*
0.050%	2 fl. oz.
0.125%	5 1/3 fl. oz.
0.250%	10 2/3 fl. oz.

* Do not dilute with spray oil.

PLANT SAFETY NOTICE: This product has demonstrated excellent plant safety; however, all varieties of the plants listed nave not been tested. When treating large numbers of plants of a single variety, treat a few plants and observe for phytotoxicity prior to full scale application.

Direct treatment to some open blooms may cause petal drop.

FOR USE INDOORS AND OUTDOORS ON, BUT NOT LIMITED TO:

Bedding plants (such as): impatiens, marigolds, petunias, geraniums, and garden mums.
Cut Flowers (such as): roses, chrysanthemums, carnations, snapdragons, and orchids.
Flowering hanging baskets (such as): fuchsia, lantana, New Guinea impatiens, ivy, and geraniums.
Foliage (such as): Boston fern, dracaena, ficus, schefflera, and philodendron.
Potted Flowering plants (such as): poinsettia, chrysanthemums, florist's azaleas, lilies, African violets, geraniums, hibiscus, kalanchoes, and begonias.
Ornamentals (such as): azalea, barberry, cotoneaster, euonymus, holly, juniper, oak, pine, rhododendron, roses, spiraea, spruce, viburnum and yew.

Do not treat vegetables.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container in secured dry storage area. Do not store above 100° F for extended periods of time. Storage below 32° F may cause freezing.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Empty container by using the product in accordance with the label directions. Do not reuse this container! Place empty container in trash or offer for recycling if available. If container is partly filled, call your local solid waste agency or call 1-800-CLEANUP for disposal instructions. NEVER PLACE UNUSED PRODUCT DOWN ANY INDOOR OR OUTDOOR DRAIN!

In addition to sites listed on the front panel, PT 3-6-10C is acceptable for use in the following sites:

INDOORS: Animal areas & quarters Apartments **Auditoriums** Bakeries Barns **Beverage Plants** Boats **Bottling Plants** Brandy Storage Warehouses **Buildings** Buses Cabins Calf Hutches Canneries **Cattle Barns** Churches Closets Commercial and Industrial Buildings **Communication Centers** Condominiums **Correctional Facilities** Dairies **Dairy & Livestock Barns** Drains & Sewers **Dried Fruit Processing Plants Dried Fruit Warehouses** Drugstores **Dumpsters Dwellings** Egg Processing Plants Factories Flour Mills Food Handling Establishments Food Processing Plants Food Storage Areas Freight Containers Fruit Packing Sheds Garbage Cans **Garbage Compactors** Goat Houses Grain Elevators Grain Handling Equipment Grain Mills Granaries Hobby Greenhouses Hog Operations

Homes Horse Barns Horse Stables Hospices Hospitals Hotels Industrial Installations Institutions Interiorscapes Jails Kennels Laboratories Libraries Livestock Barns Loafing Sheds Locker Rooms Mausoleums Meat Packing Plants Milk Rooms Milking Parlors Mills Missions Mobile Homes Motels Museums Mushroom Houses Mushroom Processing Plants Nursing Homes **Office Buildings** Passenger Rail Cars Peanut Processing Plants Peanut Storage Warehouses Playgrounds **Poultry Houses** Poultry Operations **Poultry Processing Plants** Prisons **Public Buildings** Rabbit Houses **Rabbit Processing Plants** Railroad Cars **Recreational Vehicles Research Animal Areas** Restaurants Restrooms Rice Mills Schools Shade Houses

Shipholds Ships Stable Bins Stables Stored Food in Multi-walled Paper or Cloth Bags Stored Seed Warehouses Stores Supermarkets Swine Houses **Textile Mills & Warehouses** Theaters Tobacco Plants **Tobacco Warehouses** Townhouses Trains **Transportation Equipment Truck Trailers** Trucks Utilities **Utility Rooms** Voids (Attics, Walls, Ceilings) Warehouses Wine Storage Warehouses Wineries Zoos

OUTDOORS:

Campgrounds Corrals **Drive-in Restaurants Drive-in Theaters Dumpsters** Feedlots Garbage Dumps **Golf Courses** Junkyards Kennels Landscapes Lath Houses Manure Piles Parks Plantscapes **Recreational Areas** Sewers Swineyards Zoos

v4.0.8, 1/31/05

PT 3-6-10C is also effective against the following insects when used according to label directions:

FLYING INSECTS: Aphids **Barn Flies** Bed Bugs Bees Black Flies Blow Flies Bottle Flies **Cheese Skippers** Cluster Flies Deer Flies Face Flies Flies Fruit Flies Fungus Gnats Gnats Horn Flies Hornets Horse Flies House Flies Mosquitoes Mud Daubers Mushroom Flies Skipper Flies Stable Flies Thrips Vinegar Flies Wasps Whiteflies **Yellow Jackets**

GRAIN BORERS:

Larger Grain Borers Lesser Grain Borers

CRAWLING INSECTS:

Ants Bed Bugs Booklice **Boxelder Bugs** Cabbage Loopers Cabbage Worms Carpenter Ants Centipedes Chiggers Chrysanthemum Leaf Miners Cockroaches Crickets Earwigs Fire Ants Firebrats Grasshoppers Mealy Bugs

Millipedes Palmetto Bugs Pill Bugs Psocids Scorpions Silverfish Sowbugs Spring Tails Vegetable Leaf Miners

م ب

FLEAS AND TICKS:

Brown Dog Ticks Fleas Sheep Ticks Spinos Ear Ticks Ticks

SPIDERS & MITES:

Cheese Mites Clover Mites Grain Mites Red Spider Mites Spider Mites Spiders

LICE: Lice Mounting Stages of Yellow Lice & Sucking Blue Lice

WORMS:

Dark Mealworms Lesser Mealworms Mealworms Yellow Mealworms

BEETLES:

American Black Flour Beetles Black Carpet Beetles Black Fungus Beetles **Broadhorned Flour Beetles** Brown Spider Beetles Cadelles Carpet Béetles Catorama Beetles Cereal Beetles Cigarette Beetles Confused Flour Beetles Corn Sap Beetles Darkling Beetles Depressed Flour Beetles Dermestid Beetles **Dried Fruit Beetles**

Drugstore Beetles Elmleaf Beetles Flat Grain Beetles Flour Beetles Foreign Grain Beetles Fungus Beetles Ground Beetles Hairy Fungus Beetles Hairy Spider Beetles Hide Beetles Khapra Beetles Large Black Flour Beetles Longheaded Flour Beetles Merchants Grain Beetles Mexican Grain Beetles Murmidius Beetles **Red Flour Beetles** Red Horned Grain Beetles **Rusty Grain Beetles** Saw-Toothed Grain Beetles Slenderhorned Flour Beetles **Smalleyed Flour Beetles** Spider Beetles Squarenecked Grain Beetles Trogoderma Beetles **Two Banded Beetles** White Marked Spider Beetles

WEEVILS:

Bean Weevils Broadnosed Grain Weevils Coffee Bean Weevils Granary Weevils Maize Weevils Rice Weevils

MOTHS:

Almond Moths Angoumois Grain Moths Chocolate Moths Clothes Moths Cocoa Bean Moths European Grain Moths Indian Meal Moths Mediterranean Flour Moths Moths Pink Scavenger Caterpillars Rice Moths Small Flying Moths Tobacco Moths Webbing Clothes Moths •ما

• 5

ы

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc. 3568 Tree Court Industrial Blvd. St. Louis MO 63122-6682

www.wmmg.com

© 2004 Whitmire Micro-Gen Research Laboratories, Inc.



WHITMIRE MICRO-GEN RESEARCH LABORATORIES, INC.

January 31, 2005

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504C) U. S. Environmental Protection Agency Room 266A, Crystal Mall 2 1801 S. Bell Street Arlington, VA 22202

RE: NOTIFICATION to add "a botanical insecticide" as a descriptor to Active Ingredient In compliance with PRN 98-10, Sec II, K, 2 Whitmire PT® 3-6-10C Aero-Cide Pyrethrin Concentrate; EPA Reg. No. 499-427

Dear Sir / Madam:

Enclosed in support f this action is a completed application and one copy of the revised labeling for the referenced product. The descriptor ", a botanical insecticide" has been added to the Active Ingredient statement as allowed by PRN 98-10, Section II, K, 2.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Also enclosed is a self-addressed, "No-Postage Necessary" postcard that, when filled out, will indicate the status of this notification. Please take a moment to fill in the appropriate information and drop it in the mail for our records. Your assistance in this regard is certainly appreciated

Thank you for your assistance in this matter. Please don't hesitate to contact me at (636) 861-4218 or via electronic mail at <u>wayne.moran@wmmg.com</u> should you have any questions or require additional information concerning this action.

Sincerely,

Whitmire Micro-Gen Research Laboratories, Inc.

N. akeyn Mc

H. Wayne Moran Director, Regulatory and Research & Development



3568 Tree Court Industrial Boulevard St. Louis, MO 63122-6682 800-777-8570 • FAX 636-225-3739 *www.wmma.com*