

PM 03

499-439

Whitmire
TC 148b

11/5/97

pg 1/4

Microencapsulated Concentrate

KILLS IMMATURE STAGES OF: Aphids, Fungus Gnat Larvae, Mealybugs, Mites, Scale, Thrips and Whiteflies.

FOR USE IN: Greenhouses (Shade and Lath Houses), Garden Centers, Interiorscapes, Florists, Hobby Greenhouses, Homes, Commercial Buildings and Warehouses.

ACTIVE INGREDIENT:

Pyriproxyfen [2-(1-Methyl-2-(4-phenoxyphenoxy) ethoxy) pyridine] 1.3%

INERT INGREDIENTS 98.7%

EPA Reg. No. 499-⁴³⁹UGO EPA Est. No. 499-MO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION


Statement of Practical Treatment

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF SWALLOWED: Call a physician or Poison control Center immediately. Do not induce vomiting.

IF ON SKIN: Remove contaminated clothing and wash before reuse. Wash skin with soap and warm water. Get medical attention if irritation persists.

NET WEIGHT: 1 gal (1pt.) (1qt.)



It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

TC 148b is a flowable microencapsulated concentrate containing pyriproxyfen. TC 148b may be used on any surface which will not be damaged or stained by water. Heavy applications may leave a visible deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

Pyriproxyfen is an insect growth regulator(IGR) that prevents the development of the immature stage of the insect or limits the reproductive capability of the individual if it should reach adulthood. Little or no adult kill should be expected from an application of TC 148b.

This product may be tank mixed with an approved adulticide. For any mixture check for phytotoxicity before contacting plants with the mixture.

Apply pyriproxyfen (Distance) (TC 148b) no more than two times per cropping cycle or no more than two times per six months.

If a high population of adult insects is present, the use of an adulticide may be necessary for immediate relief.

Mixing Instructions

Thoroughly clean spray equipment before using 148b. Shake concentrate well before diluting. When diluting, first add approximately one-half of the water to the spray tank and then add the proper amount of 148b. When emptying the bottles triple rinse with water, shake well and add the rinsate to the spray tank. Add the rest of the water and agitate the sprayer thoroughly. Agitate sprayer occasionally during use to insure even mixture. Shake or reagitate sprayer if spraying is interrupted or if dilution is left in sprayer overnight. If spray screens are used, they should be 50 mesh or larger.

(THIS NEXT SECTION TO BE IN A BOX)

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Waterproof gloves, and Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

(BOX TO END HERE)

Mixing and Dilution Instructions:

Heavy Infestation: 0.3 fl. oz. TC 148b per gallon of water
Preventative and Low Infestation: 0.15 fl. oz. TC 148b per gallon of water

Spray upper and lower foliage surfaces of plants uniformly. Repeat as necessary.
It is permissible to treat under benches for the control of insects.

Drenching: Make soil application to cover soil surface or use adequate volume to moisten soil to desired depth. Apply when soil is drier, but do not stress plant with dryness prior to treatment. Repeat as necessary. For plug flats or trays, use lower rate. Do not apply more than once per week.

Special Notice: This product has demonstrated excellent plant safety; however, all varieties of the plants listed have not been tested. When treating large numbers of plants of a single variety, spray a few plants several days before the main application and observe for phytotoxicity prior to full scale application.

FOR USE INDOORS:

Bedding plants (including): impatiens, marigolds, petunias, geraniums, garden mums, verbena, New Guinea impatiens, and dahlia.

Cut flowers (including): roses, chrysanthemums, carnations, snapdragons, and orchids.

Flowering hanging baskets (including): fuchsia and lantana.

Foliage (such as): Boston fern, dracaena, ficus, schefflera, and philodendron,

Potted flowering plants (including): poinsettia, chrysanthemums, florist's azaleas, lilies, African violets, geraniums, hibiscus, kalanchoes, begonias, gloxinia, cyclamen, cineraria, calceolaria, and exacum.

Trees & shrubs (including): azalea, barberry, cotoneaster, euonymus, holly, ivy, juniper, oak, pine, rhododendron, roses, spiraea, spruce, virburnum, and yew.

Direct spray to some open blooms may cause petal drop.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides or fertilizers. Do not store above 122°F for extended periods of time. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse, wrap container and put in trash.

CAUTION

4/4

Harmful if swallowed or absorbed through skin. Do not breathe vapors or spray mist. Avoid contact with skin or eyes. In case of contact, flush with plenty of water. Wash with Soap and water after use. Obtain medical attention if irritation persists. Avoid contamination of food or feedstuffs.

Do not spray where electrical short circuits might result, such as wall outlets, conduits, etc.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergents and hot water. Keep and wash PPE separately from other laundry.

Personal Protective Equipment: APPLICATORS AND OTHER HANDLERS MUST WEAR: Long-sleeved shirt and long pants, Waterproof gloves, Shoes plus socks.

(THIS NEXT SECTION TO BE IN BOX)

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

(BOX TO END HERE)

LIMITATION OF LIABILITY

Use only as directed. Buyer assumes all risks in using or handling this product in any way. WHITMIRE EXPRESSLY DISCLAIMS ALL IMPLIED WARRANTIES OF FITNESS FOR A PARTICULAR PURPOSE AND MERCHANTABILITY. IN NO CASE SHALL WHITMIRE BE LIABLE FOR CONSEQUENTIAL, SPECIAL, INCIDENTAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. DAMAGES FROM THE USE OF THIS PRODUCT ARE EXPRESSLY LIMITED TO REPLACEMENT OF THE PRODUCT OR REFUND OF THE PURCHASE PRICE.

A Prescription Treatment® brand insecticide from:
Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Ind. Blvd.
St. Louis, Missouri 63122
Copyright, 1997