

499-467

12/19/2003

1/3

TC 181

**AVERT**

WHITMIRE

TC 181

**AVERT**

WHITMIRE

WHITMIRE

**AVERT**

TC 181

KILLS:

ACTIVE INGREDIENT:

OTHER INGREDIENTS:

EPA Reg. No. 499-467 EPA Est. No. 56207-WI-2

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**



WHITMIRE MICRO-GEN

NET WEIGHT: 0.63 oz. (18 grams)

Contains 12 Cloud Resistant 9.1% Spores at 0.05 oz. (1.5 grams per unit)

WHITMIRE

**AVERT**

TC 181

TC 181 - Outer Package

© 2002 Whitmire Micro-Gen Research Laboratories, Inc.

Version 2.0 (Rev. 8/03)

**AVERT**

TC 181

ACCEPTED  
DEC 19 2003

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

499-467

## PRECAUTIONARY STATEMENTS

### FIRST AID

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor immediately for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 1-800-225-3320 for emergency medical treatment information.

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

### DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

Whitmire AVERT TC 181 bait stations contain a slow acting insect bait. For best results placement of the bait should be into or near insect harborage sites.

**FOR USE IN AND AROUND:** Apartments, Campgrounds, Food Storage Areas, Homes, Hospitals, Hotels, Meat Packing and Food Processing Plants, Motels, Nursing Homes, Resorts, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes), Utilities, Warehouses, and other Commercial and Industrial Buildings.

**A PRESCRIPTION TREATMENT® brand insecticide from:**

Whitmire Micro-Gen Research Laboratories, Inc.  
3568 Tree Court Industrial Blvd.  
St. Louis MO 63122-6682  
[www.wmrmg.com](http://www.wmrmg.com)

TC 181 - Outer Package

© 2002 Whitmire Micro-Gen Research Laboratories, Inc.

Version 7.0.1 (rev. 6/03)

### STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**STORAGE:** Store in a cool, dry area inaccessible to children.

**DISPOSAL:** Wrap container and put in trash collection.

**COCKROACHES:** Concentrate treatment at insect activity sites, and other sites where these insects are found.

1. Break bait stations apart from the plastic frame. [Alternate Directions: Remove bait station from container.]
2. For thorough roach control, use as many bait stations as necessary to achieve good coverage and control of the existing infestation for the treated area.
3. For best results, place the bait stations next to the wall along baseboards, in corners, under sinks, in cabinets and near plumbing fixtures. Use under and around equipment and appliances. If possible, place in voids.
4. Bait stations work best on horizontal surfaces touching corners and walls. Remove peel and stick backing on reverse side of bait station before applying to vertical surfaces.
5. Replace bait stations when necessary, approximately every 3 months.

**APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS:** Food handling establishments include places other than private residences in which food is held, processed, prepared or served.

**FOOD AREAS:** Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Avoid contamination of food. Use as instructed above.

**AIRCRAFT CABINS:** Place bait stations in suspected and infested cockroach areas. Place bait stations in areas between elements of construction. Use up to 12 bait stations per galley and up to 6 bait stations per rest room. In cargo holds, use 12 to 24 bait stations per 100 sq. feet. Do not use bait stations in cockpit area.

TC 181

# AVERT®


WHITMIRE

RESEARCH LABORATORIES, INC.  
800-777-8570

Contains 12 Child Resistant Bait Stations at 0.05 oz. (1.5 grams) per unit.

2/3

[BASIC PACKAGE DESIGN - STATION LABEL]



[ALTERNATE ESTABLISHMENT NUMBER]

