

499-473

3/27/2001

1/2



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
499-473

Date of Issuance:
MAR 27 2001

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance: **Unconditional**

Name of Pesticide Product:
TC 205 Injection System
Insecticide

Name and Address of Registrant (include ZIP Code):

Whitmire Micro-Gen Research Laboratories, Inc
3568 Tree Court Industrial Boulevard
St. Louis, MO 63122-6682

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5). A stamped copy of the label is enclosed for your records. If you have any questions regarding this action, please contact Dr. William Sprout of my team at (703) 308-8587.

Signature of Approving Official:

George T. LaRocca

Date:

MAR 27 2001

EPA Form 8570-6

Enclosure

7506
SPROUT

2/2

OUTDOOR TREATMENTS

SUBTERRANEAN ANTS SUCH AS ARGENTINE ANTS, FIRE ANTS, AND HARVESTER ANTS: To treat nests, use the Whitmire Micro-Gen soil injector. To maximize insecticide distribution, insert injector into the nest full length if possible. Inject the nest at 3-6 sites at a rate of 5-10 seconds per site. For very large nests (over 12" diameter), increase the number of injection sites. Space injection sites in a circular pattern on the nest surface with one site in the center. To quiet active ants or nest surface, sweep surface with spray from a distance of 12 inches. For best results, apply when temperature is 65-85°F, or in early morning or late evening hours. Treat new nests as they appear.

PERIMETER TREATMENT: For crawling insects, apply TC 205 to cracks and crevices where insects may enter premises or harbor. Areas to be treated include, but are not limited to, weep holes, under siding, wall voids, soffits around attic vents, cracks and crevices openings around windows, doors and pipes and landscape timbers. Treatment of infested fences and tree holes is permitted.

TERMITES AND CARPENTER ANTS LIVING INSIDE TREES: Determine the highest and lowest elevation of the colony. Drill holes into colony at highest elevation and inject 5 to 10 seconds. Do the same for lowest elevation. Drill intermediate holes every four feet between the upper and lower elevations if required.

AERIAL TERMITE CARTON IN TREES: Drill or punch hole into center of carton, inject for 5 to 10 seconds. If carton is greater than 12 inches across, open another injection site, inject for 5 to 10 seconds. Make one additional injection site for each 12 inches of carton.

TERMITES AND CARPENTER ANTS IN STUMPS, UTILITY POLES AND FENCES: Treat open holes or drill horizontal holes in a circular pattern where insects are suspected. Inject 2 to 5 seconds of product per hole.

TREATMENT FOR IN-GROUND SERVICE BOXES: Ants, Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Fire Ants, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders (such as Black Widow and Brown Recluse), Springtails. Use Whitmire Micro-Gen soil injector or spray actuator to treat open boxes and exposed surfaces. Do not contact any electrical wiring or mechanical parts of meter.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: This container may be recycled in aerosol recycling centers. At present, there are only a few such centers in the country. Before offering for recycling, empty the can by using the product according to the label (DO NOT PUNCTURE!). If the recycling is not available, replace cap and discard container in trash, sanitary landfill or by other approved state and local procedures. Do not incinerate or puncture.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Apply this product only as specified on the label. Spray will damage plant foliage.

PHYSICAL OR CHEMICAL HAZARDS

Extremely Flammable. Contents under pressure. Keep away from fire, sparks and heated surfaces. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces. Do not spray directly into any electronic equipment such as radios, televisions, computers, etc.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



A Prescription Treatment® brand insecticide from Whitmire Micro-Gen Research Laboratories, Inc. 3568 Tree Court Industrial Blvd. St. Louis, MO 63122-6682 www.whitmg.com © 2001 Whitmire Micro-Gen Research Laboratories, Inc.



ACCEPTED

MAR 27 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, or amended thereunder, registered under EPA Reg. No. 499-473

Whitmire Micro-Gen TC 205 Injection System

For Commercial/Institutional Use Only.

KILLS:

Ants, Booklice, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs, Elmleaf Beetles, Fire Ants, Flour Beetles, Grain Weevils, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders, Springtails, Above-Ground Termites, Ticks, Trogloderma (Cabinet, Khapra, and Warehouse Beetles), Wood Infesting Borers and Beetles, and Wood Wasps.

FOR USE IN AND AROUND:

Apartments, Campgrounds, Food Storage Areas, Homes, Hospitals, Hotels, Meat Packing and Food Processing Plants, Motels, Nursing Homes, Resorts, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes), Utilities, Warehouses, and other Commercial and Industrial Buildings. * Use in cargo areas only. Do not use in aircraft cabins.

ACTIVE INGREDIENT:

λ-Cyhalothrin 0.05%

OTHER INGREDIENTS: 99.95%

Contains petroleum distillates. TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically. Contains petroleum distillate - vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

See side panel for additional Precautionary Statements

EPA Reg. No. 499-473 • EPA Est. No. 499-MO-1

NET WEIGHT: 1 lb., 1.5 oz. (17.5 oz.)

800-777-8570



WHITMIRE MICRO-GEN RESEARCH LABORATORIES INC.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in hospital patient rooms while occupied or in any rooms occupied by the elderly or infirm. Do not apply to classrooms when in use. Do not apply to institutions (including libraries, sport facilities, etc.) in the immediate area when occupants are present.

HOW TO USE THE WHITMIRE MICRO-GEN CRACK AND CREVICE® INJECTION SYSTEM: Use Whitmire Micro-Gen TC 205 with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment. Inject TC 205 into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For closed voids calculate the void's cubic area and treat at the rate of one to five seconds per three cubic feet. Several holes may be required in long-running voids.

INDOOR TREATMENTS

CRAWLING INSECTS (Cockroaches, Dermestids, Scorpions, Spiders, Earwigs, Crickets, Sowbugs, Millipedes, Centipedes): Inject the prescribed dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewer, floor drains and meter boxes.

TICKS: Thoroughly treat infested and egg laying sites near pet resting areas including, but not limited to, cracks and crevices, behind baseboards, door and window frames. Do not use on pets.

BOOKLICE & SILVERFISH: Inject into cracks and crevices of baseboards, door frames, bookcases and other harborage sites.

ANTS: Inject cracks, crevices and voids where ants may be traveling and/or harboring, including, but not limited to, around doors, window frames, wall voids and other structural cracks and crevices.

GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, DRUGSTORE BEETLES, TROGODERMA (IN EXPOSED ADULT AND LARVAL STAGES) AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices in cabinets and pantries, wall voids, around window and door frames, under siding and in machinery.

APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS

Food Handling Establishments are places other than private residences in which food is held, processed, prepared or served.

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE® TREATMENT ONLY: Food areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) is also considered a food area.

Apply prescribed amounts of TC 205 directly into cracks and crevices or voids using injection tube or other Whitmire Micro-Gen equipment. Apply to points between different elements of construction, between equipment and floor, openings leading to voids and hollow spaces in walls, equipment legs and bases where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food, food utensils or food processing surfaces. Food must be covered or removed in areas being treated. Food contact surfaces and equipment must be cleaned with an effective cleaning compound and rinsed with potable water before using.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD-HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE® TREATMENT ARE NOT PERMITTED.

APPLICATIONS WITHIN FOOD SERVING AREAS (facilities where foods are served, such as dining rooms): Apply as a Crack and Crevice® treatment to selective surfaces such as baseboards, under elements of construction, and into crack and crevices. Avoid treating surfaces likely to be contacted by food. Food should be covered or removed in area being treated. Do not apply directly to food or allow applications to contaminate food.

INDOOR AND OUTDOOR TREATMENTS

WOOD DESTROYING INSECTS - Localized treatment of wood-infesting insects including Beetles and Borers (such as the following families: Anobiidae, Bostrichidae, Buprestidae, Dermomyiidae, Lyctidae, Oedemeridae, Sesiidae), Carpenter Ants, Carpenter Bees, Termites and Wood Wasps: Locate suspected infestation and/or insect tunnels and cavities. Drill pilot holes into insect tunnels, galleries and inaccessible closed voids. Drill holes 8 to 10 inches apart to insure thorough coverage. Inject 5 to 10 seconds per hole. Do not contact electrical wiring, plumbing, etc. when drilling and never use Whitmire Micro-Gen metal wood injectors where electrical shock could occur. Use a Whitmire Micro-Gen standard plastic wood injector if there is danger of electrical contact. For carpenter bees, seal bee entrances after application. This treatment is intended for localized termite infestations only. For severe infestations or treatments under slabs, in basements and buildings with insufficient crawl space, contact a licensed pest control operator or termite service. Not recommended as sole protection against termites. The purpose of treatments with this product is to kill workers or winged forms which may be present in the treated galleries at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement.

FOR ACTIVE AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames.