Please read instructions on reverse before completing form.	Form Approved, OMB No. 2070-0060, Approvel expires 2-28-9
United States Environmental Protection Weshington, DC 20460	Registration OPP Identifier Number Agency Amendment
Application	for Pesticide - Section I
1. Company/Product Number 499-473	2. EPA Product Manager 3. Proposed Classification George LaRocca None Restricted
4. Company/Product (Name) Whitmire Micro-Gen TC 205 Injection System	PM# 13
5. Name and Address of Applicant (Include ZIP Code) Whitmire Micro-Gen Research Laboratories 3568 Tree Court Industrial Blvd St Louis MO 63122-6682 Check if this is a new address	Inc (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name
	Section - II
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.	Final printed labels in repsonse to Agency letter dated "Me Too" Application. Other - Explain below.
EPA's letter to NSF dated July 28, 2004 and the requirements of EPA's made to the labeling or the confidential statement of formula of this prostatement to EPA. I further understand that if this notification is not confrements of 40 CFR 156.10 and 40 CFR 152.46, this product may ons 12 and 14 of FIFRA.	tter for complete details. This notification is consistent with the guidance set forth in s regulations at 40 CFR 156.10 and 40 CFR 152.46, and no other changes have been oduct. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false sistent with the guidance set forth in EPA's letter to NSF dated July 28, 2004 and the be in violation of FIFRA and I may be subject to enforcement action and penalties under
Material This Product Will Be Packaged In:	
Yes Yes No No Per	Water Soluble Packaging Yes No If "Yes" Package wgt No, per Container Additional Plastic Glass Paper Other (Specify)
3. Location of Net Contents Information 4. Size(s) Retail	Container 5. Location of Label Directions
6. Manner in Which Lebel is Affixed to Product Lithograph Paper glue Stenciled	h Other
	Section - IV
1. Contact Point (Complete items directly below for identification o	f individual to be contacted, if necessary, to process this application.)
Name Titl Dana Thomas Ma	Telephone No. (include Area Code) anager, Product Registrations 636-861-4223
Certification I certify that the statements I have made on this form and all I acknowledge that any knowlingly false or misleading statem both under applicable law.	attachments thereto are true, accurate and complete.
Alara M Homas Ma	Fitte
4. Typed Name 5. C	Jate
Dana M. Thomas	November 29, 2004

WHITMIRE MICRO-GEN TC 205 INJECTION SYSTEM

[ALTERNATE BRAND NAME: Prescription Treatment brand 221L Residual Insecticide]

KILLS:

Ants, Booklice, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs, Elmleaf Beetles, Fire Ants, Flour Beetles, Grain Weevils, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders, Springtails, Above-Ground Termites, Ticks, Trogoderma (Cabinet, Khapra, and Warehouse Beetles), Wood Infesting Borers and Beetles, and Wood Wasps.

FOR USE IN AND AROUND:

Apartments, Campgrounds, Food Storage Areas, Homes, Hospitals, Hotels, Meat Packing and Food Processing Plants, Motels, Nursing Homes, Resorts, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes†), Utilities, Warehouses, and other Commercial and Industrial Buildings.
† Use in cargo areas only. Do not use in aircraft cabins.

ACTIVE INGREDIENT:

λ-Cyhalothrin	0.05%
OTHER INGREDIENTS:	
Contains petroleum distillates.	TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional Precautionary Statements.

EPA Reg. No. 499-473
 EPA Est. No. 499-MO-1

NET WEIGHT:



PRECAUTIONARY STATEMENTS

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-225-3320 for emergency medical treatment information.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically. Contains petroleum distillate - vomiting may cause aspiration pneumonia.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Apply this product only as specified on the label. Spray will damage plant foliage.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces. Do not spray directly into any electronic equipment such as radios, televisions, computers, etc.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

SHAKE WELL BEFORE USING

See supplemental label for complete Directions for Use.

Do not apply this product in hospital patient rooms while occupied or in any rooms occupied by the elderly or infirm. Do not apply to classrooms when in use. Do not apply to institutions (including libraries, sport facilities, etc.) in the immediate area when occupants are present.

HOW TO USE WHITMIRE MICRO-GEN CRACK AND CREVICE® INJECTION SYSTEM

CRACK & CREVICE TREATMENT: Use Whitmire Micro-Gen TC 205 with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment. Inject TC 205 into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For closed voids calculate the void's cubic area and treat at the rate of one to five seconds per three cubic feet. Several holes may be required in long-running voids.

SPOT TREATMENT: Use TC 205 with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment. Hold container upright while spraying. Direct nozzle approximately 12 inches from surface to be treated. Apply directly on insects in these locations when possible.

INDOOR TREATMENTS

CRAWLING INSECTS (Cockroaches, Dermestids, Scorpions, Spiders, Earwigs, Crickets, Sowbugs, Millipedes, Centipedes): Apply as a Crack & Crevice treatment using the prescribed dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewer, floor drains and meter boxes. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

TICKS: Apply as a Crack & Crevice treatment or Spot treatment into infested and egg laying sites near pet resting areas including, but not limited to cracks and crevices, behind baseboards, door and window frames. Do not use on pets.

BOOKLICE & SILVERFISH: Apply as a Crack & Crevice treatment into cracks and crevices of baseboards, door frames, bookcases and other harborage sites. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

ANTS: Apply as a Crack & Crevice treatment into cracks, crevices and voids where ants may be traveling and/or harboring, including, but not limited to, around doors, window frames, wall voids and other structural cracks and crevices. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, DRUGSTORE BEETLES, TROGODERMAS (IN EXPOSED ADULT AND LARVAL STAGES) AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices in cabinets and pantries, wall voids, around window and door frames, under siding and in machinery. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry grea away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not puncture or incinerate! Empty container by using the product in accordance with the label directions. Offer empty container for recycling, if available, or place in trash if allowed by state and local regulations. If container is partly full, contact your local solid waste agency or call 1-800-CLEANUP for disposal instructions.



Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
www.wmmg.com
© 2004 Whitmire Micro-Gen Research Laboratories, Inc.

IADDITIONAL LABELING - ENCLOSED IN CARTON WITH PRODUCTI

INDOOR TREATMENTS APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS



Food Handling Establishments are places other than private residences in which food is held, processed, prepared or served.

In Federally inspected meat and poultry plants, the facility must not be in operation when this product is used, regardless of the method of application. In non-Federally inspected facilities, Spot and/or Crack and Crevice application may be used while the facility is in operation provided exposed food is covered or removed from the area being treated prior to application.

Do not apply directly to food. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food, food utensils or food processing surfaces. Food must be covered or removed in areas being treated. Spray should not be applied directly to surfaces or utensils that may come into contact with food. Food contact surfaces and equipment must be cleaned with an effective cleaning compound and rinsed with potable water before usina.

Reapplication may be made at 21 day intervals. Limit individual spot treatments to an area no larger than 20% of the treated surface. Individual spot treatments should not exceed two square feet.

FOOD AREAS: Food areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) is also considered a food area.

Apply as a Crack and Crevice treatment or Spot treatment directly into cracks, crevices, voids or onto surfaces where insects are resting. Use an injection tube or other Whitmire Micro-Gen equipment capable of delivering a pinstream. Apply to different elements of construction; between equipment and floor; openings leading to hollow spaces in walls, equipment legs and bases; and where insects hide or rest.

FOOD SERVING AREAS (facilities where foods are served, such as dining rooms): Apply as a Crack and Crevice treatment or Spot treatment to selective surfaces such as baseboards, under elements of construction, and into cracks and crevices. Avoid treating surfaces likely to be contacted by food. Food should be covered or removed in area being treated. Do not apply directly to food or allow applications to contaminate food.

NON-FOOD AREAS: Includes areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling).

Apply as a Crack & Crevice treatment or Spot treatment to selective surfaces such as baseboards, under elements of construction, into cracks and crevices. Avoid treating surfaces likely to be contacted by food.

INDOOR AND OUTDOOR TREATMENTS

WOOD DESTROYING INSECTS - Localized treatment of wood-infesting insects including Beetles and Borers (such as the following familles: Anobiidae, Bostrichidae, Buprestidae, Cerambycidae, Lyctidae, Oedemeridae, Sestidae), Carpenter Ants, Carpenter Bees, Termites and Wood Wasps: Locate suspected infestation and/or insect tunnels and cavities. Drill pilot holes into insect tunnels, galleries and inaccessible closed voids. Drill holes 8 to 10 inches apart to ensure thorough coverage. Inject 5 to 10 seconds per hole. Do not contact electrical wiring, plumbing, etc. when drilling and never use Whitmire Micro-Gen metal wood injectors where electrical shock could occur. Use a Whitmire Micro-Gen standard plastic void injector if there is danger of electrical contact. For carpenter bees, seal bee entrances after application. This treatment is intended for localized termite infestations only. For severe infestations or treatments under slabs, in basements and buildings with insufficient crawl space, contact a licensed pest control operator or termite service. Not recommended as sole protection against termites. The purpose of treatments with this product is to kill workers or winged forms which may be present in the treated galleries at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement.

FOR ACTIVE AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames.

OUTDOOR TREATMENTS

SUBTERRANEAN ANTS SUCH AS ARGENTINE ANTS, FIRE ANTS, AND HARVESTER ANTS: To treat nests, inject the nest at 3-6 sites at a rate of 5-10 seconds per site. For very large nests (over 12" diameter), increase the number of injection sites. Space injection sites in a circular pattern on the nest surface with one site in the center. To quiet active ants on nest surface, sweep surface with spray from a distance of 12 inches. For best results, apply when temperature is 65-85°F, or in early morning or late evening hours. Treat new nests as they appear.

PERIMETER TREATMENT: For crawling insects, apply as a Crack & Crevice treatment or Spot treatment into cracks and crevices where insects may enter premises or harbor. Areas to be treated include, but are not limited to, weep holes, under siding, wall voids, soffits, around attic vents, cracks and crevices, openings around windows, doors and pipes and landscape timbers. Treatment of infested fences and tree holes is permitted.

TERMITES AND CARPENTER ANTS LIVING INSIDE TREES: Determine the highest and lowest elevation of the colony. Drill holes into colony at highest elevation and inject 5 to 10 seconds. Do the same for lowest elevation. Drill intermediate holes every four feet between the upper and lower elevations if required.

AERIAL TERMITE CARTON IN TREES: Drill or punch hole into center of carton. Inject for 5 to 10 seconds. If carton is greater than 12 inches across, open another injection site. Inject for 5 to 10 seconds. Make one additional injection site for each 12 inches of carton.

TERMITES AND CARPENTER ANTS IN STUMPS, UTILITY POLES AND FENCES: Treat open holes or drill horizontal holes in a circular pattern where insects are suspected. Inject 2 to 5 seconds of product per hole.

TREATMENT FOR IN-GROUND SERVICE BOXES: Ants, Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Fire Ants, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders (such as Black Widow and Brown Recluse), and Springtails. Use spray actuator to treat open boxes and exposed surfaces. Do not contact any electrical wiring or mechanical parts of meter.

v4.2 - 12/04

8/21

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
www.wmmg.com
© 2003 Whitmire Micro-Gen Research Laboratories, Inc.



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

JUL ES ME

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Mr. Kenji Yano NSF International Nonfood Compounds Registration Program 789 N. Dixboro Road Ann Arbor, MI 48105

Re: Use of NSF logo

Dear Mr. Yano:

This is in response to your letter of June 17, 2004 which provides a revised NSF logo to satisfy concerns raised in a May 13, 2004 conference call between representatives of NSF and the Agency.

The following is a summary of agreements reached during the conference call:

- Use of the NSF mark with a check mark is permitted
- A NSF 6-digit listing number is permitted
- Use of the phrase "NSF listed" is permitted whereas use of "NSF registered" is unacceptable
- Use of the NSF logo on the side/back panels is permitted
- Rame Cromwell is the EPA contact for other uses of the logo

The Agency continues to have concern about the placement of the NSF mark on the front panel of a pesticide label. Logos and associated language should not be placed in a position on the label which might negate or detract from required labeling statements. The Agency would prefer the logo to be placed on the back or side panel of the label to avoid any confusion.

The Agency approves logos on a case-by-case basis. At a minimum the logo must be submitted through the Agency's notification process as described in PR Notice 98-10. Before coming in with a notification, the company desiring to use this logo should contact their product manager for

further guidance. A copy of the proposed modification of the label bearing the actual logo and any associated language must be submitted with the notification.

If you have additional questions please do not hesitate to contact me.

Sincerely,

Frank T. Sanders, Director Antimicrobials Division

11/2





January 28, 2004

Dana Thomas
WHITMIRE MICRO-GEN RESEARCH LABORATORIES, INC.
3568 TREE COURT INDUSTRIAL BLVD.
ST. LOUIS, MO 63122-6682
UNITED STATES

RE PRESCRIPTION TREATMENT® brand 221L

Category Code: F2

NSF Registration No. 132391

Dear Dana Thomas:

NSF has processed the application for Registration of PRESCRIPTION TREATMENT® brand 221L to the NSF Registration Guidelines for Proprietary Substances and Nonfood Compounds (2004), which are available at www.nsf.org/usda. The NSF Nonfood Compounds Registration Program is a continuation of the USDA product approval and listing program, which is based on meeting regulatory requirements including FDA 21 CFR for appropriate use, ingredient and labeling.

This product is acceptable for use as a Residual Pesticide (F2) provided its use is generally restricted to inedible product areas or outside premises where exposed food products are neither processed nor stored in open containers. The product may be used in edible product areas for crack and crevice treatment provided applicable directions appear on the EPA registered label. Such use is acceptable provided the pesticide is applied after processing operations have ended, exposed food products and packaging materials are removed or protected, the treated area is ventilated and cleaned thoroughly before processing operations are resumed, and the treated crack and crevices are sealed.

NSF Registration of this product is current when the NSF Registration Number, Category Code, and Registration Mark appear on the NSF-approved product label, and the registered product name is included in the current NSF White Book Listing of Nonfood Compounds at the NSF website (http://www.nsf.org/usda). The NSF Registration Mark can be downloaded from the NSF website, at http://www.nsf.org/mark/download_marks.html.

NSF Listing of all registered Nonfood compounds by NSF International is not an endorsement of those compounds, or of any performance or efficacy claims made by the manufacturer.


Registration status may be verified at any time via the NSF web site, at http://www.nsf.org/usda. Changes in formulation or label, without the prior written consent of NSF, will void registration, and will supersede the on-line listing.

Sincerely.

Carmen Grindatti

NSF Nonfood Compounds Registration Program

Company No: N10443



To Joyce Edwards/DC/USEPA/US@EPA cc bcc

Subject 499-473; Notification to add NSF Logo 11-29-04

Hi Joyce,

Attached is the revised labeling for the Notification submitted 11/29/04 to add the NSF logo and listing number. The offending associated text has been removed. Do you need me to send a letter or revised application? I can do that by e-mail as well. It will only take a few minutes to do, so just please let me know.

Also, is this the only one you have? I'm asking because I sent in similar notifications for two other products on the same day (but different PMs handle those products – EPA Reg #s 499-290 and 499-406). If I need to revise those labels, I can, but I don't want to jump the gun.

Thanks for your help – look forward to hearing from you soon.

Dana

Dana Thomas
Manager, Product Registrations
Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis, MO 63122-6682
Phone: (636) 861-4223
Fax: (636) 225-3739

e-mail: dana.thomas@wmmg.com

•••

000499-00473.20050127.tc205.v42_FPL.pdf

November 29, 2004

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504C) U.S. Environmental Protection Agency Room 266A, Crystal Mall 2 1801 Bell Street Arlington, VA 22202

RE:

NOTIFICATION TO ADD NSF LISTING

PER AGENCY LETTER TO NSF DATED 7/28/04 (copy enclosed)

Whitmire Micro-Gen TC 205 Injection System EPA Reg. No. 499-473

NOTIFICATION

JAN 2 7 2005

Dear Sir/Madam:

Enclosed in support of this action is a completed application and one copy revised labeling for the referenced product. The NSF logo, listing number and associated text has been added to the product label in proximity to the directions for use in food/feed areas of food/feed handling establishments.

This notification is consistent with the guidance set forth in EPA's letter to NSF dated July 28, 2004 and the requirements of EPA's regulations at 40 CFR 156.10 and 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance set forth in EPA letter to NSF dated July 28, 2004 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Also enclosed is a self-addressed, "No Postage Necessary" postcard that, when filled out, would indicate the status of this notification. Please take a moment to fill in the appropriate information and drop it in the mail for our records. Your assistance in this regard is certainly appreciated.

Thank you for your assistance in this matter. Please don't hesitate to contact me at (636) 861-4223 or, via electronic mail at <u>dana.thomas@wmmg.com</u>, should you have any questions or require additional information concerning this action.

Sincerely,

WHITMIRE MICRO-GEN RESEARCH LABORATORIES INC.

Dana M. Thomas

Manager, Product Registrations

Hara M. Thomas

Encl.

v4.2 - 12/04 15/21

WHITMIRE MICRO-GEN TC 205 INJECTION SYSTEM

[ALTERNATE BRAND NAME: Prescription Treatment brand 221L Residual Insecticide]

KILLS:

Ants, Booklice, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chocólate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs, Elmleaf Beetles, Fire Ants, Flour Beetles, Grain Weevils, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders, Springtails, Above-Ground Termites, Ticks, Trogoderma (Cabinet, Khapra, and Warehouse Beetles), Wood Infesting Borers and Beetles, and Wood Wasps.

FOR USE IN AND AROUND:

Apartments, Campgrounds, Food Storage Areas, Homes, Hospitals, Hotels, Meat Packing and Food Processing Plants, Motels, Nursing Homes, Resorts, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes†), Utilities, Warehouses, and other Commercial and Industrial Buildings.

† Use in cargo areas only. Do not use in aircraft cabins.

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional Precautionary Statements.

EPA Reg. No. 499-473 • EPA Est. No. 499-MO-1

NET WEIGHT:



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.
800-777-8570

PRECAUTIONARY STATEMENTS

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-225-3320 for emergency medical treatment information. NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically. Contains petroleum distillate - vomiting may cause aspiration pneumonia.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal áreas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Apply this product only as specified on the label. Spray will damage plant foliage.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces. Do not spray directly into any electronic equipment such as radios, televisions, computers, etc.

4.2 - 12/04

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

SHAKE WELL BEFORE USING

See supplemental label for complete Directions for Use.

Do not apply this product in hospital patient rooms while occupied or in any rooms occupied by the elderly or infirm. Do not apply to classrooms when in use. Do not apply to institutions (including libraries, sport facilities, etc.) in the immediate area when occupants are present.

HOW TO USE WHITMIRE MICRO-GEN CRACK AND CREVICE® INJECTION SYSTEM

CRACK & CREVICE TREATMENT: Use Whitmire Micro-Gen TC 205 with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment. Inject TC 205 into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For closed voids calculate the void's cubic area and treat at the rate of one to five seconds per three cubic feet. Several holes may be required in long-running voids.

SPOT TREATMENT: Use TC 205 with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment. Hold container upright while spraying. Direct nozzle approximately 12 inches from surface to be treated. Apply directly on insects in these locations when possible.

INDOOR TREATMENTS

CRAWLING INSECTS (Cockroaches, Dermestids, Scorpions, Spiders, Earwigs, Crickets, Sowbugs, Millipedes, Centipedes): Apply as a Crack & Crevice treatment using the prescribed dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewer, floor drains and meter boxes. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

TICKS: Apply as a Crack & Crevice treatment or Spot treatment into infested and egg laying sites near pet resting areas including, but not limited to cracks and crevices, behind baseboards, door and window frames. Do not use on pets.

BOOKLICE & SILVERFISH: Apply as a Crack & Crevice treatment into cracks and crevices of baseboards, door frames, bookcases and other harborage sites. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

ANTS: Apply as a Crack & Crevice treatment into cracks, crevices and voids where ants may be traveling and/or harboring, including, but not limited to, around doors, window frames, wall voids and other structural cracks and crevices. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

18/21

GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, DRUGSTORE BEETLES, TROGODERMAS (IN EXPOSED ADULT AND LARVAL STAGES) AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices in cabinets and pantries, wall voids, around window and door frames, under siding and in machinery. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not puncture or incinerate! Empty container by using the product in accordance with the label directions. Offer empty container for recycling, if available, or place in trash if allowed by state and local regulations. If container is partly full, contact your local solid waste agency or call 1-800-CLEANUP for disposal instructions.



Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc. 3568 Tree Court Industrial Blvd.

St. Louis MO 63122-6682

www.wmmg.com

© 2004 Whitmire Micro-Gen Research Laboratories, Inc.

[ADDITIONAL LABELING - ENCLOSED IN CARTON WITH PRODUCT]

INDOOR TREATMENTS APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS



TC 205 [Prescription Treatment brand 221L] is NSF listed and is acceptable for use as a Residual Pesticide (F2).

Food Handling Establishments are places other than private residences in which food is held, processed, prepared or served.

In Federally inspected meat and poultry plants, the facility must not be in operation when this product is used, regardless of the method of application. In non-Federally inspected facilities, Spot and/or Crack and Crevice application may be used while the facility is in operation provided exposed food is covered or removed from the area being treated prior to application.

Do not apply directly to food. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food, food utensils or food processing surfaces. Food must be covered or removed in areas being treated. Spray should not be applied directly to surfaces or utensils that may come into contact with food. Food contact surfaces and equipment must be cleaned with an effective cleaning compound and rinsed with potable water before using.

Reapplication may be made at 21 day intervals. Limít individual spot treatments to an area no larger than 20% of the treated surface. Individual spot treatments should not exceed two square feet.

FOOD AREAS: Food areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) is also considered a food area.

Apply as a Crack and Crevice treatment or Spot treatment directly into cracks, crevices, voids or onto surfaces where insects are resting. Use an injection tube or other Whitmire Micro-Gen equipment capable of delivering a pinstream. Apply to different elements of construction; between equipment and floor; openings leading to hollow spaces in walls, equipment legs and bases; and where insects hide or rest.

FOOD SERVING AREAS (facilities where foods are served, such as dining rooms): Apply as a Crack and Crevice treatment or Spot treatment to selective surfaces such as baseboards, under elements of construction, and into cracks and crevices. Avoid treating surfaces likely to be contacted by food. Food should be covered or removed in area being treated. Do not apply directly to food or allow applications to contaminate food.

NON-FOOD AREAS: Includes areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling).

Apply as a Crack & Crevice treatment or Spot treatment to selective surfaces such as baseboards, under elements of construction, into cracks and crevices. Avoid treating surfaces likely to be contacted by food.

19/21

v4.2 – 12/04

INDOOR AND OUTDOOR TREATMENTS

WOOD DESTROYING INSECTS - Localized treatment of wood-infesting insects including Beetles and Borers (such as the following families: Anobiidae, Bostrichidae, Buprestidae, Cerambycidae, Lyctidae, Oedemeridae, Seslidae), Carpenter Ants, Carpenter Bees, Termites and Wood Wasps: Locate suspected infestation and/or insect tunnels and cavities. Drill pilot holes into insect tunnels, galleries and inaccessible closed voids. Drill holes 8 to 10 inches apart to ensure thorough coverage. Inject 5 to 10 seconds per hole. Do not contact electrical wiring, plumbing, etc. when drilling and never use Whitmire Micro-Gen metal wood injectors where electrical shock could occur. Use a Whitmire Micro-Gen standard plastic void injector if there is danger of electrical contact. For carpenter bees, seal bee entrances after application. This treatment is intended for localized termite infestations only. For severe infestations or treatments under slabs, in basements and buildings with insufficient crawl space, contact a licensed pest control operator or termite service. Not recommended as sole protection against termites. The purpose of treatments with this product is to kill workers or winged forms which may be present in the treated galleries at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement.

FOR ACTIVE AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames.

OUTDOOR TREATMENTS

SUBTERRANEAN ANTS SUCH AS ARGENTINE ANTS, FIRE ANTS, AND HARVESTER ANTS: To treat nests, inject the nest at 3-6 sites at a rate of 5-10 seconds per site. For very large nests (over 12" diameter), increase the number of injection sites. Space injection sites in a circular pattern on the nest surface with one site in the center. To quiet active ants on nest surface, sweep surface with spray from a distance of 12 inches. For best results, apply when temperature is 65-85°F, or in early morning or late evening hours. Treat new nests as they appear.

PERIMETER TREATMENT: For crawling insects, apply as a Crack & Crevice treatment or Spot treatment into cracks and crevices where insects may enter premises or harbor. Areas to be treated include, but are not limited to, weep holes, under siding, wall voids, soffits, around attic vents, cracks and crevices, openings around windows, doors and pipes and landscape timbers. Treatment of infested fences and tree holes is permitted.

TERMITES AND CARPENTER ANTS LIVING INSIDE TREES: Determine the highest and lowest elevation of the colony. Drill holes into colony at highest elevation and inject 5 to 10 seconds. Do the same for lowest elevation. Drill intermediate holes every four feet between the upper and lower elevations if required.

AERIAL TERMITE CARTON IN TREES: Drill or punch hole into center of carton. Inject for 5 to 10 seconds. If carton is greater than 12 inches across, open another injection site. Inject for 5 to 10 seconds. Make one additional injection site for each 12 inches of carton.

TERMITES AND CARPENTER ANTS IN STUMPS, UTILITY POLES AND FENCES: Treat open holes or drill horizontal holes in a circular pattern where insects are suspected. Inject 2 to 5 seconds of product per hole.

TREATMENT FOR IN-GROUND SERVICE BOXES: Ants, Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Fire Ants, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders (such as Black Widow and Brown Recluse), and Springtails. Use spray actuator to treat open boxes and exposed surfaces. Do not contact any electrical wiring or mechanical parts of meter.

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
www.wmmg.com
© 2003 Whitmire Micro-Gen Research Laboratories, Inc.

21/21