

499-475

0011512006

1/14

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 499-475	2. EPA Product Manager Richard Gebken	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Pro-Control Pyrethrin 6EC	PM# 10	
5. Name and Address of Applicant (Include ZIP Code) Whitmire Micro-Gen Research Laboratories Inc. 3568 Tree Court Industrial Blvd. St. Louis, MO 63122-6682 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
FEB 15 2006

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of minor reformatting of label text to accomodate container sizes. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the CSF of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.


Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No			
* Certification must be submitted If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt No. per container		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Patricia A Hodge	Title Registration Specialist	Telephone No. (Include Area Code) (636)861-4272
---------------------------------	---	---

Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Specialist	
4. Typed Name Patricia A Hodge	5. Date 2/1/2006	

2/14

PRO-CONTROL® PYRETHRIN 6EC

NOTIFICATION
FEB 15 2006

[ALTERNATE BRAND NAME #1: Prescription Treatment® brand Pyreth-It® Formula 2]
[Text in this color applies only to this marketplace label]

[ALTERNATE BRAND NAME #2: Prescription Treatment® brand ULD® BP Booster 6-60]
[Text in this color applies only to this marketplace label; numbers in brackets denote order of directions for use]

[ALTERNATE BRAND NAME #3: Prescription Treatment® brand Pyreth-It® 6-60]

[Black text applies to all marketplace labels.]

- For use on growing crops, ornamentals, stored products, and fruit fly control on harvested fruits and vegetables.
- May be used alone as a clean-up spray or a pre-harvest spray.
- To Provide Quick Knockdown of Insects
- Quicker and More Complete Control
- Use Where Resistance May Be a Problem
- May be Combined with other Insecticides
- Food Processing Operations may continue when this product is applied as a Surface Spray

KILLS: Ants, Cadelles, Cigarette Beetles, Confused Flour Beetles, Cockroaches, Crickets, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Flies, Fruit Flies, Grain Moths, Gnats, Hornets, Indian Meal Moths, Lesser Grain Borers, Mediterranean Flour Moths, Mosquitoes, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles, Spider Beetles, Spiders, Square Necked Grain Beetles, Ticks, Wasps, Yellow mealworms

FOR USE IN AND AROUND: Bakeries, Cargo Ships, Food Processing Plants, Grain Storage buildings, Homes, Industrial Installations, Restaurants, Tobacco Warehouses, Trucks, Warehouses.

See full label for a complete list of insects and sites.

ACTIVE INGREDIENT:

Pyrethrins, a botanical insecticide.....	6.00%
Piperonyl butoxide, technical*	60.00%

OTHER INGREDIENTS:	34.00%
Contains petroleum distillate.	TOTAL: 100.00%

* Equivalent to 48% (butylcarbityl)(6-propylpiperonyl) ether and 12% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for ~~Precautionary Statements.~~

~~[Alternate text: See inside booklet for Precautionary Statements and Directions for Use.]~~

EPA Reg. No. 499-475 • EPA Est. No. 499-MO-1

**ULTRA LOW DOSAGE THROUGH MICRON GENERATION
NET CONTENTS:**



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.
800-777-8570 www.wmmg.com



See full label for complete Precautionary Statements and Directions for Use.

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 1-800-225-3320 for emergency medical treatment information.

NOTE TO PHYSICIAN: Contains petroleum distillate - vomiting may cause aspiration pneumonia.


PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. Avoid breathing spray mist. Causes eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove pets and birds, and cover and turn off fish aquariums before spraying. When using in an enclosed area, do not remain in treated area. Ventilate area after treatment is completed. When using this product as a space spray in food processing plants, food should be removed or covered during treatment. Do not apply as a space spray while food processing is underway. Food processing surfaces and equipment must be covered during treatment or cleaned with a suitable detergent and rinsed with potable water before reuse. Except in Federally inspected meat and poultry plants, food processing operations may continue when this product is applied as a surface or pin stream Crack and Crevice spray with care and in accordance with the directions and precautions given on the full label.

A Prescription Treatment® brand insecticide from:
Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
www.wmmg.com

© 2006 Whitmire Micro-Gen Research Laboratories, Inc.



4/14

PRO-CONTROL® PYRETHRIN 6EC

[ALTERNATE BRAND NAME #1: Prescription Treatment® brand Pyreth-It® Formula 2]
[Text in this color applies only to this marketplace label]

[ALTERNATE BRAND NAME #2: Prescription Treatment® brand ULD® BP Booster 6-60]
[Text in this color applies only to this marketplace label; numbers in brackets denote order of directions for use]

[ALTERNATE BRAND NAME #3: Prescription Treatment® brand Pyreth-It® 6-60]

[Black text applies to all marketplace labels.]

- For use on growing crops, ornamentals, stored products, and fruit fly control on harvested fruits and vegetables.
- May be used alone as a clean-up spray or a pre-harvest spray.
 - To Provide Quick Knockdown of Insects
 - Quicker and More Complete Control
 - Use Where Resistance May Be a Problem
 - May be Combined with other Insecticides
 - Food Processing Operations may continue when this product is applied as a Surface Spray

KILLS: Ants, Caddisflies, Cigarette Beetles, Confused Flour Beetles, Cockroaches, Crickets, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Flies, Fruit Flies, Grain Moths, Gnats, Hornets, Indian Meal Moths, Lesser Grain Borers, Mediterranean Flour Moths, Mosquitoes, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles, Spider Beetles, Spiders, Square Necked Grain Beetles, Ticks, Wasps, Yellow mealworms

FOR USE IN AND AROUND: Bakeries, Cargo Ships, Food Processing Plants, Grain Storage buildings, Homes, Industrial Installations, Restaurants, Tobacco Warehouses, Trucks, Warehouses.

ACTIVE INGREDIENT:

Pyrethrins, a botanical insecticide.....	6.00%
Piperonyl butoxide, technical*	60.00%

OTHER INGREDIENTS:	34.00%
---------------------------------	---------------

Contains petroleum distillate.	TOTAL: 100.00%
--------------------------------	----------------

* Equivalent to 48% (butylcarbityl)(6-propylpiperonyl) ether and 12% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 499-475 • EPA Est. No. 499-MO-1

ULTRA LOW DOSAGE THROUGH MICRON GENERATION NET CONTENTS:



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.
800-777-8570 www.wmmg.com

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 1-800-225-3320 for emergency medical treatment information.

NOTE TO PHYSICIAN: Contains petroleum distillate - vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed or inhaled. Avoid breathing spray mist. Causes eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove pets and birds, and cover and turn off fish aquariums before spraying. When using in an enclosed area, do not remain in treated area. Ventilate area after treatment is completed. When using this product as a space spray in food processing plants, food should be removed or covered during treatment. Do not apply as a space spray while food processing is underway. Food processing surfaces and equipment must be covered during treatment or cleaned with a suitable detergent and rinsed with potable water before reuse. Except in Federally inspected meat and poultry plants, food processing operations may continue when this product is applied as a surface or pin stream Crack and Crevice spray with care and in accordance with the directions and precautions given on this label.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers who may be exposed to the dilute and/or concentrate through application or other tasks must wear: Long-sleeved shirt and long pants; Chemical-resistant gloves, such as, Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton; Shoes plus socks; Protective Eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240 (d) (4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS**Users should:**

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift from treated areas may be hazardous to organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

6/14

DIRECTIONS FOR USE**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT WITH ITS LABELING.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

PRO-CONTROL PYRETHRIN 6EC may be used on most crops because its active ingredients are exempt from tolerances when applied to growing crops. The crop grouping scheme used on this label was devised by the Environmental Protection Agency to expedite minor use pesticide registration. Each crop grouping on this label contains the phrase "Including, but not limited to." and then lists a number of crops in each group. This wording allows the use of PRO-CONTROL PYRETHRIN 6EC on crops that may not be specifically listed on this label (providing that the group to which the crop belongs is listed).

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls; Chemical-resistant gloves, such as Barrier Laminate, Nitrile Rubber, Neoprene Rubber, or Viton; Shoes plus socks; Protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR, Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

[1] CONTROLS INSECTS: Such as Ants, Aphids, Apple Maggots, Armyworms, Artichoke Plume Moths, Asparagus Beetles, Beet Armyworms, Bagworms, Bean Beetles, Blister Beetles, Blow Flies, Biting Flies, Boll Weevils, Cabbage Loopers, Cankerworms, Carrot Weevils, Caterpillars, Clover Mites, Clover Weevils, Cockroaches, 12-spotted Cucumber Beetles, Codling Moths, Colorado Potato Beetles, Corn Earworms, Crickets, Crane Flies, Crickets, Cross-striped Cabbageworms, Cucumber Beetles, Deer Flies, Deer Ticks, Earwigs, Diamondback Larvae, Eastern Tent Caterpillars, Elm Leaf Beetles, European Corn Borers, European Pine Tip Moths, Face Flies, Fall Webworms, Fire Ants, Firebrats, Fireworms, Flea Beetles, Flies, Forest Tent Caterpillars, Fungus Gnats, Fruit Flies, Fruittree Leafrollers, Grape Leafhoppers, Grape Leaf Skeletonizers, Grasshoppers, Green Fruit Worms, Green Peach Aphids, Greenhouse Thrips, Gypsy Moths (adults & larvae), Harlequin Bugs, Heliothis sp, Hornets, Horn Flies, Hornworms, Horse Flies, House Flies, Imported Cabbageworms, Indian Meal Moths, Japanese Beetles, Katydid, Lace Bugs, Leafhoppers, Leafrollers, Leaf-tiers, Lice, Loopers, Lygus, Mealy Bugs, Mediterranean Flour Moths, Mexican Bean Beetles, Midges, Millipedes, Mosquitoes, Mushroom Flies, Navel Orangeworms, Onion Maggots, Pear Psylla, Potato Leafhoppers, Psyllids, Rice Weevils, Saw Toothed Grain Beetles, Silverfish, Skippers, Sowbugs, Spiders, Stable Flies, Stink Bugs, Tabanidae, Tarnished Plant Bugs, Thrips, Tomato Hornworms, Vinegar Flies, Wasps, Webworms, Whiteflies and Yellow Jackets.

[2] GENERAL USE DIRECTIONS

USED ALONE: This concentrate can also be used as a clean-up or a pre-harvest spray where other materials cannot be used because of residue restrictions. Contains natural pyrethrins.

USED IN COMBINATION WITH OTHER INSECTICIDES: PRO-CONTROL PYRETHRIN 6EC may be combined with other insecticides and acaricides where resistance may be a problem and to provide a flushing of insects from hiding and into contact with other spray residues for quicker and more complete control. The application must conform to the accepted use precautions and directions for both products.

Prior to tank-mixing, a compatibility test should be conducted using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture.

Tank mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

Apply 2 to 16 ounces per acre and repeat if required to maintain effective control. Use in sufficient water for thorough coverage of upper and lower leaf surfaces unless otherwise noted. This product may be applied by air in no less than 2 gallons of water per acre and by ground in no less than 10 gallons of water per acre. It is recommended that the final spray mix be buffered to a pH of 5.5 - 7.0.

This concentrate is relatively non-toxic to honey bees. To avoid possible harm to honey bees, it is advisable to apply in the early morning or late evening hours.

USE THROUGH IRRIGATION SYSTEMS (CHEMIGATION)

Apply this product only through sprinkler (including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for the operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the pump motor stops. The irrigation line or water pump must include a functional pressure valve which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must be a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of more dilute suspension per unit of time.

GROWING CROPS (OUTDOORS AND IN GREENHOUSES)

ROOT AND TUBER VEGETABLES: Including (but not limited to) Arracacha, Arrowroot, Purple Arrowroot, Japanese Artichoke, Jerusalem Artichoke, Garden Beets, Sugar Beets, Edible Burdock, Edible Canna, Carrots, Cassava (bitter or sweet), Celeriac (celery root), Chayote, Chervil (turnip rooted), Chicory, Chufa, Dasheen, Ginger, Ginseng, Horseradish, Leren, Parsley (turnip rooted), Parsnip, Potato, Radish, Japanese Radish (Daikon), Rutabaga, Salsify, Black Salsify, Spanish Salsify, Sklrret, Sweet Potato, Tanier, Turmeric, Turnip, Yam (true), Yam Bean.

LEAVES OF ROOT AND TUBER VEGETABLES: Including (but not limited to) Garden Beet, Sugar Beet, Edible Burdock, Carrot, Cassava (bitter or sweet), Celeriac (celery root), Chervil (Turnip Rooted), Chicory, Dasheen (taro), Parsnip, Radish, Japanese Radish (Daikon), Rutabaga, Black Salsify, Sweet Potato, Tanler, Turnip, Yam (true).

BULB VEGETABLES: (*Allium* spp.): Including (but not limited to) Garlic, Great-headed Garlic, Leek, Onion (dry bulb and green), Welch, Shallot.

LEAFY VEGETABLES: Including (but not limited to) Amaranth (leafy Amaranth, Chinese Spinach, Tampala), Arrugula, Cardoon, Celery, Chinese Celery, Celluce, Chervil, Cilantro, Com Salad, Chrysanthemum (edible leaved), Chrysanthemum (garland), Cress (garden, water), Upland Cress (yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Fennel (Florence), lettuce (head and leafy), Orach, Parsley, Purslane (garden & winter), Radicchio, Rhubarb, Spinach, Vine Spinach (Malabar, Indian), Spinach (New Zealand) and Swiss Chard.

Greens, Mustard Spinach and Rape Greens.

LEGUME VEGETABLES (SUCCULENT OR DRIED): Including (but not limited to) Adzuki Beans, Field Beans, Kidney Beans, Lima Beans, Moth Beans, Mung Beans, Navy Beans, Pinto Beans, Rice Beans, Runner Beans, Snap Beans, Tepary Beans, Urd Beans, Wax Beans, Asparagus Beans, Black-eyed Peas, Catjang, Chinese Longbeans, Cowpeas, Crowder Peas, Southern Peas, Yardlong Beans, Broad Beans (Fava Beans), Chick Peas (Garbanzo Beans), Guar, Jackbean (Sword Bean), Lablab Bean (Hyacinth Bean), Lentils, Peas (Edible Pod Pea, Garden Peas, Field Peas, Sugar Snap Peas, English Pea, Snow Pea), Pigeon Peas, Soybeans, Sweet Lupin Beans, White Lupin Beans, White Sweet Lupin and Sword Bean.

FOLIAGE OF LEGUME VEGETABLES: Including (but not limited to) Plant parts of any legume vegetable included in the legume vegetable group that will be used as animal feed including any variety of Beans, Field Peas or Soybeans.

FRUITING VEGETABLES: Including (but not limited to) Eggplant, Ground Cherry, Okra, Pepino, Pepper (Bell Pepper, Chili Peppers, Cooking Peppers, Pimentos, Sweet Peppers), Tomatillo and Tomato.

CUCURBIT VEGETABLES: Including (but not limited to) Balsam Apple, Balsam Pear (Bitter Melon), Chayote, Chinese Waxgourd (Chinese preserving melon), Chinese Cucumber, Citron Melon, Cucumber, Gherkin, Edible Gourds, Muskmelons (including hybrids, Cantaloupe, Casaba, Crenshaw, Golden Pershaw Melon, Honeydew Melons, Honey Balls, Mango Melon, Persian Melon, Pineapple Melon, Santa Claus Melon, Snake Melon), Pumpkin, Squash (summer & winter) and Watermelon (including hybrids).

CITRUS FRUITS: Including (but not limited to) Calamondin, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquats, Lemons, Limes, Mandarin (Tangerine), Orange (sweet & sour), Pummelo, Satsuma Mandarin (Citrus spp. Includes Chironja, Tangelos, Tangors).

POME FRUITS: Including (but not limited to) Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear and Quince.

STONE FRUITS: Including (but not limited to) Apricot, Cherry (sweet & sour), Nectarine, Peach, Plum, Prune, Chickasaw Plum, Damson Plum, Japanese Plum and Plumcot.

SMALL FRUITS AND BERRIES: Including (but not limited to) Blackberry, Blueberry, Cranberry, Currant, Dewberry, Elderberry, Gooseberry, Grape, Huckleberry, Loganberry, Olallie Berry, Raspberry (black & red), Strawberry and Youngberry.

TREE NUTS: Including (but not limited to) Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory Nut, Macadamia Nut (Bush Nut), Pecan, Pistachio, Walnut, Black and English (Persian).

ORIENTAL VEGETABLES: Including (but not limited to) Acerola, Atemoya, Balsam Pear (bitter melon), Carambola, Japanese Artichoke, Chinese Broccoli (Gai Lan), Chinese Cabbage (Bok Choy, Napa), Chinese Mustard Cabbage (Gai Choy), Dasheen, Ginger, Ginseng, Chinese Longbeans, Mung Beans, Citron Melon, Balsam Pear (Bitter Melon), Japanese Radish (Daikon), Chinese Spinach, Chinese Waxgourd, Cilantro, Citron Melon, Rambutan and Water Chestnuts.

SUBTROPICAL FRUITS: Avocado, Banana, Carob, Barbados Cherry, Cherimoya, Dates, Durian (Jackfruit), Feijoa, Figs, Guava, Kiwifruit, Lychee Mango, Papaya, Passion Fruit, Paw Paw, Persimmon, Pineapple and Pomegranate.

ADDITIONAL CROPS: Including Artichoke, Asparagus, Avocado, Coffee, Cotton, Hops, Jojoba, Mushroom, Okra, Peanuts, Pineapple, Safflowers, Sesame, Sugar Cane, Sunflowers and Tea.

CEREAL GRAINS: Including (but not limited to), Barley, Buckwheat, Corn (sweet and field), Millet, Proso, Oats, Pearl, Popcorn, Rice, Rye, Sorghum (milo), Teosine, Triticale, Wheat, Wild Rice.

FORAGE FODDER AND STRAW OF CEREAL GRAINS: Including (but not limited to), Barley, Buckwheat, Corn (Sweet and Field), Millet, Proso, Oats, Pearl, Popcorn, Rice, Rye, Sorghum (Milo), Teosine, Triticale, Wheat and Wild Rice.

GRASSES FOR SEED- FORAGE, FODDER AND HAY: Including (but not limited to) any grass (Gramineal family, (green or cured), except sugarcane and those listed in the cereal grains group) that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage, Bermuda Grass, Bluegrass, Bromegrass and Fescue.

NON-GRASS ANIMAL FEEDS: Including (but not limited to) Alfalfa, Velvet Bean, Clover, Kudzu, Lespedeza, Lupin, Sainfoin, Trefoil, Crown Vetch and Wilk Vetch.

HERBS AND SPICES: Including (but not limited to) Allspice, Angelica, Anise (Anise seed), Annatto, Balm, Basil, Borage, Burnet, Camomile, Caper Buds, Caraway, Black Caraway, Cardamon, Cassia bark, Cassia buds, Catnip,

9/14

Celery Seed, Chervil dried, Chives, Chinese Chive, Clary, Clove Buds, Coriander (cilantro or Chinese parsley leaf), Coriander (cilantro seed), Costmary, Culantro, Cumin, Curry Leaf, Dill (dill weed), Dill (seed), Fennel (Italian and Sweet), Fenugreek, Grains of Paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf & seed), Mace, Marigold, Sweet Marjoram, Wild Marjoram, Mustard (seed), Nasturtium, Nutmeg, Parsley, Oregano, Mint, Paprika, Parsley, Pennyroyal, Pepper (black & white), Poppy seed, Rosemary, Rue, Saffron, Sage, Savory, Summer and Winter Savory, Sweet Bay (Bay Leaf), Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff and Wormwood.

ORNAMENTALS: Including (but not limited to): African Violet, Ageratum, Andromeda, Arborvitae, Ash, Aster, Azalea, Beech, Begonia, Birch, Boxwood, Cacti, Calceolaria, Calendula, Calla, Camellia, Carnation, Ceanothus, Cineraria, Chrysanthemum, Coleus, Cotoneaster, Crabapple, Cyclamen, Cypress, Daffodil, Dahlia, Delphinium, Dogwood, Elm, Euonymus, Fern, Ficus, Fir, Firethorn, Foliage Plants, Forsythia, Fuschia, Gardenia, Geranium, Gladiolus, Gloxinia, Gypsophila, Hawthorn, Hemlock, Hickory, Holly, Honey Locust, Horse Chestnut, Hyacinth, Hydrangea, Iris, Lilies, Maidenhair Fern, Juniper, Larch, Laurel, Lilac, Linden, Marigold, Mimosa (Silk Tree), Myrtle, Narcissus, Oak, Palm, Pansy, Pelargonium, Peony, Petunia, Philodendron, Phlox, Pine, Privet, Pyracantha, Rhododendron, Roses, Rubber Plant, Snapdragon, Stock, Sweet Pea, Tulip, Tulip Tree, Viburnum, Wandering Jew, Willow, Yew, Zinnia.

GREENHOUSE FRUITS, VEGETABLES, FLOWERS AND FOLIAGE PLANTS

USED ALONE: Combine 12 to 24 ounces of PRO-CONTROL PYRETHRIN 6EC with 100 gallons of water for applications with conventional hydraulic sprayers or 1 to 2 teaspoons per gallon of water for applications with compressed air sprayers.

USED IN COMBINATION WITH OTHER INSECTICIDES: To provide quick knockdown of insects when used with a residual insecticide, tank mix 1 to 4 ounces of PRO-CONTROL PYRETHRIN 6EC with the proper amount of companion insecticide in 100 gallons of water and apply with a conventional hydraulic sprayer. Applications must be made in accordance with the more restrictive label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

IMPORTANT NOTE: Plant safety is an important consideration when using insecticides in a greenhouse. However, it is not possible to evaluate the phytotoxicity of PRO-CONTROL PYRETHRIN 6EC towards numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of PRO-CONTROL PYRETHRIN 6EC, treat a limited number of plants and observe for phytotoxicity over a 10 day period.

[6] OUTDOOR USE ON TREES, SHRUBS, FLOWERS AND FOLIAGE PLANTS

USED ALONE: Combine 12 to 24 ounces of PRO-CONTROL PYRETHRIN 6EC with 100 gallons of water for applications with conventional hydraulic and airblast sprayers or 12 to 24 ounces of PRO-CONTROL PYRETHRIN 6EC with 10 gallons of water for applications with low volume mist blowers or 1 to 2 teaspoons per gallon water for applications with compressed air sprayers.

USED IN COMBINATION WITH OTHER INSECTICIDES: To provide quick knockdown of insects when used with a residual insecticide, tank mix 1 to 4 ounces of PRO-CONTROL PYRETHRIN 6EC with the proper amount of companion insecticide in 100 gallons of water (10 gallons of water for low volume application with mist blowers) and apply with conventional hydraulic or airblast sprayers.

Applications must be made in accordance with the more restrictive label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

FOR CONTROL OF GYPSY MOTH CATERPILLARS AND ADULTS: Combine 8 to 12 ounces of PRO-CONTROL PYRETHRIN 6EC with 100 gallons of water for applications with conventional hydraulic sprayers or 8 to 12 ounces of PRO-CONTROL PYRETHRIN 6EC with 10 gallons of water for applications with airblast sprayers. To provide quick knockdown on gypsy moth caterpillars when used with a residual insecticide, tank-mix 1 to 4 ounces of PRO-CONTROL PYRETHRIN 6EC with the proper amount of companion insecticide in 100 gallons of water (10 gallons of water for airblast sprayers) and apply with a conventional hydraulic sprayer.

Applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

[5] INDOOR USE ON TREES, SHRUBS, FLOWERS AND FOLIAGE PLANTS

USED ALONE: Combine 12 to 24 ounces of PRO-CONTROL PYRETHRIN 6EC with 100 gallons of water for applications with conventional hydraulic sprayers or 1 to 2 teaspoons of PRO-CONTROL PYRETHRIN 6EC per gallon of water for applications with compressed air sprayers.

10/14

USED IN COMBINATION WITH OTHER INSECTICIDES: To provide quick knockdown of insects when used with a residual insecticide, tank-mix 1 to 4 ounces of PRO-CONTROL PYRETHRIN 6EC with the proper amount of companion insecticide in 100 gallons of water and apply with a conventional hydraulic sprayer.

Applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

HARVESTED FRUITS AND VEGETABLES

Including Apples, Blackberries, Blueberries, Boysenberries, Cherries, Crabapples, Currants, Dewberries, Figs, Gooseberries, Grapes, Guavas, Loganberries, Mangoes, Muskmelons, Oranges, Peaches, Pears, Peas, Pineapples, Plums, Raspberries and Tomatoes.

DIRECT SPRAY TO FRUITS IN BASKETS, ON TRUCKS OR IN PROCESSING PLANTS: To control *Drosophila* spp., Tephritid spp., Fruit Flies, Vinegar Flies and other nuisance pests, dilute this concentrate at the rate of 1 part with 1,200 parts water (1 pint per 150 gallons or 1 teaspoon per 12.5 pints water). Thoroughly mix the emulsion in the spray tank and treat as follows:

- 1) Apply liberally to fruits and vegetables in baskets, on trucks and in processing plants. Use sprayers at a high pressure for applying at the rate of 5 or 6 pints of diluted spray to a 2 ton load of produce. Direct the spray for maximum coverage of the baskets or hampers. It is important to spray between and beneath the containers. It not only kills the flies, but the emulsion loosens any dead flies so they are readily washed from the fruit.
- 2) Spray the raw stock stacked in the yard.
- 3) Dip baskets in the diluted spray, after dumping the produce, to kill adhering larvae and pupae.

FOR USE IN CANNERIES: The entire space inside of the cannery should be sprayed after washing and cleaning up and just before bringing produce into it, with this product diluted 1 part to 29 parts of water (1 quart with 7.5 gallons water) up to 1 part to 11 parts of water (1 quart with 3 gallons water). Use 1 gallon of the spray per 750 square feet, directing it on walls, ceiling, and floors paying special attention to forcing the spray into all cracks and crevices for the control of ants, roaches, silverfish, crickets, spiders and cheese mites. This same dilution used as a space spray will give excellent control of fruit flies, house flies, hornets, grain moths, gnats, mosquitoes, and skipper flies. Use 1 ounce diluted spray per 1000 cubic feet of space. Do not spray while the plant is in operation as dead flies may fall into containers or the products being processed.

FOR USE WITH HYDROPONICALLY GROWN VEGETABLES AS A WATER SYSTEM TREATMENT: To control aquatic diptera larvae, apply PRO-CONTROL PYRETHRIN 6EC to the water at the rates outlined in the following table:

Pyrethrins Concentration	Pro-Control Pyrethrin 6EC	Amount of Water
0.1 ppm	2.18 fl. oz. (64.6 ml)	10,000 gal.
0.01 ppm	0.218 tsp. (6.46 ml)	10,000 gal.
0.001 ppm	0.022 tsp. (0.646 ml)	10,000 gal.

[3] STORED PRODUCTS

This concentrate can be used at the rate of 1 part to 29 parts up to 1 part to 11 parts water or deodorized base oil can be used on rice, barley, beans, birdseed, buckwheat, cocoa beans, corn, cottonseed, flax, oats, grain nuts, dried fruit, almond nutmeat and shells, walnut nutmeat and shells, pistachios, dried prunes, dried apricots, raisins, figs, wheat, oats, corn, barley, rye, sorghum, wheat, tobacco and peanuts held in storage for control of the accessible stages of Almond Moths, Angoumois Grain Moths, Cadelle Beetles, Cigarette Beetles, Confused Flour Beetles, Flat Grain Beetles, Granary Weevils, Indian Meal Moths, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Saw-toothed Grain Beetles, Square Necked Grain Beetles, and Tobacco Moths.

FOR USE ON SWEET POTATOES IN STORAGE: For control of Fruit Flies and Vinegar Flies dilute this concentrate at 1 part to 19 parts water (6.4 fluid ounces per gallon). Apply as a space fog with a mechanical fogger capable of producing particles of aerosol size at the rate of 1 gallon diluted spray per 100,000 cubic feet of space. Apply only when flying insects are present. Several applications may be necessary during periods of heavy infestation, but do not make more than 10 applications.

SURFACE TREATMENT OF STORED GRAIN AND SEED: To control Indian Meal Moths, Angoumois Grain Moths and Mediterranean Flour Moths, monthly inspections should be made after the grain is placed in storage. If the top two or three inches are infested, dilute 1 part PRO-CONTROL PYRETHRIN 6EC with 19 parts of water and apply at the rate of 1 to 2 gallons per 1,000 square feet of grain. Rake the mixture into the grain to a depth of 4 inches.

FOR USE AS A GRAIN PROTECTANT: This concentrate when diluted with water and sprayed directly on grain will effectively protect the grain against grain storage insects for a full season or approximately 8 months. Dilute at the rate of 1 part to 29 parts water (1 quart with 7.5 gallons water / 1 pint with 3.75 gallons water). Thoroughly mix

the emulsion and apply at the rate of 4 to 5 gallons per 1,000 bushels of grain as it is carried along a belt or as it enters the auger or elevator. This concentrate may be used in combination with a registered fumigant for use on heavily infested stored products.

ON ALMONDS, PEANUTS, AND WALNUTS IN BULK OR IN BAGS: To control stored product insects such as Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cigarette Beetles, Confused Flour Beetles, Drugstore Beetles, Flat Grain Beetles, Granary Weevils, Indian Meal Moths, Lesser Grain Borers, Maize Weevils, Mediterranean Flour Moths, Merchant Grain Beetles, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Sawtoothed Grain Beetles and Square Necked Grain Beetles, dilute 1.5 ounces of PRO-CONTROL PYRETHRIN 6EC per gallon of water and apply as a coarse wet spray over the top of stored nuts or the outside surface of stacked bagged nuts at the rate of 4 gallons per 1,000 square feet. Apply at weekly intervals for about 6 weeks and then at 15 day intervals. The first two applications should be applied at the rate of 4 gallons per 1,000 square feet and subsequent treatments should be applied at the rate of 2 gallons per 1,000 square feet.

FOR USE IN STORAGE SITES: This concentrate can be used to treat grain and seed in warehouse bins and trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment as a clean up prior to using them for storage. In mills and elevators, all grain infested accumulations should be removed from the bin hoppers. All storage areas and conveying equipment should be thoroughly cleaned by sweeping out the waste grain, cobwebs and other debris from the walls and rafters as well as on the floor and door frames with special attention to material lodged in the cracks and crevices. All of the debris should be removed and burned to kill eggs and insects that might be present.

For farms, particular attention should be given to cleaning up around the used feed and grain bags, grain residues from wagons, harvesting equipment and feed troughs. Newly harvested grain should not be placed in the same bin with carry-over grain and all carry-over grain stocks that are not treated with grain protectant should be fumigated. These cleaning operations should be done within two or three weeks before harvest.

After above sanitation measures have been employed, spray all areas prior to use for storage with 1 part to 29 parts water (1 quart with 7.5 gallons water / 1 pint with 3.75 gallons water) up to 1 part to 11 parts (1 quart with 3 gallons water / 1 pint with 1.5 gallons water). Apply at the rate of 1 gallon per 750 square feet on walls, floors, ceilings and partition boards of bins, paying particular attention to forcing the spray into all cracks and crevices.

Monthly inspections should be made. If the top 2 or 3 inches are found to be infested, re-treat applying at the rate of 1 to 2 gallons of diluted material per 1,000 bushels of stored product.

[4] OTHER INDOOR USES

CRAWLING AND FLYING INSECTS: For control of accessible, exposed stages of CRAWLING INSECTS including but not limited to, Ants, Cockroaches, Cadelles, Cigarette Beetles, Confused Flour Beetles, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Grain Mites, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles, Spider Beetles, Yellow Mealworms and FLYING INSECTS including, but not limited to, Angoumois Grain Moths, Cheese Skippers, Fruit Flies, Fungus Gnats, Gnats, House Flies, Indian Meal Moths, Mosquitoes, Mediterranean Flour Moths, Small Flying Moths, Tobacco Moths: dilute 1 part PRO-CONTROL PYRETHRIN 6EC with 11 parts of water or oil (10.67 ounces per gallon) and apply at the rate of 1 ounce per 1,000 cubic feet of space. Direct the spray towards the ceiling and upper corners of the area and behind obstructions. Vacate the treated area and keep the area closed for at least 30 minutes after treatment. Ventilate the area before reoccupying. Repeat treatment as necessary.

AS A SPACE SPRAY IN FOOD AND NONFOOD AREAS OF FOOD PROCESSING PLANTS, INDUSTRIAL INSTALLATIONS, BAKERIES, RICE AND WHEAT MILLS, RESTAURANTS, TOBACCO WAREHOUSES, GRAIN ELEVATORS, HOMES AND WAREHOUSES: To kill flying insects such as fruit flies, house flies, hornets, wasps, grain moths, gnats, mosquitoes and skipper flies, dilute this concentrate at the rate of 1 part with 29 parts water (1 quart with 7.5 gallons water / 1 pint with 3.75 gallons water) up to 1 part to 11 parts water (1 quart with 3 gallons water / 1 pint with 1.5 gallons water). Use at the rate of 1/2 to 1 ounce of diluted spray per 1,000 cubic feet of space. Direct the space treatment upward whenever practical, and keep doors and windows closed for at least 10 minutes after application. The use of this product in food processing or food handling establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatments. All food processing surfaces should be covered during treatment or thoroughly cleaned before use.

Where oil residues are not undesirable, this product can be diluted at the rate of 1 part to 29 parts up to 1 part to 11 parts in deodorized base oil instead of water and applied as a space spray with any good type applicator such as mechanical or ULV fogger capable of producing particles of aerosol size.

USE AS A SURFACE SPRAY IN HOMES, RESTAURANTS, FOOD PRECESSING PLANTS, INDUSTRIAL INSTALLATIONS AND WAREHOUSES to control accessible, exposed stages of crawling insects including, but not limited to, Ants, Cockroaches, Cadelles, Cigarette Beetles, Confused Flour Beetles, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Grain Mites, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Spider Beetles and Yellow Mealworms: dilute 1 part PRO-CONTROL PYRETHRIN 6EC with 59 parts water and

12/14

apply at the rate of 1 gallon to 750 square feet, paying special attention to force the spray into all cracks and crevices.

Except in Federally inspected meat and poultry plants, food processing operations may continue when this product is applied as a surface spray with care and in accordance with the directions and precautions given above.

FOR USDA INSPECTED FACILITIES: To control accessible, exposed stages of crawling insects including, but not limited to, Ants, Cockroaches, Caddisflies, Cigarette Beetles, Confused Flour Beetles, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Grain Mites, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Spider Beetles and Yellow Mealworms: dilute 1 part PRO-CONTROL PYRETHRIN 6EC with 19 parts of water and apply at the rate of 1 gallon to 750 square feet, paying special attention to force the spray into all cracks and crevices.

[7] OUTDOOR USE AROUND HOMES AND OTHER BUILDINGS

In grassy undeveloped areas use this concentrate at 1 part to 59 parts water to control foraging fire ants. Also spray grassy areas around yard borders liberally to control ticks that may carry Lyme Disease.

[8] OUTDOOR USE - MOSQUITO CONTROL

PRO-CONTROL PYRETHRIN 6EC may be used for mosquito control programs involving residential, industrial, recreational and agricultural areas as well as swamps, marshes, (and) overgrown waste areas, roadsides and pastures where adult mosquitoes occur. PRO-CONTROL PYRETHRIN 6EC may be used over agricultural crops. For best results, apply when meteorological conditions create a temperature inversion and wind speed does not exceed 5 miles per hour. The application should be made so the wind will carry the insecticidal fog into the area being treated. Treatment may be repeated as necessary to achieve the desired level of control.

When used in cold aerosol generators, mechanical or electronic foggers, or misting equipment that produce a fog with the majority of droplets in the 5 - 50 micron range, PRO-CONTROL PYRETHRIN 6EC should be diluted with light mineral oil (specific gravity of approximately 0.8 at 60°F, boiling point: 500-840°F). An N.F. grade oil is preferred.

GROUND APPLICATION: To control adult mosquitoes and biting flies, apply up to 0.0025 pounds of pyrethrins per acre (use a 300 foot swath width for acreage calculations). Treatment of vegetation at the water's edge is permitted by allowing the wind to move the product into the vegetation.

TRUCK-MOUNTED ULV APPLICATION: Dilute 5 parts of PRO-CONTROL PYRETHRIN 6EC with 1 part of oil and apply at the rate of 2 to 2.25 ounces per minute while the machine is traveling 5 miles per hour. The nozzle should be positioned approximately 30 degrees above the horizontal off the side of the truck bed. The delivery rate and truck speed may be varied as long as the application rate is 0.002 to 0.0025 pounds of pyrethrins per acre (use a 300 foot swath width for acreage calculations).

BACKPACK SPRAYER APPLICATION: Apply 0.002 to 0.0025 pounds of pyrethrins per acre. Dilute 1 part PRO-CONTROL PYRETHRIN 6EC with 12 parts of oil and apply at the rate of 7 ounces per acre (based on a 50 foot swath, 7 ounces should be applied while walking 870 feet).

AERIAL APPLICATION (FIXED WING AND HELICOPTER): To control adult mosquitoes and biting flies, apply up to 0.0025 pounds of pyrethrins per acre with equipment designed and operated to produce a ULV spray application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry place. Keep container closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Empty container by using the product in accordance with the label directions. Do not reuse this container! Place empty container in trash or offer for recycling if available. If container is partly filled, call your local solid waste agency or call 1-800-CLEANUP for disposal instructions. **NEVER PLACE UNUSED PRODUCT DOWN ANY INDOOR OR OUTDOOR DRAIN!**

A Prescription Treatment® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.

3568 Tree Court Industrial Blvd.

St. Louis MO 63122-6682

www.wmmg.com

© 2006 Whitmire Micro-Gen Research Laboratories, Inc.

13/14



Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
Attn: Richard Gebken PM 10
Room 266A, Crystal Mall #2
1801 South Bell Street
Arlington, VA 22202

February 2, 2006

**RE: NOTIFICATION per PR notice 98-10 of Reformatting of label text
Pro-Control Pyrethrin 6EC
EPA Reg. No: 499-475**

Dear Mr. Gebken,

Enclosed in support of above action is a completed application and two (2) copies revised labeling [one copy has been highlighted (new or revised text only) for easy reference] for the referenced product.

Reformatted label to include front panel, back panel, and complete label, and included the following statements:

- ⌘ Addition of 'First Aid and' to the statement on Front Panel "See back panel for First Precautionary Statements".
- ⌘ Addition of the Alternate Text statement on front panel "See inside booklet for First Aid Precautionary Statements and Directions for Use.
- ⌘ Addition of 'full label' and 'and Directions for Use' on back panel, to the statement "See full label for complete Precautionary Statements and Directions for Use".

We have also added alternate dilution instructions for pint size containers; i.e. wherever the label reads "1 quart with 7.5 gallons water" or "1 quart with 3 gallons water", it now reads "1 quart with 7.5 gallons water / 1 pint with 3.75 gallons water" or "1 quart with 3 gallons water / 1 pint with 1.5 gallons water" in the directions for use to accommodate the different container sizes for this product. The different container sizes are 6-64 oz, 1 gallon, and 55 gallon.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 156.10 and 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notices 83-1, 84-1, 98-10 and 2001-6 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

14/14


February 2, 2006
Page 2

Also enclosed is a self-addressed, "No Postage Necessary" postcard that, when filled out, would indicate the status of this notification. Please take a moment to fill in the appropriate information and drop it in the mail for our records. Your assistance in this regard is certainly appreciated.

Thank you for your assistance in the review and approval process. Please do not hesitate to contact me at (636) 861-4272 or, via electronic mail, at patricia.hodge@wmmg.com, should you have any questions or require additional information.

Best Regards,

WHITMIRE MICRO-GEN RESEARCH LABORATORIES, INC



Patricia Hodge
Registration Specialist
encl.

