

499-477

11/15/2005

1/10

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 499-477	2. EPA Product Manager 	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Optachield CS Controlled Release Premise Insecticide-AG	PM# 	
5. Name and Address of Applicant (Include ZIP Code) Whitmire Micro-Gen Research Laboratories Inc. 3568 Tree Court Industrial Blvd. St. Louis, MO 63122-6682 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

NOTIFICATION
NOV 15 2005

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of relocation of Precautionary Statement to its preferred location before "Hazards to Humans" and modification of "Storage and Disposal" Statement. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the CSF of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:					
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. _____	No. per container _____	If "Yes" Package wgt _____	No. per container _____
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 		5. Location of Label Directions <input type="checkbox"/> _____	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Patricia A Hodge	Title Registration Specialist	Telephone No. (Include Area Code) 	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Specialist		
4. Typed Name Patricia A Hodge	5. Date 10/26/05		

210

v3.1 10/05

[FRONT PANEL]

NOTIFICATION

NOV 15 2005

**OPTASHIELD® CS
CONTROLLED RELEASE PREMISE INSECTICIDE
Microencapsulated Insecticide**

Recommended for Use by Commercial Applicators

For broad spectrum residual control of crawling, flying, and wood infesting insect pests, indoors, and outdoors.

Optashield CS is intended for use as a general surface, spot, Crack & Crevice® or void application in, on and around buildings and structures and their immediate surroundings and in trucks and trailers and other transportation vehicles.

PERMITTED AREAS OF USE INCLUDE:

Calf Hutches, Calving Pens and Parlors, Dairy Areas, Dog Kennels, Hog barns, Homes, Horse Barns, Milk Rooms, Poultry Houses, Trucks/Trailers, Warehouses, and Other Commercial and Industrial Buildings.

ACTIVE INGREDIENT

Cyfluthrin 6.0%

OTHER INGREDIENTS: 94.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN


CAUTION

See back panel for additional Precautionary Statements.

**EPA Reg. No. 499-477 • EPA Est. No. 499-MO-1
NET CONTENTS: 3.75 Qt. (120 fl. oz.)**



**WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.
800-777-8570**



[BACK PANEL]

OPTSHIELD® CS**FIRST AID**

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-225-3320 for emergency medical treatment advice.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Do not apply to humans, their clothing or bedding. Do not allow children or pets to contact treated surfaces until spray has dried. Do not contaminate food or use on household pets.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

See supplemental label for complete Directions for Use and Storage and Disposal.

Cover fish tanks prior to application. Do not apply this product in patient rooms while occupied or in any rooms while occupied by the elderly or infirm. Do not apply to classrooms when in use. Avoid contamination of food, feedstuffs, or water supply. Do not contaminate food preparation surfaces, kitchen utensils, dishes, or feed storage containers. All food/feed contact surfaces and cooking utensils should be covered during treatment or thoroughly cleaned before using. Do not spray where electrical short circuits might result, such as wall outlets, conduits, etc.

USE IN AND AROUND LIVESTOCK HOUSING

For use by pest control operators or livestock producers in livestock housing structures to control flying and crawling insects. Apply as a General Surface Application and/or Spot or Crack and Crevice® or Void Application. Make applications with a low pressure sprayer with suitable nozzles to achieve uniform coverage or with fogging or misting equipment. Control will be enhanced when facilities are cleaned and indoor treatments are supplemented with outdoor perimeter applications. Do not make indoor applications in areas of facility where animals other than cattle or horses are present. Do not apply directly to animal feedstuffs or watering equipment.

FOGGING OR MISTING APPLICATION: Remove all animals from the facility to be treated. Cover all feeders and waterers, and close all vents, windows and doors. After application, wait one half hour and ventilate facility before entering.

FLIES: For residual control of flies such as house flies and stable flies on outdoor surfaces and in animal areas such as dairy barns, dairy areas, milkrooms, calf hutches, calving pens and parlors, poultry houses, hog barns,

4/10

v3.1 10/05

horse barns, and kennels. As a general surface application treat normal fly resting areas such as ceilings, screens, upper corners of the areas, around windows, doors and light fixtures and other surfaces known to attract flies. For best results initiate treatments at the beginning of the fly season and repeat during periods of heavy fly activity. Treat no more than twice per week and at least 3 days apart.

DARKLING BEETLES: For residual control of darkling beetles and other insects in poultry houses, at cleanouts and in between poultry grow-out periods. Use as a General Surface application on floors, litter, walls, ceilings, posts and other surfaces where pests live and cause damage. Use 1.0 to 2.0 oz. per 1,000 sq. ft. Use the high rate for heavy infestations.

A PRESCRIPTION TREATMENT® brand insecticide from:
Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
www.wmmg.com
© 2005 Whitmire Micro-Gen Research Laboratories, Inc.

[SUPPLEMENTAL LABEL]

OPTASHIELD CS
CONTROLLED RELEASE PREMISE INSECTICIDE

Microencapsulated Insecticide

Recommended for Use by Commercial Applicators

For broad spectrum residual control of crawling, flying, and wood infesting insect pests, indoors, and outdoors. Optashield CS is intended for use as a general surface, spot, Crack & Crevice® or void application in, on and around buildings and structures and their immediate surroundings and in trucks and trailers and other transportation vehicles.

PERMITTED AREAS OF USE INCLUDE:

Calf Hutches, Calving Pens and Parlors, Dairy Areas, Dog Kennels, Hog barns, Homes, Horse Barns, Milk Rooms, Poultry Houses, Trucks/Trailers, Warehouses, and Other Commercial and Industrial Buildings.

ACTIVE INGREDIENT:

Cyfluthrin 6.0%

OTHER INGREDIENTS: 94.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-225-3320 for emergency medical treatment information.

See page 2 for additional Precautionary Statements.

EPA Reg. No. 499-477 • EPA Est. No. 499-MO-1



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.
800-777-8570

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Do not apply to humans, their clothing or bedding. Do not allow children or pets to contact treated surfaces until spray has dried. Do not contaminate food or use on household pets.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Cover fish tanks prior to application. Do not apply this product in patient rooms while occupied or in any rooms while occupied by the elderly or infirm. Do not apply to classrooms when in use. Avoid contamination of food, feedstuffs, or water supply. Do not contaminate food preparation surfaces, kitchen utensils, dishes, or feed storage containers. All food/feed contact surfaces and cooking utensils should be covered during treatment or thoroughly cleaned before using. Do not spray where electrical short circuits might result, such as wall outlets, conduits, etc.

GENERAL INFORMATION

OPTASHIELD CS is a liquid capsule suspension containing 6% of cyfluthrin which will provide extended residual control of the pests listed on this label. This product may be applied using compressed air, backpack, or power spray equipment. Thoroughly clean spray equipment before using this product. Shake concentrate well before mixing. First add approximately one-half the required water to the tank, then add the proper amount of OPTASHIELD CS. Agitate, add the remaining water and thoroughly agitate the solution again. When emptying the bottle, triple rinse with water and add the rinsate to the spray tank. Continue to agitate solution occasionally during use to ensure even coverage. Shake or re-agitate solution if spraying is interrupted or if dilution is left in sprayer overnight. If spray screens are used they should be 50 mesh or larger. Water may be used at temperatures up to 160° F.

OPTASHIELD CS may be sprayed on any surface which will not be damaged or stained by water. Heavy applications may leave a visible deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

Do not apply this product through any type of irrigation system and do not formulate this product into other end-use products.

Do not apply water based sprays of this product to conduits, motor housing, junction and switch boxes or other electrical equipment because of possible shock hazard.

General Surface Application

Use 1.0 to 2.0 fl. oz. of OPTASHIELD CS per 1,000 sq. ft. in sufficient water to adequately cover the area being treated, without dripping or run off. The higher rate should be used for clean-outs, for longer residual control or for heavy insect infestations.

Mixing Chart

Low Rate - 1.0 fl. oz. per 1,000 sq. ft.

OPTASHIELD CS	Finished Spray	Spray Rate	Coverage
1.0 fl. oz.	1 gallon	1 gal/1,000 sq. ft.	1,000 sq. ft.
1.0 fl. oz.	2.5 gallons	2.5 gal/1,000 sq. ft.	1,000 sq. ft.

20.0 fl. oz.	20 gallons	1 gal/1,000 sq. ft.	20,000 sq. ft.
20.0 fl. oz.	50 gallons	2.5 gal/1,000 sq. ft.	20,000 sq. ft.
High Rate - 2.0 fl. oz. per 1,000 sq. ft.			
OPTASHIELD CS	Finished Spray	Spray Rate	Coverage
2.0 fl. oz.	1 gallon	1 gal/1,000 sq. ft.	1,000 sq. ft.
2.0 fl. oz.	2.5 gallons	2.5 gal/1,000 sq. ft.	1,000 sq. ft.
40.0 fl. oz.	20 gallons	1 gal/1,000 sq. ft.	20,000 sq. ft.
40.0 fl. oz.	50 gallons	2.5 gal/1,000 sq. ft.	20,000 sq. ft.

Apply as a coarse wet spray.

Applications can be made to walls, floors, ceilings, in and around cupboards, between, behind beneath equipment, appliances, around floor drains, window and door frames and light fixtures, and on the underside of shelves, drawers and in similar areas. Applications may be made to floor surfaces, along walls and around air ducts.

As a perimeter spray to help prevent pest infestations treat a band of soil 6 to 10 feet wide around and adjacent to the building or structure. Also treat the building foundation and sides where pests are active and may find entrance. Treat other insect entry points such as doors, windows, under siding and eaves of structure.

Spot, or Crack and Crevice® or Void Applications

Mix 1.0 to 2.0 fl. oz. of OPTASHIELD CS in one gallon of water. The higher rate should be used for clean-outs, for longer residual control or for heavy insect infestations.

	OPTASHIELD CS	Finished Spray	A.I. Concentration
Low Rate	1.0 fl. oz.	1 gallon	0.05%
High Rate	2.0 fl. oz.	1 gallon	0.10%

Use a low pressure system with a pin point or variable pattern nozzle to apply the suspension in specific areas such as cracks and crevices in or behind baseboards, in floor walls and expansion joints, areas around water and sewer pipes, wall voids or voids in table legs and equipment where pests can hide as well as other similar areas. Spot treatments may also be made to areas, including but not limited to, storage areas, closets, doors and windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment, the underside of shelves, drawers and similar areas.

INDOOR TREATMENTS

For general indoor pests and occasional invaders make applications in a manner as previously described under General Surface Application and/or Spot, or Crack and Crevice or Void Applications. Particular attention should be given to treating entry points and harborage areas.

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food/feed is exposed or facility is in operation. Serving areas include areas where prepared foods are served such as dining rooms and exclude areas where foods may be prepared or held. In the home, all food/feed processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Nonfood/feed areas are areas such as garbage rooms, rest rooms, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Flies: Make spot applications of OPTASHIELD CS on surfaces where flies rest in the dark at night or frequent during the day, such as window frames, screens, trash cans, ceilings and upper wall surfaces.

Ants: Inject cracks, crevices and voids where ants may be traveling and/or harboring, including but not limited to, around doors, window frames, and other structural cracks, crevices and voids. Spot treatment of trails and feeding areas may also be made.

Other crawling insects, including but not limited to Centipedes, Cockroaches, Crickets, Dermestids, Earwigs, Firebrats, Millipedes, Psocids, Silverfish, Sowbugs, Spiders: Make spot treatment and/or Crack and Crevice injections into hiding places such as cracks and crevices, void areas, moist areas, openings around pipes and sinks and under refrigerators.

Ticks: Apply as a spot and/or Crack and Crevice treatment in infested and egg laying areas such as pet resting quarters, nearby cracks and crevices, behind baseboards, door and window frames. Replace pet's old bedding with clean fresh bedding after treatment. Do not treat pets with this product.

OUTDOOR TREATMENTS

OPTASHIELD CS can be used as a General Surface Application on and around outdoor surfaces and/or a Spot or Crack and Crevice or Void Application treatment of buildings, porches, window frames, eaves, patios, garages, garbage dumps, and other areas where pests congregate or enter premises.

As a perimeter treatment make applications in a manner previously described under General Surface Application.

Flies: Treat outside surfaces of screens, doors, window frames or wherever flies enter. Also treat surfaces on porches, in garages, and other places where insects light or congregate. Treat normal fly resting areas such as ceilings, upper corners of the area, around windows and doors, and other surfaces. Treat inside and outside surfaces of dumpsters and other trash holding containers. Treat trash storage areas. Treat so as to lightly moisten the target surface. For best results, begin treatments at the start of the fly season and repeat during periods of heavy fly activity. Treat no more than twice per week and at least 3 days apart.

Crawling Insects, including but not limited to Ants, Cockroaches, Crickets, Earwigs, Sowbugs, and Scorpions: For treatment of infestations of these insects in outdoor areas which are landscaped, planted, mulched or unmaintained. For ants treat mounds and runways. Repeat applications as infestation warrants and as reinfestations occur. Apply OPTASHIELD CS to cracks and crevices where insects may enter premises. Areas to be treated include, but are not limited to, weep holes, under siding, wall voids, soffits, around attic vents, cracks and crevices openings around windows, doors and pipes. Treatment of infested fences and tree holes is permitted.

Fleas: Treat infested areas. OPTASHIELD CS will kill fleas on contact but does not provide residual control.

INDOOR AND OUTDOOR TREATMENTS

Carpenter Ants: Make spot and/or Crack and Crevice applications to any trails, inside or outside around doors and windows, and other places where ants enter premises. Areas which may be treated include but are not limited to atriums, crawl spaces, ceilings, garages, soffits, wall voids, weep holes, and other void areas. Where possible, treat ant nest or infested wood directly.

Bees, wasps and yellow jackets: As a preventative or remedial treatment to help prevent these insects from building nests or workers from congregating on structures follow perimeter treatment directions under General Surface Application section of label. Pay particular attention to soffit and eave areas.

For treatment of a yellow jacket or wasp nest, liberally treat the nest, inside the nest for yellow jackets and the surrounding area. For treatment of nests inside wall voids inject into the nest sufficient quantity to treat the complete nest. More than one point of application maybe required. Most applications should require no more than half a gallon of diluted material. Pay particular attention to what the water may contact such as electrical wiring. Do not use where there is a shock hazard.

OPTASHIELD CS DOES NOT PROVIDE INSTANT KNOCKDOWN. THE BEST TIME TO TREAT IS LATE IN THE DAY. COLONY SHOULD BE DEAD IN 2-3 DAYS. DEPENDING ON THE SITUATION PROTECTIVE EQUIPMENT FOR THE APPLICATOR MAY BE REQUIRED.

USE IN AND AROUND LIVESTOCK HOUSING

For use by pest control operators or livestock producers in livestock housing structures to control flying and crawling insects. Apply as a General Surface Application and/or Spot or Crack and Crevice or Void Application. Make applications with a low pressure sprayer with suitable nozzles to achieve uniform coverage or with fogging or misting equipment. Control will be enhanced when facilities are cleaned and indoor treatments are supplemented with outdoor perimeter applications.

Do not make indoor applications in areas of facility where animals other than cattle or horses are present. Do not apply directly to animal feedstuffs or watering equipment.

Fogging or misting application: Remove all animals from the facility to be treated. Cover all feeders and waterers, and close all vents, windows and doors. After application, wait one half hour and ventilate facility before entering.

Flies: For residual control of flies such as house flies and stable flies on outdoor surfaces and in animal areas such as dairy barns, dairy areas, milkrooms, calf hutches, calving pens and parlors, poultry houses, hog barns, horse barns, and kennels. As a general surface application treat normal fly resting areas such as ceilings, screens, upper corners of the areas, around windows, doors and light fixtures and other surfaces known to attract flies.

For best results initiate treatments at the beginning of the fly season and repeat during periods of heavy fly activity. Treat no more than twice per week and at least 3 days apart.

Darkling Beetles: For residual control of darkling beetles and other insects in poultry houses, at cleanouts and in between poultry grow-out periods.

9/10

v3.1 10/05

Use as a General Surface application on floors, litter, walls, ceilings, posts and other surfaces where pests live and cause damage.

Use 1.0 to 2.0 oz. per 1,000 sq. ft. Use the high rate for heavy infestations.

PESTS CONTROLLED BY OPTASHIELD CS:

	Indoor	Outdoor		
Crawling	Ants	Firebrats	Ants	Fleas
	Cent pedes	Millipedes	Cockroaches	Scorpions
	Cockroaches	Psocids	Crickets	Ticks
	Darking Beetles	Silverfish	Fire Ants	
	Dermestids	Sowbugs		
	Earwigs	Spiders		
	Flies		Bees	
Flying	Gnats		Flies	
	Mosquitoes		Hornets	
	Moths		Moths	
	Yellow jackets		Yellow jackets	
	Wasps		Wasps	
Wood Infesting Pests	Carpenter Ants			
	Beetles			

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides or fertilizers. Do not store above 122°F for extended periods of time. Preferred storage is above 32°F. Storage below 32°F may result in solidification. If warmed to above 32°F, product will return to original form. Freezing does not adversely affect this product. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL (1 gal.): Empty container by using the product in accordance with the label directions. Do not reuse container! Place empty container in trash or offer for recycling if available. If container is partly filled, call your local solid waste agency or call 1-800-CLEANUP for disposal instructions. **NEVER PLACE UNUSED PRODUCT DOWN ANY INDOOR OR OUTDOOR DRAIN!**

CONTAINER DISPOSAL (Greater than 1 gal.): Empty container by using the product in accordance with the label directions. Do not reuse container! Place empty container in trash or offer for recycling if available. If container is partly filled, call your local solid waste agency or call 1-800-CLEANUP for disposal instructions. **NEVER PLACE UNUSED PRODUCT DOWN ANY INDOOR OR OUTDOOR DRAIN!**

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.

3568 Tree Court Industrial Blvd.

St. Louis MO 63122-6682

www.wmmg.com

© 2005 Whitmire Micro-Gen Research Laboratories, Inc.

10/10



Document Processing Desk (NOTIF)
Office of Pesticide Programs(7504C)
U.S. Environmental Protection Agency
Room 266A, Crystal Mall #2
1801 South Bell Street
Arlington, VA 22202

NOTIFICATION

NOV 15 2005

October 26, 2005

RE: NOTIFICATION per PR notice 98-10 and 2001-6 of Movement of Precautionary Statement and modification of "Storage and Disposal " Statement Optashield CS Controlled Release Premise Insecticide-AG EPA Reg. No: 499-477

Dear Sir or Madam:

Enclosed in support of above action is a completed application and two (2) copies revised labeling [one copy has been highlighted (new or revised text only) for easy reference] for the referenced product.

- ✓ Relocation of "Precautionary Statement" to its preferred location before "Hazards to Humans"
- ✓ Modification of "Storage and Disposal" to be consistent with PRN 2001-6
- ✓ Movement of "Storage and Disposal" to end of DIRECTIONS FOR USE as recommended by EPA.

This notification is consistent with the provisions of PR Notice 98-10 and 2001-6 and EPA regulations at 40 CFR 156.10 and 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notices 83-1, 84-1, 98-10 and 2001-6 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Also enclosed is a self-addressed, "No Postage Necessary" postcard that, when filled out, would indicate the status of this notification. Please take a moment to fill in the appropriate information and drop it in the mail for our records. Your assistance in this regard is certainly appreciated.

Thank you for your assistance in the review, and approval process. Please don't hesitate to contact me at (636) 861-4272 or, via electronic mail, at patricia.hodge@wmmg.com, should you have any questions or require additional information.

Best Regards,

WHITMIRE MICRO-GEN RESEARCH LABORATORIES, INC

Patricia Hodge
Registration Specialist

patricia.hodge@wmmg.com
encl.

3568 Tree Court Industrial Boulevard
St. Louis, MO 63122-6682
800-777-8570 • FAX 636-225-3739
www.wmmg.com

