

**ACCEPTED**

SEP 16 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. **499-483**

Use TC 179 at the rate of one can (9 fluid oz.) for up to 2,100 sq. ft. Carpeted areas may be vacuumed before treatment and after space treatment has dried. Treat infested areas or areas which could be infested; these include rugs, carpets, upholstered furniture, pet beds and pet resting areas. Hold can at arms length and approximately 36 inches away from the surface being treated. Use a sweeping motion to apply product and then vacate treated area. An area of 80 - 100 sq. ft. can be treated in approximately 10 seconds with TC 179. When treating upholstered furniture, treat areas where flea development occurs including under cushions and beneath furniture.

Prior to treating, cover aquariums and fish bowls and remove birds from area. Treated areas should be vacated during application. **EXCEPT FOR APPLICATOR DO NOT PERMIT HUMANS OR PETS TO CONTACT TREATED SURFACES UNTIL THE SPRAY HAS DRIED.** Do not treat pets with this product.

**NOTICE:** In transportation equipment, application of products onto certain plastics may result in whitening or dulling of the treated surface. Apply only to carpets and cloth upholstered areas, or test in an inconspicuous area before treating.

TC 179 has been tested on several types of carpet, fabric and other household furnishings without adverse effects. In a few instances, waxed surfaces have been whitened or dulled. Holding the can 36 inches away from the target surface normally prevents any whitening from occurring. Because of the wide variety of floor types and finishes, treat a small inconspicuous area before treating the entire area. Avoid spray contact with antique finishes.

**STORAGE & DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool dry area away from heat or open flame and inaccessible to children.

**PESTICIDE DISPOSAL:** Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** This container may be recycled in the few but growing number of communities where steel aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the label (DO NOT PUNCTURE!). If recycling is not available, wrap the container and discard in the trash.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



**A PRESCRIPTION TREATMENT® brand insecticide from:**  
Whitmire Micro-Gen Research Laboratories, Inc.  
3568 Tree Court Industrial Blvd.  
St. Louis MO 63122-6682  
www.wmrg.com

© 2002 Whitmire Micro-Gen Research Laboratories, Inc.



**Contains NYLAR\***

**KILLS:** Active flea infestations and prevents infestations from developing.

**FOR USE IN:** Apartments, Automobiles, Homes, Hospitals and Nursing Homes, Hotels, Kennels, Motels, Offices, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks), Warehouses, Utilities, Veterinary Clinics and Other Commercial and Industrial Buildings.

**ACTIVE INGREDIENTS:**

Permethrin ..... 2.50%  
Pyriproxyfen ..... 0.18%  
Piperonyl Butoxide, Technical ..... 10.00%

**OTHER INGREDIENTS:** ..... 87.32%

Contains petroleum distillate. **TOTAL: 100.00%**

\*NYLAR - Reg. TM of McLaughlin Gormley King Co.

**EPA REG. No. 499-483**  
**EPA Est. No. 499-MO-1**  
**KEEP OUT REACH OF CHILDREN**  
**CAUTION**

See side panel for additional Precautionary Statements.

**NET WEIGHT: 9 oz.**



**WHITMIRE MICRO-GEN**  
RESEARCH LABORATORIES, INC.  
800-777-8570

**PRECAUTIONARY STATEMENTS**

**FIRST AID**

**IF SWALLOWED:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-225-3320 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** Contains petroleum distillate - vomiting may cause aspiration pneumonia.

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Do not allow children or pets to contact treated surfaces until spray has dried. Avoid contamination of food, utensils, and food preparation surfaces. Remove pets and birds and cover or remove aquariums before use.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and other aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Apply this product only as specified on the label. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

**Extremely flammable.** Contents under pressure. Keep away from heat, sparks and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray directly into any electronic equipment such as radios, televisions, computers, etc.

**DIRECTIONS FOR USE**

**IT IS A VIOLATION TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

TC 179 Flea Killer and Insect Growth Regulator kills fleas and prevents flea infestations for up to 7 months. Adult fleas may be seen in the treated areas when brought in on infested animals or when adults emerge from pupal cases. However, a population will not develop in the treated area.