

499-503

12/07/2005

1/5



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Avenue NW
Washington, D.C. 20460

EPA Reg. Number:

499-503

Date of Issuance:

12/07/05

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Whitmire Micro-Gen TC 240
Insecticide

Name and Address of Registrant (include ZIP Code):

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Boulevard
St. Louis, MO 63122-6682

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following labeling changes before you release the product for shipment:
Revise the EPA Registration Number to read: "EPA Registration No. 499-503"

Signature of Approving Official:

George LaRocca
George LaRocca
Product Manager (RM 13)
Insecticide Branch
Registration Division (7505C)

Date:

December 07, 2005

2/5

3. Submit to the Agency, data on one year storage stability testing (830.6317), and corrosion characteristics study (830-6320) at 0, 3, 6, 9 and 12 months of warehouse storage.
4. Submit two (2) copies of the final printed label bearing the above stated revisions prior to releasing this product for sale.

If you fail to comply with the above stated conditions, this registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the stamped label is enclosed for your records. If you have any questions concerning this action, please contact Dr. B. A. Akinlosotu at (703) 605-0653.

Enclosure:

and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For closed voids calculate the void's cubic area and treat at the rate of one to five seconds per three cubic feet. Several holes may be required in long-running voids.

SPOT TREATMENT: Use TC 240 with the supplied actuator and injection tubes or Whitmire Micro-Gen System Equipment (such as System III.) Hold container upright while spraying. Direct nozzle approximately 12 inches from surface to be treated. Apply directly on insects in these locations when possible.

INDOOR TREATMENTS

COCKROACHES, DERMESTIDS, SCORPIONS, SPIDERS, EARWIGS, CRICKETS, SOWBUGS, MILLIPEDES, CENTIPEDES: Apply as a Crack & Crevice treatment using the prescribed dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewer, floor drains and meter boxes. Spot treatments may also be made to areas including, but not limited to, storage areas, closets, around water pipes, doors and windows, around refrigerators, cabinets, sinks, stoves and other equipment, shelves, drawers and similar areas.

TICKS: Apply as a Crack & Crevice treatment or spot treatment into infested and egg laying sites near pet resting areas including, but not limited to cracks and crevices, behind baseboards, door and window frames. Do not use on pets. Also, treat your yard with a product registered for yard application and your pet with a product registered for pet treatment.

BOOKLICE & SILVERFISH: Apply as a Crack & Crevice treatment into cracks and crevices of baseboards, door frames, bookcases and other harborage sites. Spot treatments may also be made to areas including, but not limited to, storage areas, closets, around water pipes, doors and windows, around refrigerators, cabinets, sinks, stoves and other equipment, shelves, drawers and similar areas.

ANTS: Apply as a Crack & Crevice treatment into cracks, crevices and voids where ants may be traveling and/or harboring, including, but not limited to, around doors, window frames, wall voids and other structural cracks and crevices. Spot treatments may also be made to areas including, but not limited to, storage areas, closets, around water pipes, doors and windows, around refrigerators, cabinets, sinks, stoves and other equipment, shelves, drawers and similar areas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.
PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Do not puncture or incinerate! Empty container by using the product according to the label directions. Offer empty container for recycling, if available, or place in trash if allowed by state and local regulations. If container is partly filled, contact your local solid waste agency or call 1-800-CLEANUP for disposal instructions.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



A **PRESCRIPTION TREATMENT**® brand insecticide from:
 Whitmire Micro-Gen Research Laboratories, Inc.
 3558 Tree Court Industrial Blvd
 St. Louis MO 63122-6682
 www.wmrg.com

© 2004 Whitmire Micro-Gen Research Laboratories, Inc.
 TC 240 Container v1 0.3 (3/04)



KILLS:

Ants, Bed Bugs, Bees, Booklice, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs, Elmleaf Beetles, Fire Ants, Flies, Flour Beetles, Grain Weevils, Indian Meal Moths, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders, Springtails, Above-Ground Termites**, Ticks, Trogoderma (Cabinet, Khapra, and Warehouse Beetles), Wasps and Yellow Jackets.

**Refer to termites section on complete label for restrictions regarding these pests.

FOR USE IN AND AROUND:

Apartments, Campgrounds, Food Storage Areas, Homes, Hospitals*, Hotels, Meat Packing and Food Processing Plants, Motels, Nursing Homes*, Resorts, Restaurants and other Food Handling Establishments, Schools*, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes*), Utilities, Warehouses, and other Commercial and Industrial Buildings.

* Refer to "General Use Restrictions" section for restrictions regarding these sites.

ACTIVE INGREDIENT:

A-Cyhalothrin	0.05%
Prallethrin	0.03%
OTHER INGREDIENTS:	99.92%
Contains petroleum distillate.	TOTAL 100.00%

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

See side panel for additional Precautionary Statements.

EPA Reg. No. 499-LNG • EPA Est. No. 499-MO-1

NET WEIGHT: 8 oz.

* [ALTERNATE TEXT: "NET WEIGHT 1 lb 1.5 oz. (17.5 oz.)"]



**WHITMIRE MICRO-GEN
 RESEARCH LABORATORIES INC.
 800-777-8570**

**ACCEPTED
 with COMMENTS
 In EPA Letter Dated:
 DEC 7 2005**

**Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under EPA Reg. No.
 499-503**

PRECAUTIONARY STATEMENTS

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-225-3320 for emergency medical treatment information.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically. Contains petroleum distillate - vomiting may cause aspiration pneumonia.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Apply this product only as specified on the label. Spray will damage plant foliage.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

SHAKE WELL BEFORE USING

See complete label for additional Directions for Use.

GENERAL USE RESTRICTIONS

- Do not apply to classrooms when in use.
- Use in aircraft cargo areas only. Do not use in aircraft cabins.
- Do not apply this product in hospital patient rooms while occupied or in any rooms occupied by the elderly or infirm.
- Do not use in patient rooms in hospitals and nursing homes for treatment of bed bugs.
- Do not apply to institutions (including libraries, sport facilities, etc.) in the immediate area when occupants are present.
- Do not spray on plastic, painted, or varnished surfaces.
- Do not spray directly into any electronic equipment such as radios, televisions, computers, etc.
- Keep children and pets out of treated areas until surfaces are dry.
- For food handling establishments, reoccupation may be made at 21 day intervals.

**HOW TO USE WHITMIRE MICRO-GEN
 CRACK AND CREVICE® INJECTION SYSTEM**

CRACK & CREVICE TREATMENT: Use Whitmire Micro-Gen TC 240 with the supplied actuator and injection tubes or Whitmire Micro-Gen System Equipment (such as System III) injected TC 240 into cracks

INDOOR AND OUTDOOR TREATMENTS

FOR ACTIVE AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames.

OUTDOOR TREATMENTS

SUBTERRANEAN ANTS SUCH AS ARGENTINE ANTS, FIRE ANTS, AND HARVESTER ANTS: To treat nests, inject the nest at 3-6 sites at a rate of 5-10 seconds per site. For very large nests (over 12" diameter), increase the number of injection sites. Space injection sites in a circular pattern on the nest surface with one site in the center. To quiet active ants on nest surface, sweep surface with spray from a distance of 12 inches. For best results, apply when temperature is 65-85°F, or in early morning or late evening hours. Treat new nests as they appear.

BEES, WASPS AND YELLOW JACKETS: Apply directly on insects and their nests from approximately 18 inches. Contacted insects will fly away. Applications should be made in late evening when insects are at rest. Disperse into hiding and breeding places contacting as many insects as possible.

PERIMETER TREATMENT: For crawling insects, apply as a Crack & Crevices Treatment or Spot Treatment into cracks and crevices where insects may enter premises or harbor. Areas to be treated include, but are not limited to, weep holes, under siding, wall voids, soffits, around attic vents, cracks and crevices, openings around windows, doors and pipes and landscape timbers. Treatment of infested fences and tree holes is permitted.

TERMITES AND CARPENTER ANTS LIVING INSIDE TREES: Determine the highest and lowest elevation of the colony. Drill holes into colony at highest elevation and inject 5 to 10 seconds. Do the same for lowest elevation. Drill intermediate holes every four feet between the upper and lower elevations if required.

AERIAL TERMITE CARTON IN TREES: Drill or punch hole into center of carton. Inject for 5 to 10 seconds. If carton is greater than 12 inches across, open another injection site. Inject for 5 to 10 seconds. Make one additional injection site for each 12 inches of carton.

TERMITES AND CARPENTER ANTS IN STUMPS, UTILITY POLES AND FENCES: Treat open holes or drill horizontal holes in a circular pattern where insects are suspected. Inject 2 to 5 seconds of product per hole.

**The purpose of such applications of this product for termites is to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but are merely a supplement. This product is not recommended as sole protection against termites. For active termite infestations, get a professional inspection.

TREATMENT FOR IN-GROUND SERVICE BOXES: Ants, Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Fire Ants, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders (such as Black Widow and Brown Recluse), and Springtails. Use spray actuator to treat open boxes and exposed surfaces. Do not contact any electrical wiring or mechanical parts of meter.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not puncture or incinerate! Empty container by using the product according to the label directions. Offer empty container for recycling, if available, or place in trash if allowed by state and local regulations. If container is partly filled, contact your local solid waste agency or call 1-800-CLEANUP for disposal instructions.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

DEC 7 2005

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

499-503

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
www.wmmg.com

© 2004 Whitmire Micro-Gen Research Laboratories, inc.

WHITMIRE MICRO-GEN

TC 240

INSECTICIDE

KILLS:

Ants, Bed Bugs, Bees, Booklice, Boxelder Bugs, Carpenter Ants, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Drug Store Beetles, Earwigs, Elmleaf Beetles, Fire Ants, Flies, Flour Beetles, Grain Weevils, Indian Meal Moths, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders, Springtails, Above-Ground Termites**, Ticks, Trogloderma (Cabinet, Khapra, and Warehouse Beetles), Wasps and Yellow Jackets.

** Refer to termite section for restrictions regarding these pests.

FOR USE IN AND AROUND:

Apartments, Campgrounds, Food Storage Areas, Homes, Hospitals*, Hotels, Meat Packing and Food Processing Plants, Motels, Nursing Homes*, Resorts, Restaurants and other Food Handling Establishments, Schools*, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes*), Utilities, Warehouses, and other Commercial and Industrial Buildings.

* Refer to "General Use Restrictions" section for restrictions regarding these sites.

ACTIVE INGREDIENT:

λ-Cyhalothrin	0.05%
Prallethrin	0.03%

OTHER INGREDIENTS:

Contains petroleum distillates.	99.92%
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See page 2 for additional Precautionary Statements.

EPA Reg. No. 499-LNG • EPA Est. No. 499-MO-1



**WHITMIRE MICRO-GEN
RESEARCH LABORATORIES INC.
800-777-8570**

4/5



TC 240 INSECTICIDE

PRECAUTIONARY STATEMENTS

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-225-3320 for emergency medical treatment information.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically. Contains petroleum distillate - vomiting may cause aspiration pneumonia.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Apply this product only as specified on the label. Spray will damage plant foliage.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

GENERAL USE RESTRICTIONS

- Do not apply to classrooms when in use.
- Use in aircraft cargo areas only. Do not use in aircraft cabins.
- Do not apply this product in hospital patient rooms while occupied or in any rooms occupied by the elderly or infirm.
- Do not use in patient rooms in hospitals and nursing homes for treatment of bed bugs.
- Do not apply to institutions (including libraries, sport facilities, etc.) in the immediate area when occupants are present.
- Do not spray on plastic, painted, or varnished surfaces.
- Do not spray directly into any electronic equipment such as radios, televisions, computers, etc.
- Keep children and pets out of treated areas until surfaces are dry.
- For food handling establishments, reapplication may be made at 21 day intervals.

HOW TO USE THE WHITMIRE MICRO-GEN CRACK AND CREVICE[®] INJECTION SYSTEM

CRACK & CREVICE TREATMENT: Use Whitmire Micro-Gen TC 240 with the supplied actuator and injection tubes or Whitmire Micro-Gen System Equipment (such as System III[®]). Inject TC 240 into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For closed voids calculate the void's cubic area and treat at the rate of one to five seconds per three cubic feet. Several holes may be required in long-running voids.

SPOT TREATMENT: Use Whitmire Micro-Gen TC 240 with the supplied actuator and injection tubes or Whitmire Micro-Gen System Equipment (such as System III). Hold container upright while spraying. Direct nozzle approximately 12 inches from surface to be treated. Apply directly on insects in these locations when possible.

INDOOR TREATMENTS

COCKROACHES, DERMESTIDS, SCORPIONS, SPIDERS, EARWIGS, CRICKETS, SOWBUGS, MILLIPEDES, CENTIPEDES: Apply as a Crack & Crevice treatment using the prescribed dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewer, floor drains and meter boxes. Spot Treatments may

also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

TICKS: Apply as a Crack & Crevice treatment or Spot Treatment into infested and egg laying sites near pet resting areas including, but not limited to cracks and crevices, behind baseboards, door and window frames. Do not use on pets. Also, treat your yard with a product registered for yard application and your pet with a product registered for pet treatment.

BOOKLICE & SILVERFISH: Apply as a Crack & Crevice treatment into cracks and crevices of baseboards, door frames, bookcases and other harbor-age sites. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

ANTS: Apply as a Crack & Crevice treatment into cracks, crevices and voids where ants may be traveling and/or harboring, including, but not limited to, around doors, window frames, wall voids and other structural cracks and crevices. Spot Treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, FLIES, CLUSTER FLIES AND CLOVER MITES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices in cabinets and pantries, wall voids, around window and door frames, under siding and in machinery. Spot treatments may also be made to areas including, but not limited to, storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas.

CHOCOLATE MOTHS, DRUGSTORE BEETLES, GRAIN WEEVILS, FLOUR BEETLES, INDIAN MEAL MOTHS, TROGODERMAS (IN EX-POSED ADULT AND LARVAL STAGES): Make Spot and/or Crack and Crevice treatment to localized areas including shelves, cupboards, walls and floors where food products suspected of being infested are stored. Remove or cover food before treatment. This treatment is supplemental to good housekeeping, since it will have no effect on insects within packaging. Infested food packaging should be destroyed.

BED BUGS: Take bed apart and treat cracks, joints and interior of framework. Treat mattresses and box springs; especially tufts, folds, and edges. Also treat picture frames, moldings and cracks and crevices in the room. Allow all treated articles to dry thoroughly before use.

APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS

Food Handling Establishments are places other than private residences in which food is held, processed, prepared or served.

In Federally inspected meat and poultry plants, the facility must not be in operation when this product is used, regardless of the method of application. In non-Federally inspected facilities, Spot and/or crack and crevice application may be used while the facility is in operation provided exposed food is covered or removed from the area being treated prior to application.

Do not apply directly to food. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food, food utensils or food processing surfaces. Food must be covered or removed in areas being treated. Spray should not be applied directly to surfaces or utensils that may come into contact with food. Food contact surfaces and equipment must be cleaned with an effective cleaning compound and rinsed with potable water before using.

Reapplication may be made at 21 day intervals. Limit individual spot treatments to an area no larger than 20% of the treated surface. Individual spot treatments should not exceed two square feet.

FOOD AREAS: Food areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) is also considered a food area.

Apply as a Crack and Crevice Treatment or Spot Treatment directly into cracks, crevices, voids or onto surfaces where insects are resting. Use an injection tube or other Whitmire Micro-Gen equipment capable of delivering a pinstream. Apply to different elements of construction; between equipment and floor; openings leading to hollow spaces in walls, equipment legs and bases; and where insects hide or rest.

FOOD SERVING AREAS (facilities where foods are served, such as dining rooms): Apply as a Crack and Crevice Treatment or Spot Treatment to selective surfaces such as baseboards, under elements of construction, and into crack and crevices. Avoid treating surfaces likely to be contacted by food. Food should be covered or removed in area being treated. Do not apply directly to food or allow applications to contaminate food.

NON-FOOD AREAS: Includes areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling).

Apply as a Crack & Crevice Treatment or Spot Treatment to selective surfaces such as baseboards, under elements of construction, into cracks and crevices. Avoid treating surfaces likely to be contacted by food.

5/5