499-528

12-20-2011



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

> > DEC 2 0 2011

Ms. Dana Thomas Whitmire Micro-Gen Research Laboratories Inc. 3568 Tree Court Industrial Blvd. St. Louis, MO 63122-6682

Dear Ms. Thomas:

Subject: Application for Pesticide Notification (PRN 98-10) Comply with Agency pyrethroid labeling letter dated June 4, 2009, Revise per PR Notice 2007-4, Add optional marketing claims, Revise EPA Est. No. and Warranty statement, Revise copyright year, Remove WMG logo, Add telephone number and manufactured for/by statements TC 251B EPA Reg. No. 499-528 Your submission dated July 27, 2011

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated July 27, 2011 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on <u>nonrefillable</u> containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions regarding this letter, please contact Dana Pilitt, Ph.D. of my staff at (703) 305-7071 or via e-mail at <u>pilitt.dana@epa.gov</u>.

Sincerely. Charl Ceb C

Richard Gebken Product Manager 10 Insecticide Branch Registration Division (7505P)

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060

€EPA	ection Agence	tion Agency		Registration Amendment Other		OPP Identifier Number		
	Appli	cation for Pe	sticide - Se	ction	I			
1. Company/Product Number	ır	2.	EPA Product M			3. Pr	roposed Classi	fication
499-528			Richard Gebken	ו		-1	None	Restricted
4. Company/Product (Name) TC 251B		PI	<b>PM#</b> 10					
3568 Tree Court Indus St. Louis, MO 63122-0	esearch Laboratories Inc. trial Blvd.	(b to E	Expedited Re (i)(i), my produce EPA Reg. No. Product Name	t is sim	ilar or idention	cal in co	mposition ar	
		Sectio	Contraction of the local division of the loc					
Amendment - Explain below.			Final prin Agency le "Me Too" Other - E	etter dat * Applica	ation.	to		
This label has been revised via 2007-1 (or non-notification), ar understand that it is a violation	8-10 and 2007-1 and revisions per a Notification of label changes per ad the requirements of EPA's regul of 18 U.S.C. Sec. 1001 to willfully 66, this product may be in violation	EPA letter dated June lations at 40 CFR 156 make any false state	e 4, 2009 along wit b. No other change ement to EPA. I fur be subject to enfor	th minor r s have be ther unde	evisions allowe een made to the rstand that if th	e labeling e amende	or the CSF for t	his product. I onsistent with
1. Material This Product Wil	Pa Backaged Inc	Section	n - III					
Child-Resistant Packaging Yes No * Certification must be submitted	Unit Packaging Yes No If "Yes" No. p Unit Packaging wgt. conta	er If "Yes"	No. pe		2. Type of C	Metal Plastic Glass Paper	Specify)	
3. Location of Net Contents	Information 4. Size	(s) Retail Container	<u>_</u>	5. Lo	cation of Labe	Directio		C C C
6. Manner in Which Label is	Affixed to Product	Lithograph Paper glued Stenciled	[_] Oti	her				¢
		Section	n - IV	111		Lecce.	C C C C C C C C C C C C C C C C C C C	C
1. Contact Point (Complete	items directly below for identi	fication of individua	d to be contacted	d, if nec	essary, to pro	ess this	application.)	
<sub>Name</sub> Dana Thomas		Title Product R	Title Product Regulatory Mgr		1	Telephone No. ((השושל Area Code) 636-861-4223		
	ments I have made on this for hy knowlinglly false or misleadi						6. Date Appl Received (Stan	
2. Signature Aara M Mondes Dana Thomas Digitaly signed by Data Themas Dana Thomas Digitaly signed by Data Themas Digitaly si		3. Title Produc	з. тіче Product Regulatory Mgr					
4. Түреd Name Dana M. Thomas		5. Date July 2	<sup>5. Dete</sup> July 27, 2011					

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

White - EPA File Copy (original)

## Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL			
EPA Registration #	Date Submitted to EPA	Electronic file name	
499-528	7/27/11	000499-00528.20110725v21.Pyrethroid Non-AgOutdr.pdf	

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

Signature

7/27/11 Date

Dana M. Thomas Name (typed)

Product Regulatory Mgr. Title



July 27, 2011

Document Processing Desk (NOTIF - PYRETHROID) Office of Pesticide Programs (7504P) U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington VA 22202-4501

# RE: Label Notification per PRNs 98-10 and 2007-1 and revisions per EPA letter dated 6/4/09 regarding Pyrethroid Non-Agricultural Outdoor Products TC 251B, EPA Reg. No. 499-528

Enclosed in support of this action is a completed application and two copies revised labeling; one highlighted, one clean (an electronic copy and label certification is also enclosed).

The label has been revised as follows:

- 1. Added optional marketing claims for SmartCap™ Technology
- 2. Changed EPA Est. # to 7969-MO-1
- 3. Revised the Environmental Hazards section as outlined in EPA letter dated 6/4/09
- 4. Moved the Use Restrictions section to above the General Information section
- 5. Added Use Restrictions as outlined in EPA letter dated 6/4/09
- 6. Revised Container Disposal (corrected >5 gal statement and container handling) per PRN 2007-1, 2007-4 and the LRM for products registered for uses other than household or residential
- 7. Added standard company warranty statement
- 8. Removed WMG logo, added "Manufactured for/by statements, added company telephone number for questions and revised the copyright year

This label has been revised via Notification of label changes per EPA letter dated June 4, 2009 along with minor revisions allowed via Notification per PRNs 98-10 and 2007-1 (or non-notification), and the requirements of EPA's regulations at 40 CFR part 156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR part 156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Sincerely,

Aana M Shemas

Dana M. Thomas Product Regulatory Mgr. & Authorized Agent for Whitmire Micro-Gen Research Laboratories, Inc. (636) 861-4223, dana.thomas@basf.com

CCCCC

occcc

BASF Corporation Pest Control Solutions 3568 Tree Court Industrial Blvd. St. Louis, MO 63122 Tel: (800) 777-8570 www.PestControl.basf.us

## Prescription Treatment<sup>®</sup> brand TC 251B Controlled Release Permethrin

[Alternate Brand Name: Prescription Treatment<sup>®</sup> brand Perma-Cap CS<sup>™</sup> {added via Notification 5/1/08} Descriptor: Controlled Release Permethrin]

#### **Optional Marketing Claims:**

- SmartCap<sup>™</sup> Technology
- With SmartCap<sup>™</sup> Technology
- Using SmartCap<sup>™</sup> Technology

#### KILLS:

Ants (foraging Carpenter, Fire and Harvester Ants; excluding Pharaoh ants), Bedbugs, Bees, Boxelder Bugs, Cadelles, Carpenter Bees, Centipedes, Chocolate Moths, Cigarette Beetles, Clover Mites, Cluster Flies, Cockroaches, Crickets, Darkling Beetles, Dermestids, Drugstore Beetles, Earwigs, Elm Leaf Beetles, Firebrats, Fleas, Flies, Flour Beetles, Grain Weevils, Hide Beetles, House Flies, Indianmeal Moths, Larder Beetles, Lesser Grain Borers, Lesser Mealworms, Mediterranean Flour Moths, Merchant Grain Beetles, Mealworms, Millipedes, Mosquitoes, Pill Bugs, Rice Weevils, Sawtoothed Grain Beetles, Scorpions, Silverfish, Sow Bugs, Spiders (excluding Black Widow and Brown Recluse), Springtails, Ticks, Warehouse Moths, Wasps, Webbing Clothes Moths, Wood Wasps, Wood-infesting Borers and Beetles

#### FOR USE IN AND AROUND:

Apartments, Calf Hutches, Calving Pens and Parlors, Campgrounds, Empty Chicken Houses, Dairy Areas, Dog Kennels, Hog Barns, Homes, Horse Barns, Hospitals<sup>†</sup>, Hotels, Milkrooms, Motels, Non-Food Areas of (Food Storage Areas, Grain Mills, Granaries, Meat Packing Plants, and Supermarkets), Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes<sup>†</sup>), Utilities, Warehouses, Commercial, Institutional<sup>†</sup> and Industrial Buildings

\* Refer to the Use Restrictions section for special instructions regarding the use of this product in these sites.

ACTIVE INGREDIENT:	
Permethrin	
OTHER INGREDIENTS:	
	TOTAL: 100.0%

## EPA Reg. No. 499-528 • EPA Est. No. 7969-MO-1 KEEP OUT OF REACH OF CHILDREN CAUTION

See page 2 for First Aid, Precautionary Statements and Directions for Use.

#### NET CONTENTS:

#### PRECAUTIONARY STATEMENTS

#### ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply this product or allow it to drift to crops or weeds on which bees are actively foraging. Additional information may be obtained from your Cooperative Extension Service.

This product is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors and anywhere such containers are present. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hr will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

Personal Protective Equipment (PPE): Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

NOTIFICATION

DEC 2 0 2011

#### Master Label

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.

#### DIRECTIONS FOR USE

#### IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANOR INCONSISTENT WITH ITS LABELING. USE RESTRICTIONS

- Do not apply this product through any type of irrigation system.
- Do not make outdoor applications during rain.
- Do not water the treated area to the point of run-off.
- All outdoor applications must be limited to spot or Crack & Crevice<sup>®</sup> treatments only, except for the following permitted uses:
  - 1. Treatment to soil or vegetation around structures;
  - 2. Applications to lawns, turf and other vegetation;
  - 3. Applications to building foundations, up to a maximum height of 3 ft.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors and eaves) are limited to spot and Crack & Crevice applications only.

- Do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies or aquatic habitat can occur. Except as directed by this label.
- Do not apply water-based sprays of this product to wall outlets, conduits, motor housing, junction and switch boxes or other electrical equipment where electrical short circuits might result because of possible shock hazard.
- Do not apply in aircraft cabins.
- Do not apply to humans, their clothing or bedding.
- Do not apply to firewood.
- Do not contaminate food or use on household pets. Cover fish tanks prior to application.
- Do not allow children or pets to contact treated surfaces until spray has dried.
- Do not apply in patient rooms while occupied or in any rooms while occupied by the elderly or infirm.
- Do not contaminate food, feedstuffs or water supply. Do not contaminate food preparation surfaces, kitchen utensils, dishes, or food storage containers. Cover any food/feed contact surfaces and cooking utensils, in the treatment area, during treatment or thoroughly clean before using.
- Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed (see General Information section for additional information).
- Do not apply to classrooms when in use.
- · Do not apply when occupants are present in the immediate area in institutions such as libraries, sport facilities, etc.

#### **GENERAL INFORMATION**

#### Kills pests on contact.

Apply this product only as specified on this label. This product is a flowable microencapsulated concentrate containing 2 lbs of permethrin per gallon. This product may be used on any surface which will not be damaged or stained by water. Heavy applications may leave a visible deposit. On some surfaces, this deposit can be removed with a damp cloth or sponge.

In situations where conventional application methods have not or are not likely to provide adequate coverage, foam generating equipment or similar machines can be used to provide a continuous coverage. Use drier foam inside wall voids to avoid any water damage or electrical short circuits.

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food/feed is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home,

#### Master Label

remove or cover all food/feed processing surfaces and exposed utensils, in the treatment area, prior to application or wash thoroughly before use; remove or cover any exposed food/feed prior to application. Non food/feed areas are areas such as garbage rooms, restrooms, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

#### MIXING INSTRUCTIONS

Thoroughly clean spray equipment before using this product. Shake concentrate well before diluting. When diluting, first add approximately 1/2 of the water to the spray tank and then add the proper amount of product. When emptying the bottles, triple rinse with water, shake well and add the rinsate to the spray tank. Add the rest of the water and agitate the sprayer thoroughly. Agitate sprayer occasionally during use to ensure even mixture. Shake or reagitate sprayer if spraying is interrupted or if dilution is left in sprayer overnight. If spray screens are used, they should be 50 mesh or larger.

**FOAM APPLICATION USE DIRECTIONS:** This product can be converted to a foam and the foam used to treat structural voids to kill insect and arthropod pests in walls, under slabs or in other void areas. Apply as a dry foam (approximately 25:1). Fill void with foam so that all surfaces are contacted. Mix end-use dilution with manufacturer's recommended volume of foaming agent. Use the recommended rate suspension converted to a foam to achieve the desired foam expansion ratio. Verify before treatment that the foaming agent is compatible with product.

#### SPRAY DILUTION CHART

Amount of	Spray Concentrate			
Finished Spray	0.5% Spray	1.0% Spray		
1 gal	2 2/3 fl oz	5 1/3 fl oz		

Use the higher rate for clean-outs and high insect infestations.

#### INDOOR TREATMENTS

Dilute product to 0.5% or 1.0% with water using the spray dilution chart.

Use as a Crack & Crevice product where insects may be hiding, living and breeding and/or for selected spot application.

Non-food/feed areas are areas such as garbage rooms, restrooms, floor drains (to sewers) entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

CENTIPEDES, COCKROACHES, CRICKETS, DERMESTIDS, EARWIGS, FIREBRATS, MILLIPEDES, PSOCIDS, SILVERFISH, SOWBUGS, SPIDERS (excluding Black Widow and Brown Recluse): Make spot treatment and/or Crack & Crevice injections into hiding places such as cracks and crevices, void areas, moist areas, openings around pipes and sinks and under refrigerators. Repeat as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

FLIES: Make spot applications of product on surfaces where flies rest in the dark at night or frequent during the day, such as window frames, screens, trash cans, ceilings and upper wall surfaces.

**CARPET BEETLES (Dermestids):** Treat along edges of carpet, under carpet edge, furniture, throw rugs and as a Crack & Crevice application in closets and other shelving where infested items are stored. Do not treat clothing with this product. Repeat application as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

**BED BUGS:** Take bed apart. Treat joints and channels if hollow, such as square or round tubing, and see that the interior framework is treated. Bed Bugs may also harbor in areas of the room away from the bed. Apply as a Crack & Crevice or spot treatment to night stands and other furniture in the room, wall voids, behind wall hangings, floor moldings, window casings, carpet edges, popcorn ceilings and other cracks & crevices. Apply this product until the surface is wet, but not to the point of runoff.

TICKS: Apply as a spot and/or Crack & Crevice treatment in infested and egg laying areas such as pet resting quarters, nearby cracks and crevices, behind baseboards, doors and window frames. Replace old pet bedding with clean, fresh bedding after treatment. Do not treat pets with this product. Repeat application as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

ANTS (foraging Carpenter, Fire and Harvester ants; excluding Pharaoh ants): Inject cracks, crevices and voids where ants may be traveling and/or harboring, including but not limited to, around doors, window frames, and other structural cracks, crevices and voids. Spot treatment of trails and feeding areas may also be made.

**STORED PRODUCT PESTS (exposed Adult and Immature stages):** Before storing products, treat warehouses, production facilities, storage areas, rail cars, truck beds and other areas where products are stored. Apply as a spot and/or Crack & Crevice treatment to cracks, crevices and surfaces where pests have been seen or may have harborage. This product may be used to treat grain storage facilities and other listed areas for stored product pests. Cleaning areas prior to using this product will enhance kill. Remove and destroy any infested foodstuffs.

#### INDOOR AND OUTDOOR TREATMENTS

Select one of the following dilutions, 0.5% or 1.0%, for insects listed below unless otherwise noted.

HIBERNATING STAGES OF BOXELDER BUGS, ELM LEAF BEETLES, CLUSTER FLIES AND CLOVER MITES: Inject cracks, crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices in weep holes, wall voids, around window and door frames and under siding. Treat wall area around entry points on exterior where insects congregate. Repeat application as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

**CARPENTER ANTS:** Make spot and/or Crack & Crevice applications to any trails, inside or outside around doors and windows, and other places where ants enter premises. Areas which may be treated include, but are not limited to, atriums, crawl spaces, ceilings, garages, soffits, wall voids, weep holes and other void areas. Where possible, treat ant nest or infested wood directly. Repeat application as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

BEES AND WASPS: As a preventative treatment to help prevent these insects from building nests or workers from congregating on structures, follow perimeter treatment directions under **Outdoor Treatments** section of label. Pay particular attention to soffit and eave areas. For treatment of nest inside wall voids, inject into the nest a sufficient quantity to treat the entire nest. More than one point of application may be required. Most applications should require no more than ½ gal of diluted material. Pay particular attention to what the water may contact, such as electrical wiring. Do not use where there is a shock hazard. Apply this product until the surface is wet, but not to the point of runoff.

THIS PRODUCT DOES NOT PROVIDE INSTANT KNOCKDOWN. The best time to treat is late in the day. The colony should be dead in 2 – 3 days. Depending on the situation, protective equipment for the applicator may be required (is advisable).

**MOSQUITOES:** Apply at the 0.5% rate. Apply directly to walls, ceilings, window screens, and other resting areas as a surface treatment. May be used inside residential buildings as well as in and around carports, garages, and storage sheds; also see **Outdoor Treatments**. Apply as a general structural perimeter spray to landscape plantings, turf, and building foundations (up to a maximum height of 3 ft) to treat against mosquitoes. Yards or other frequented areas enclosed by landscaping can benefit from the creation of a mosquito barrier to reduce invading mosquitoes by the treatment of perimeter vegetation. Higher volumes applied result in better coverage and, as a rule, will improve kill. Application to vegetation away from structures may require additional certification, e.g. in turf or ornamental categories. Consult your state regulatory agency for requirements.

WOOD DESTROYING INSECTS: Localized treatment of wood-infesting insects – Beetles and Borers (such as the following families: Anobiidae, Bostrichidae, Buprestidae, Cerambycidae, Lyctidae, Dedemerldae, Sesiidae): Locate suspected infestation and/or insect tunnels and cavities. Drill pilot holes into insect tunnels, gallery and inaccessible closed voids. Drill holes 8 – 10" apart to ensure thorough coverage. Inject 5 – 10 sec/hole. Do not contact electrical wiring, plumbing, etc. Use the 1.0% rate.

BROADCAST TREATMENT OF WOOD FOR USE AGAINST WOOD-INFESTING INSECTS AND NUISANCE PESTS OUTSIDE OF STRUCTURE: Apply a 0.5% emulsion with a fan spray using a maximum of 25 psi. Apply just to the point of runoff. To treat against wood-infesting insects active inside trees, utility poles and/or fence posts, drill to find the interior infested cavity and inject a 0.5% emulsion. To treat against bees apply in late evening when insects are at rest. Aim spray at nest openings in ground, bushes and in cracks and crevices which may harbor nests, saturating nest openings and contacting as a many insects as possible.

#### OUTDOOR TREATMENTS

KILLS: Ants (foraging Carpenter, Fire and Harvester ants; excluding Pharaoh ants), Boxelder Bugs, Carpenter Bees, Carpet Beetles, Centipedes, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Earwigs, Elm Leaf Beetles, Flies, Millipedes, Mosquitoes, Psocids, Scorpions, Silverfish, Sowbugs, Spiders (excluding Black Widow and Brown Recluse) and Ticks

Not for use in Outdoor Residential Misting Systems.

Follow the instructions for outdoor applications under Use Restrictions. Apply to and around buildings, porches, window frames, eaves, patios, garages, garbage dumps and other areas where pests congregate or enter premises. Repeat treatment as necessary to maintain effectiveness, but not more than twice per week. Treatments must be at least 3 days apart. Use the 0.5% rate.

This product can be applied against pests in outdoor void areas such as meter boxes and manholes. Repeat treatment as necessary, but not more than twice per week. Treatments must be at least 3 days apart. Use the 0.5% rate.

**MOSQUITOES:** To treat against adult mosquitoes, use hand held, back-pack or truck mounted sprayers only. Not for use in wide area local, state or federal control programs. Not for use with truck mounted cold aerosol ULV sprayers, hand held cold aerosol ULV sprayers or thermal fogging devices. Do not apply by aircraft. Do not apply through any kind of irrigation and/or lawn sprinkler systems. Use up to 150 gal of diluted spray per acre. Treat shrubbery, vegetation and other areas where mosquitoes may rest. Shrubbery and vegetation around stagnant pools, marshy areas, ponds and shorelines may be treated if applied as a wet spray. Direct application of this product to any body of water is prohibited.

#### TREATMENT OF OUTSIDE SURFACES AND AROUND BUILDINGS

Follow the instructions for outdoor applications under Use Restrictions. Apply using a 0.5% emulsion to outside surfaces of buildings including, but not limited to, exterior siding, foundation, porches, window frames, eaves, patios, garages, refuse dumps, lawn areas adjacent or around private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, fence lines, storage sheds, barns, other residential structures, commercial, industrial and institutional buildings, soil, trunks of woody ornamentals and other areas where pests congregate or have been seen. Repeat treatment as necessary to maintain effectiveness.

**PERIMETER TREATMENT:** To help prevent infestations of buildings, treat a band of soil 5 - 10 ft wide around and adjacent to the building. Also treat the building foundation to a height of 2 - 3 ft where pests are active and may find entrance. Treat other insect entry points such as doors, windows, under siding and eaves of structure. May be applied in compressed air, back-pack or power spray equipment. An application rate of  $2 \ 2/3$  fl oz/1,000 ft<sup>2</sup> is desirable. Apply in a sufficient amount of water to adequately cover the area being treated, but avoiding dripping or run-off. Alternate directions are to use 10 - 20 fl oz of product per 50 gal of water.

DILUTION CHART					
	Tank Size (gallons)	Suggested Amount of Product per tank			
	15	4 – 8 fl oz			
	25	7 – 14 fl oz			
	50	10 – 20 fl oz			
	100	20 – 40 fl oz			

FLIES: Treat outside surfaces of screens, doors, window frames or wherever flies enter. Also treat surfaces on porches, in garages and other places where insects light or congregate. Repeat as necessary. Treat normal fly resting areas such as ceilings, upper corners of the area, around windows and doors, and other surfaces. Treat inside and outside surfaces of dumpsters and other trash holding containers. Treat trash storage areas. Treat so as to lightly moisten the target surface. For best results, begin treatments at the start of the fly season and repeat during periods of heavy fly activity. Treat no more than twice per week and at least 3 days apart.

CRAWLING INSECTS – including Ants (foraging Carpenter, Fire and Harvester ants; excluding Pharaoh ants), Cockroaches, Crickets, Earwigs, Sowbugs and Scorpions: For treatment of infestations of these insects in outdoor areas which are landscaped, planted, mulched or unmaintained. For ants, treat mounds and runways. Repeat application as infestation warrants and as reinfestation occurs. Apply to cracks and crevices where insects may enter premises. Areas to be treated include, but are not limited to, weep holes, under siding, wall voids, soffits, around attic vents, cracks and crevices, openings around windows, doors and pipes. Treatment of infested fences and tree holes is permitted.

FLEAS: Treat infested areas with the 0.5% rate.

FOAM APPLICATION USE DIRECTIONS: See instructions on page x.

PESTS UNDER SLABS: Infestations of arthropods, such as Ants (foraging Carpenter, Fire, and Harvester ants; excluding Pharaoh ants), Cockroaches and Scorpions inhabiting the area under slabs may be killed by drilling and injecting or horizontal rodding. Using a 0.5% – 1.0% emulsion, inject 1 gal/10 ft<sup>2</sup> or 2 gal/10 ft.

**CRAWL SPACES:** Broadcast product at 0.5% to all surfaces in crawl space to treat against Ants (foraging Carpenter, Fire, and Harvester ants; excluding Pharaoh ants), Fleas, Roaches, Scorpions or other arthropods. Product may also be applied under the structure through insecticidal delivery systems such as piping or flexible tubing mounted under the structure. Freat surfaces thoroughly but do not allow runoff to occur. Keep children and pets off surface until dry.

#### USE IN AND AROUND LIVESTOCK HOUSING

MAY BE USED IN AND ON OUTSIDE SURFACES OF BUILDINGS (Barns, Dairies, Feedlots, Livestock Houses, Foultry Houses, Stables and Swine Houses): Do not apply to lactating or food animals. Do not contaminate milk; food or drinking water. Remove all animals prior to making application to housing. Do not apply or allow insecticide to drift onto animals. Do not allow animals to enter treated areas until surfaces are dry.

**HOUSE FLIES AND STABLE FLIES:** For residential treatment of these pests on outdoor surfaces and in animal areas such as dog kennels, hog barns, horse barns, empty chicken houses, rabbit hutches, dairy areas, milk rooms, calf hutches, calving pens and parlors: Treat normal fly resting areas such as ceilings, screens, upper corners of the areas, around windows and doors, light fixtures and other surfaces known to attract flies. Apply as a coarse, wet spray. Using a 0.5% or 1.0% emulsion, apply at the rate of 1 gal/1,000 ft<sup>2</sup> of surface. For best results, initiate treatments at the beginning of the fly season and repeat during periods of heavy fly activity. Treat no more than twice per week and at least 3 days apart.

DARKLING BEETLES: Apply 2 2/3 - 5 1/3 fl oz concentrate/1,000 ft<sup>2</sup>. Use adequate water to ensure distribution.

**PERIMETER TREATMENT:** To treat against other pests and to help prevent infestations of buildings, treat a band of soil 6 – 10 ft wide around and adjacent to the building. Also treat the building foundation and sides where pests are active and may find

#### Master Label

e e e e

C C C C C C

entrance. Treat other insect entry points such as doors, windows, under siding and eaves of structure. Product may be applied in compressed air, backpack or power spray equipment. An application rate of 1.0 fl oz product/1,000 ft<sup>2</sup> is desirable. Apply in a sufficient amount of water to adequately cover the area being treated, but avoiding dripping or run-off.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides or fertilizers. Do not store above 122°F for extended periods of time. Preferred storage is above 32°F. Storage below 32°F may result in solidification. If warmed to above 32°F, product will return to original form. Freezing does not adversely affect this product. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Non-refillable container. Empty container by using the product according to the label directions. Do not reuse or refill this container! Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

[(<5.0 gal) Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure 2 more times.]

[(>5.0 gal) Empty remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times.

Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### CONDITIONS OF SALE AND WARRANTY

Follow the **Directions for Use**. It is impossible to eliminate all risks inherently associated with use of this product, and therefore all such risk shall be assumed by the Buyer. Whitmire warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use**, subject to the inherent risks, referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW: (A) WHITMIRE MAKES NO OTHER WARRANTIES EXPRESS OR IMPLIED, INCLUDING WARRANTIES OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY, (B) BUYER'S EXCLUSIVE REMEDY AND WHITMIRE'S AND SELLER'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT, AND (C) WHITMIRE AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, INCIDENTAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. Whitmire and the Seller offer this product, and the Buyer accepts it, subject to these **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of Whitmire.

Manufactured for: Whitmire Micro-Gen Research Laboratories, Inc. by BASF Corporation 3568 Tree Court Industrial Blvd. St. Louis, MO 63122-6682 Questions? Call 1-800-777-8570

© 2011 Whitmire Micro-Gen Research Laboratories, Inc.