


12-14-67

454



Net Weight 1-lb.

U. S. D. A. Reg. No 505-4

KILLS RATS AND MICE

WITH NEW AND IMPROVED

COWLEY'S Rat & Mouse Bait

contains *PROlin.*

most notable advance in rodenticides
since warfarin

Sure-death for rats and mice! Does away with dirty traps!

U. S. PATENTS 2,427,578 AND 3,113,071 OTHER PATENT APPLICATIONS PENDING

CAUTION Keep Out Of Reach Of Children

Protect children, pets and domestic animals from bait. If swallowed by humans, animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and upon repeated feedings, causes fatal hemorrhage in rodents. It also contains an antibacterial agent, sulfaquinoxaline, which inhibits bacteria that produce vitamin K, an antidote for warfarin, thereby increasing effectiveness against hard-to-kill rats.

ACTIVE INGREDIENTS: 3-(Alpha-Acetylbenzyl)-4-hydroxycoumarin (warfarin) 0.025 %
N1-2Quinoxalinylsulfanilamide (sulfaquinoxaline) 0.025 %
INERT INGREDIENTS 99.950 %

Prolin (warfarin plus sulfaquinoxaline) is manufactured under license from the Wisconsin Alumni Research Foundation, U. S. Patent No. 2,427,578 and 3,113,071, other patent applications pending.

The antibacterial agent, Sulfaquinoxaline, increases the effectiveness of this bait and assures more thorough control of rats and mice. Especially recommended for difficult control infestations.

DIRECTIONS FOR USE

MICE: Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8-12 feet are recommended. Assure uninterrupted supply of bait for a period of not less than 12 days.

RATS: Place 4 to 6 ounces of bait in dry containers next to burrows, in runways or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped.

Replace stale, damp and moldy baits with fresh baits.

Manufactured by
S. L. COWLEY & SONS MFG. CO., Inc.
Hugo, Oklahoma 74743