

LINCK'S LIQUID KILZ-ALL.

APPLICATION RATES TO USE WITH ANY KIND OF SPRAY EQUIPMENT, INCLUDING WATERING CAN, TANK SPRAYER, HOSE SYPHONER.

Liquid Measure of Linck's Liquid Kilz-All	Add water to make	Apply to
2½ ozs.	1 gal. water	250 sq. ft.
5 ozs.	2 gals. water	500 sq. ft.
10 ozs.	4 gals. water	1000 sq. ft.

Wash all spray equipment thoroughly after use to avoid corrosion of metal parts.

GENERAL INFORMATION: Linck's Liquid Kilz-All is useful for general post-emergent weed control. It is non-selective in its action. Its phytotoxic properties are quickly inactivated on contact with soil. It is a combination of herbicide and surfactant. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young actively growing weeds. It produces top-kill only, so repeat applications are required for season long weed control of perennials.

MIXING INSTRUCTIONS: Linck's Liquid Kilz-All must be thoroughly dissolved. Fill the spray equipment reservoir about half full with water and add the required amount of herbicide with agitation. Finish filling the reservoir with water and apply.

ACCEPTED

4-1-68

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER NO. 506-112
TO ATTACHED COMMENTS.

**COVERS
3200
Sq. Ft.**

USDA REG.
NO. 506-112



LINCK'S
Liquid Kilz-All

ACTIVE INGREDIENTS:
Sodium Cacodylate 22.6%
Dimethylarsinic Acid (Cacodylic Acid) 3.9%
INERT INGREDIENTS 73.5%
Total Arsenic (as elemental) all in water soluble form 12.7%
Product contains 2.48 lbs. cacodylic acid equivalent per gallon.

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS.

**NOW CONTAINS
A SURFACTANT**

CONTENTS ONE U S LIQUID QU.

DIRECTIONS FOR USE

LAWN RENOVATION — Mow lawn to be renovated about 1 inch high before treatment. Follow dilution rates on side panel. Cover foliage thoroughly, preferably on a warm sunny day. Foliage will usually turn brown in about five days. Bermudagrass and other deep-rooted perennials, or uncovered foliage, may require a second treatment. Reapply five days later if green areas remain.

When foliage is all brown, matted areas should be raked and bare areas scarified. About five days after last treatment, water thoroughly and then seed at rate recommended for new lawns. Seed must be raked into soil. Keep seeded areas well watered.

KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL AFTER FIRST RAIN OR SPRINKLING FOLLOWING TREATMENT. DO NOT TRACK FROM TREATED TO UNTREATED AREAS.

NON-CROP — Linck's Liquid Kilz-All is useful for general weed control on drainage ditchbanks and rights-of-way, along sidewalks, driveways and fences, around buildings and ornamentals, and on similar non-crop areas. Mix 1 to 2 quarts of Linck's Liquid Kilz-All in 25 gallons of water. Spray unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs reapply as required.

DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION, OR BY ANIMALS, WILDLIFE, AND AQUATIC LIFE, OR FOR IRRIGATION PURPOSES. DO NOT GRAZE TREATED AREAS TO LIVESTOCK.

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into soil.

Do not reuse empty container. Destroy by burying in a safe place.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

NOTICE: Use of this material in a manner, or at any time, other than in accordance with the directions, may produce plant injury or other undesirable results. The Linck Company makes no express or implied warranties other than specified on the label.

MANUFACTURED BY
O. E. LINCK DIVISION — WALCO-LINCK CORPORATION
Clifton, New Jersey
MADE AND PRINTED IN U.S.A.