

APR 11, 1973
5/3-55
COMM-F-12

FRUIT TREE AND VEGETABLE SPRAY

32° BAUME WHEN PACKED AT 60° F.

ACTIVE INGREDIENTS:

Calcium Polysulfide 30.0%

Calcium Thiosulfate 1.5%

INERT INGREDIENTS: 68.5%

100.0%

DIRECTIONS

For dormant spraying of fruit trees,
(mite and aphid clean-up)
12 gallons to 100 gallons of water

For delayed dormant or pink spray,
(for mites and aphids)
7½ gallons liquid lime-sulphur
¾ pint Black Leaf 40
100 gallons of water

For psyllids of potatoes,
2½ gallons liquid lime-sulphur
100 gallons of water

For psyllids of tomatoes,
1 gallon liquid lime-sulphur
4 pounds wettable sulphur
80 gallons of water

NOTE: Do not apply lime-sulphur when trees are in bloom, and do not apply after bloom within two weeks of any oil spray. Do not apply during periods of high temperature, since sulphur sprays may burn foliage under such conditions. Use with care near buildings since discoloration of painted surfaces may result.

Follow Directions Carefully—Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of

use, storage or handling of this material not in strict accordance with directions given herewith.

**DO NOT REUSE EMPTY
DRUM**

Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

**KEEP OUT OF THE REACH
OF CHILDREN**

Net 54 Gallons

Directions for Dipping Cattle and Sheep Against Psoroptic and Sarcoptic Scab

In dipping cattle and sheep, use 1 gallon Sterling Liquid Lime-Sulphur to each 15 gallons of water.

Add Sterling Liquid Lime-Sulphur to offset loss from steam condensing through natural causes.

The repeat treatment time intervals should be 7 days and visibly affected animals should be held in the vat 2 to 3 minutes and their heads submerged at least once. Clean dipping vats thoroughly after each dipping.

- CAUTION**
- Do not dip sick or debilitated animals.
 - Do not dip in cold or stormy weather.
 - Do not dip animals having fresh wounds.

DIRECTIONS FOR USE

(a) Change the dipping vat solution as soon as it becomes excessively contaminated regardless of the number of animals dipped. (b) Do not use dipping solutions more than 10 days old. (c) Unless information is present to assure yourselves that heating is not necessary, the temperature of the dip should be maintained in the range of 95° to 105° F.

NOTE: Does not require additional wetting agents. Contains special wetting agent—your sprays will go farther and wet the trees and foliage better.

CAUTION!!

AVOID CONTACT WITH SKIN OR EYES

**Avoid long continued or frequently repeated contact—
in case of contact, wash thoroughly with plenty of
water.**

KEEP FROM FREEZING

EPA Reg. No. 513-55

ACCEPTED WITH COMMENTS