

Revised

44-15

121

BEST AVAILABLE COPY

DIRECTIONS FOR USE

Vapona insecticide may be used for control of face flies, stable and houseflies, mosquitoes, roaches, ants, gnats, crickets and certain other small flying insects.

FACE FLY SOLUTION: Mix 8 fl. oz. with 1 gallon water and 1 lb. sugar.
1% SOLUTION: Mix 8 fl. oz. with 1 gallon water.
0.5% SOLUTION: Mix 4 fl. oz. with 1 gallon water.
FOGGING SOLUTION: Mix 8 fl. oz. with 1 gallon petroleum kerosene.

DO NOT MIX UP MORE DILUTED SPRAY THAN IS NEEDED FOR THE APPLICATION

DILUTED SOLUTION

FACE FLIES—Face Fly Solution with sugar. Apply 2 fluid ounces of face fly diluted solution with sugar to face and neck of beef or dairy cattle each morning for about two weeks and thereafter as needed. Mix the diluted solution well and shake the sprayer occasionally during the application. Use a sprayer with adequate capacity.

LACTATING DAIRY & BEEF CATTLE, HORSES 1%—To protect cattle and horses from houseflies, stable flies, horn flies, mosquitoes, and gnats, use the 1% diluted solution daily and cover the animal with spray with particular attention to the back, flank and legs. Daily spray should exceed 2 ounces per head. Do not soak the skin. For best results, spray 1 ounce morning and evening.

HOGS, SHEEP AND GOATS—1%—Apply 1% diluted solution daily to face, back, flanks, belly and legs. Do not use more than 1 ounce per head daily. Reduce dosage for younger animals.

FOGGING AND SURFACE SPRAY ANIMAL BUILDINGS, DAIRY BARN, MILK ROOMS, STABLES, HOG HOUSES, SHELTERS AND POULTRY HOUSES 1%—For control of Flies, Mosquitoes, Roaches, Ants, Gnats and Crickets. Use 1 pint of 1% diluted fogging solution per each 8000 cu. ft. Repeat as necessary. Always shut doors and be enclosed if they have been opened for ventilation with 18 hours. Do not spray directly on to the animals. Always use 1 pint of 1% diluted fogging solution per 1000 square feet of weekly, or less, fogging. Cattle, window sills, and doors. **DO NOT CONTAMINATE MILK, WATER, FEED OR UTENSILS.**

FLY BREEDING AREAS, MANURE PILES, CARPARK DUMPS 1%—Use 1 pint of 1% diluted fogging solution per each 8000 cu. ft. Treat every 4 to 5 days. Do not spray directly on to the animals.

FEED LOTS, BEEF & HOG STOCKYARDS FOR CONTROL OF FLIES, MOSQUITOES, GNATS—Use 1 pint of 1% diluted fogging solution per each 8000 cu. ft. Treat every 4 to 5 days. Do not spray directly on to the animals.

MILK RANCHES 0.5%—Use 1 pint of 0.5% diluted solution per 1000 square feet. Treat every 4 to 5 days. Do not spray directly on to the animals.

MINK RANCHES 0.5%—Use 1 pint of 0.5% diluted solution per 1000 square feet. Treat every 4 to 5 days. Do not spray directly on to the animals.


DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.



VAPORA™

18.5% VAPONA CONCENTRATE

For use on lactating Dairy cattle, Beef cattle, Hogs, Sheep and Goats.



Kills: Face flies, Houseflies, Stable flies, Horn flies, Gnats, Mosquitoes, Ants, Roaches

ACTIVE INGREDIENTS	18.5%
2,2-Dichlorovinyl dimethyl phosphate	18.5%
Related compounds	0.5%
Xylene range aromatic solvent	78.5%
INERT INGREDIENTS	81.5%
Total	100.000%

Prepared by Roberts Vapona Insecticide Company, a Division of Roberts Laboratories, Inc., 2000 Roberts Road, Roberts, Kansas 66560.

DANGER

KEEP OUT OF REACH OF CHILDREN

(See additional information on side panel)

May be fatal if swallowed, inhaled or absorbed through skin or eyes. Vapona insecticide concentrate and the diluted emulsion are poisonous, therefore, read warnings and labels and antitoxin state nearby prominently on the side panel before use.

NET CONTENTS ONE GALLON (128 fl. oz.)



FOR AGRICULTURAL USE ONLY - OR - ONLY FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS THE TERM SERVICE PERSON INCLUDES BUT IS NOT LIMITED TO, A JANITOR, PEST CONTROL OPERATOR, MAINTENANCE PERSON, LAWN CARE AND LANDSCAPING PERSONNEL

USE PRIOR TO:

PRECAUTIONS IN USING

Do not use in areas where children play. Store the concentrate in a tightly closed container. It must not be able to reach children. Do not use in ventilated rooms. Wash all equipment that has been used in spraying with water before use. Do not use in areas where children play.

Do not use in areas where children play. Do not use in areas where children play. Do not use in areas where children play.

DANGER—Do not get in eyes. Vapona insecticide is poisonous. It will be absorbed through skin or eyes. Do not get on clothes. Wash all contaminated clothing with soap and hot water before reuse. Do not reuse spray bottles. Do not use in areas where children play. Do not use in areas where children play. Do not use in areas where children play.

This product is for use on animals. Do not use on humans. Do not use on humans. Do not use on humans.

ANTIDOTE—If swallowed, give milk or water. If in eyes, wash with water. If on skin, wash with soap and water. **CALL A PHYSICIAN IMMEDIATELY.**

NOTICE OF WARRANTY

ROBERTS LABORATORIES, INC. warrants that the product is free from defects in material and workmanship at the time of shipment.

ROBERTS LABORATORIES, INC.

UIN

BEST AVAILABLE COPY