



**ROBERTS**

**MALATHION 57%**

**PREMIUM GRADE**


Emulsifiable Concentrate to be used as a grain protectant and for the control of Lice, Ticks, Hornflies on Beef Cattle; Lice on Hogs. Lice, Mites, Ticks on Poultry. For use as a fly Bait. Flies and Fleas on Dogs and Cats. Certain Household and Garden Insects. Can be used in Back Rubber and Stock Oilers.



# MALATHION 57%

## PREMIUM GRADE

Emulsifiable Concentrate to be used as a grain protectant and for the control of Lice, Ticks, Hornflies on Beef Cattle; Lice on Hogs. Also, Mites, Ticks on Poultry. For use as a fly Bait. Flies and Fleas on Dogs and Cats. Certain Household and Garden Insects. Can be used in Back Rubber and Stock Oilers.



**Residual Spray - Boxcars for Loading and Transporting Bagged Flour and Packaged Cereals:**

For control of the exposed stages of confused flour beetle, rice weevil, granary weevil, saw-toothed grain beetle, flat grain beetle, red flour beetle, rusty grain beetle, lesser grain borer, Indian meal moth and mite pest infesting empty boxcars into which bagged flour and packaged cereals are to be loaded and transported, apply Malathion Grain Protectant No. 57 as follows:

#### **DIRECTIONS**

1. Clean the boxcar thoroughly, then remove and burn all sweepings and debris.
2. Spray walls and floor to the point of runoff with either 1 gallon of Malathion Grain Protectant No. 57 in 25 gallons of water, or 1 gallon of this product in 19 gallons of deodorized kerosene. (The deodorized kerosene solution should be made up of 4 parts Deosol or other aromatic hydrocarbon-type solvent.)
3. Let the sprayed boxcar stand empty with the doors open until the spray has thoroughly dried.

#### **FLOWERS, SHRUBS AND SHADE TREES:**

Use at the rate of 1 pint per 25 gallons of water (2 teaspoonfuls) per gallon. Mix well and spray plants thoroughly. This spray will kill aphids, red spider mites, bagworms, leafhoppers, thrips, scale, mealybugs, white flies, leaf miners, leaf bugs, adult Japanese beetles, lace bugs, plant bugs and tent caterpillars. Repeat treatment as necessary. Injury may occur to certain tender plants such as certain ferns and some species of crassula. Do not treat these.

**FRUIT TREES:** Use at the rate of 2 teaspoonfuls per gallon of water. Mix well and spray plants thoroughly. Repeat as necessary. This spray will kill the following:

**ON PEACHES:** Spider mites, plum curculio, Oriental fruit moth. Do not apply later than 7 days before harvest.


**ON PEARS:** Codling moth, pear payila, plum curculio, leaf roller. Do not apply later than 3 days before harvest. May cause injury in Bosc pear under certain conditions in the Northwest.

DE TREES.  
5 gall  
n. M  
is spr  
warm  
white  
panels  
tent  
p  
re

te  
Mix w  
s  
g  
un  
ly

Es  
ly  
to

25 40



**ROBERTS**

**MALATHION**

**57%**