

CAUTION

Do not apply in areas where animals have received direct application of DDT within 4 hours.

Do not contaminate water or feed stuffs.

Avoid contact with skin. Avoid breathing spray mist. Harmful if swallowed, inhaled or absorbed through the skin.

If swallowed induce vomiting immediately and get medical attention. ATROPINE IS ANTIDOTAL.

Wash thoroughly with soap and water after handling and before eating or smoking.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

Do not reuse empty drum. Return drums to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

Do not spray or store near fire or flame. Do not contaminate milk or milk equipment.

NOTICE OF WARRANTY

ROBERTS LABORATORIES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED, concerning this product or its use, which extends beyond the statements on this label.

Manufactured by
ROBERTS LABORATORIES, INC.
Rockford, Illinois



'FLY-DIE'

LIVESTOCK SPRAY WITH VAPONA

ACTIVE INGREDIENTS	BY WEIGHT
2,2-Dichlorovinyl dimethyl phosphate460%
Related Compounds040%
Piperonyl Butoxide* (Technical)250%
Pyrethrins025%
Mineral Oil	49.057%
Petroleum Derivative Solvents	50.118%
INERT INGREDIENTS050%
Total	100.000%

*Equivalent to (Butylcarbityl) 6-Propylpiperonyl) Ethers 0.20% and related compounds 0.05%

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

DIRECTIONS FOR USE

FOR DAIRY CATTLE, BEEF CATTLE AND HORSES: To protect livestock from face flies, house flies, stable flies, horse flies, mosquitos and gnats, spray the animal daily using one to two ounces per head. Cover the animal with spray with particular attention to the back, flanks, legs and shoulders. Do not use in excess of 2 ounces per adult animal. Do not soak the skin or wet the hides.

FOR BARN, MILK ROOMS AND STABLES: When used as a space sprayer direct power sprayer toward upper portion of the enclosure until area is filled with a fine mist. If a coarse spray is used apply wherever flies congregate or alight, until surfaces are moistened. Avoid contact with food or milk utensils. Close doors and windows before treatment when possible. This treatment will kill resting adult flies. Do not remain in treated areas and ventilate the area after treatment is complete.



'FLY-DIE'

TRADE MARK

**LIVESTOCK SPRAY
WITH VAPONA**

Front Panel

ROBERTS F1

KILLS FLIES
OTHER SMALL
W.

HOMES, CABINETS
A

ACTIVE INGREDIENTS :
2,2-dichlorovinyl d
Related compounds .

INERT INGREDIENTS . .

KEEP OUT OF REACH OF CHILDREN
See Back Panel for Directions
Contains Pesticides

* Equivalent to 20.0%
**Reg. T.M. Shell Chemical Co.

Net Weight 3.7 ounces

ROBERTS

RO

Back Panel

FLYCON INSECT BAR

Controls flies, mosquitoes, gnats and other small flying insects.

Use in homes, motels, dairy barns, animal shelters and garbage cans.

DIRECTIONS FOR USE

HOMES

MOTELS

To control flies, mosquitoes, gnats and other small flying insects, place one Flygon Bar per average size room, (10 x 10 x 10 feet) or per 100 square feet of floor area. Normally this is effective for three months. Replace strips if effectiveness diminishes.

For control of flies in garbage cans attach $\frac{1}{2}$ Flygon Bar inside the lid of an average size garbage can (20 gallon capacity). One unit will be effective for the entire fly season. Keep lid on garbage can.

ANIMAL BUILDINGS

To control flies, mosquitoes, gnats and other small flying insects in horse barns, dairy barns, calf barns, stables, pig parlors, dog kennels, poultry houses and other farm buildings, suspend one Flygon Bar (10" strip 3.7 oz.) for each 100 square feet of floor space. If more than one strip is required, distribute them equally over area to be treated. Weather and other conditions such as strong drafts may alter the performance, but this treatment is usually effective for one season. Replace the strips if effectiveness diminishes.

Back Panel (cont'd.)

NOTICE

WARNING: Do not use in kitchens, restaurants, or areas where food is prepared or served. Do not place in hospital or clinic rooms such as patient rooms, wards, nurseries, operating or emergency areas. Do not use in any rooms where infants, the sick, or aged are or will be present for any extended period of confinement. Do not get in mouth, harmful if swallowed. After prolonged storage, a small amount of liquid may form on strip. Do not get liquid in eyes. Wash hands thoroughly with soap and water after handling strip. Do not contaminate food, water, food containers or cooking utensils. Keep out of reach of children.

NOTICE OF WARRANTY

ROBERTS LABORATORIES INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE STATEMENTS ON THIS LABEL.

Use of the packaged compositions is covered by Shell's U.S. Patent 2,956,073 and by pending patent applications. Sales of these compositions carries with it no license to use such compositions for any purpose other than insecticide generators.

Distributed by:

ROBERTS LABORATORIES

ROCKFORD, ILLINOIS 61101

EPA REG. NO. 523-45